



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:08 PM GMT

PDB ID : 4V02
Title : MinC:MinD cell division protein complex, Aquifex aeolicus
Authors : Ghosal, D.; Lowe, J.
Deposited on : 2014-09-10
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	<div> <div>76%</div> <div>16%</div> <div>• 5%</div> </div>
1	B	258	<div> <div>78%</div> <div>13%</div> <div>• 6%</div> </div>
2	C	128	<div> <div>2%</div> <div>70%</div> <div>21%</div> <div>• 5%</div> </div>
2	D	128	<div> <div>2%</div> <div>70%</div> <div>18%</div> <div>5%</div> <div>7%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MG	A	700	-	-	-	X
3	MG	B	700	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5666 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SITE-DETERMINING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	244	Total	C	N	O	S	0	0	0
			1891	1212	319	355	5			
1	B	242	Total	C	N	O	S	0	0	0
			1874	1203	317	349	5			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	252	SER	-	EXPRESSION TAG	UNP O67033
A	253	HIS	-	EXPRESSION TAG	UNP O67033
A	254	HIS	-	EXPRESSION TAG	UNP O67033
A	255	HIS	-	EXPRESSION TAG	UNP O67033
A	256	HIS	-	EXPRESSION TAG	UNP O67033
A	257	HIS	-	EXPRESSION TAG	UNP O67033
A	258	HIS	-	EXPRESSION TAG	UNP O67033
A	40	ALA	ASP	ENGINEERED MUTATION	UNP O67033
B	252	SER	-	EXPRESSION TAG	UNP O67033
B	253	HIS	-	EXPRESSION TAG	UNP O67033
B	254	HIS	-	EXPRESSION TAG	UNP O67033
B	255	HIS	-	EXPRESSION TAG	UNP O67033
B	256	HIS	-	EXPRESSION TAG	UNP O67033
B	257	HIS	-	EXPRESSION TAG	UNP O67033
B	258	HIS	-	EXPRESSION TAG	UNP O67033
B	40	ALA	ASP	ENGINEERED MUTATION	UNP O67033

- Molecule 2 is a protein called PROBABLE SEPTUM SITE-DETERMINING PROTEIN MINC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	122	Total	C	N	O	S	0	0	0
			929	587	170	169	3			
2	D	119	Total	C	N	O	S	0	0	0
			908	575	166	164	3			


There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	202	GLY	-	EXPRESSION TAG	UNP O67034
C	203	SER	-	EXPRESSION TAG	UNP O67034
C	204	HIS	-	EXPRESSION TAG	UNP O67034
C	205	HIS	-	EXPRESSION TAG	UNP O67034
C	206	HIS	-	EXPRESSION TAG	UNP O67034
C	207	HIS	-	EXPRESSION TAG	UNP O67034
C	208	HIS	-	EXPRESSION TAG	UNP O67034
C	209	HIS	-	EXPRESSION TAG	UNP O67034
D	202	GLY	-	EXPRESSION TAG	UNP O67034
D	203	SER	-	EXPRESSION TAG	UNP O67034
D	204	HIS	-	EXPRESSION TAG	UNP O67034
D	205	HIS	-	EXPRESSION TAG	UNP O67034
D	206	HIS	-	EXPRESSION TAG	UNP O67034
D	207	HIS	-	EXPRESSION TAG	UNP O67034
D	208	HIS	-	EXPRESSION TAG	UNP O67034
D	209	HIS	-	EXPRESSION TAG	UNP O67034

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Mg 1 1	0	0
3	A	1	Total Mg 1 1	0	0

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 31	C 10	N 5	O 13	P 3	0	0
4	B	1	Total 31	C 10	N 5	O 13	P 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: SITE-DETERMINING PROTEIN




• Molecule 1: SITE-DETERMINING PROTEIN




• Molecule 2: PROBABLE SEPTUM SITE-DETERMINING PROTEIN MINC



• Molecule 2: PROBABLE SEPTUM SITE-DETERMINING PROTEIN MINC





4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	130.62Å 130.62Å 298.51Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.17 – 2.70 49.75 – 2.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.17-2.70) 100.0 (49.75-2.70)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.89 (at 2.69Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.188 , 0.236 0.206 , 0.205	Depositor DCC
R_{free} test set	2133 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	53.1	Xtriage
Anisotropy	0.128	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 42196 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5666	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.79	0/1914	0.83	1/2586 (0.0%)
1	B	0.68	0/1897	0.77	1/2563 (0.0%)
2	C	0.77	0/939	0.81	0/1261
2	D	0.66	0/916	0.80	0/1227
All	All	0.73	0/5666	0.80	2/7637 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	79	ARG	NE-CZ-NH1	6.95	123.77	120.30
1	B	201	ILE	CB-CA-C	-5.30	101.01	111.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1891	0	1998	25	0
1	B	1874	0	1987	18	0
2	C	929	0	985	22	0
2	D	908	0	966	22	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
4	A	31	0	12	0	0
4	B	31	0	12	0	0
All	All	5666	0	5960	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:140:ASP:OD1	2:D:142:SER:HB3	1.86	0.76
1:A:176:ILE:HG13	1:A:186:MET:CE	2.18	0.73
1:A:176:ILE:HG13	1:A:186:MET:HE1	1.70	0.72
2:D:140:ASP:OD1	2:D:142:SER:N	2.24	0.71
2:C:177:VAL:HG22	2:C:190:ILE:HD11	1.74	0.68
1:B:10:GLY:HA2	1:B:157:VAL:HG21	1.75	0.67
1:A:79:ARG:H	1:A:80:GLY:HA2	1.62	0.64
2:C:151:GLU:N	2:C:152:PRO:CD	2.60	0.64
2:C:140:ASP:OD2	2:C:142:SER:HB3	2.00	0.61
1:B:171:VAL:HG22	1:B:200:ILE:HD13	1.83	0.61
1:A:176:ILE:HA	1:A:186:MET:HE1	1.83	0.60
2:C:151:GLU:N	2:C:152:PRO:HD3	2.18	0.59
2:C:118:LEU:CD2	2:C:134:LYS:HD2	2.33	0.58
1:B:105:THR:O	1:B:109:ILE:HG13	2.05	0.56
1:B:79:ARG:H	1:B:80:GLY:HA2	1.70	0.56
2:D:140:ASP:OD1	2:D:142:SER:CB	2.53	0.56
2:C:148:LEU:HD21	2:D:148:LEU:HD21	1.88	0.55
1:B:6:VAL:HG11	1:B:134:VAL:HG22	1.89	0.55
1:A:137:ALA:O	1:A:166:LYS:NZ	2.39	0.54
2:C:148:LEU:CD2	2:D:148:LEU:HD21	2.37	0.54
2:C:148:LEU:HD21	2:D:148:LEU:CD2	2.38	0.53
1:A:169:TYR:C	1:A:170:LYS:HG2	2.28	0.53
1:A:46:LEU:HD22	1:A:50:LEU:HD22	1.90	0.53
2:C:146:VAL:HG22	2:C:177:VAL:HG13	1.90	0.53
2:D:138:ILE:HG23	2:D:138:ILE:O	2.07	0.53
1:A:79:ARG:N	1:A:80:GLY:HA2	2.24	0.53
1:B:96:VAL:HG12	1:B:97:ILE:N	2.24	0.52
2:C:177:VAL:HG22	2:C:190:ILE:CD1	2.40	0.52
2:D:151:GLU:N	2:D:152:PRO:CD	2.73	0.52
2:D:191:GLU:OE2	2:D:195:ARG:NH1	2.42	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:LEU:HD11	1:A:65:GLU:OE2	2.10	0.52
2:D:118:LEU:CD2	2:D:134:LYS:HD2	2.40	0.51
2:D:151:GLU:N	2:D:152:PRO:HD3	2.25	0.51
2:C:174:TYR:HB2	2:C:175:PRO:HD2	1.93	0.51
1:A:36:LEU:CD2	1:A:119:LEU:HB2	2.41	0.50
1:A:17:THR:OG1	1:A:121:ASP:OD2	2.19	0.50
1:B:141:LEU:HD22	1:B:201:ILE:HD11	1.92	0.50
2:D:134:LYS:HG2	2:D:156:GLN:HB3	1.93	0.49
2:C:118:LEU:HD22	2:C:134:LYS:HD2	1.93	0.49
1:A:189:VAL:HG23	2:C:200:LEU:HD11	1.93	0.49
2:C:200:LEU:O	2:C:200:LEU:HD22	2.13	0.49
1:A:98:ASP:HB3	1:A:101:LYS:HD2	1.95	0.48
1:A:83:LEU:HD23	1:A:83:LEU:C	2.34	0.48
2:C:138:ILE:O	2:C:138:ILE:HG23	2.12	0.48
1:A:42:GLY:O	1:A:44:ARG:HG3	2.14	0.47
1:B:181:VAL:HG23	1:B:186:MET:HE2	1.95	0.47
1:A:17:THR:HG22	1:A:49:ILE:HD12	1.97	0.47
2:D:140:ASP:OD1	2:D:142:SER:CA	2.63	0.47
1:A:171:VAL:HG22	1:A:200:ILE:HD13	1.97	0.46
2:C:113:LYS:O	2:C:114:ASP:HB2	2.15	0.46
1:B:210:LEU:HD12	1:B:220:ILE:HG21	1.96	0.46
2:D:102:ARG:NH1	2:D:138:ILE:HD11	2.31	0.45
2:D:154:LEU:HD12	2:D:163:ILE:HG22	1.98	0.45
1:A:8:THR:C	1:A:16:LYS:HD3	2.37	0.45
2:D:146:VAL:HG22	2:D:177:VAL:HG13	1.98	0.45
1:A:105:THR:O	1:A:109:ILE:HG13	2.16	0.45
1:A:8:THR:OG1	1:A:9:SER:N	2.50	0.44
2:D:140:ASP:CG	2:D:142:SER:HB3	2.38	0.44
1:A:43:LEU:HD12	1:A:43:LEU:HA	1.69	0.44
1:B:8:THR:HA	1:B:16:LYS:HD2	2.00	0.43
1:B:79:ARG:N	1:B:80:GLY:HA2	2.32	0.43
2:C:169:ARG:HA	2:C:169:ARG:HD3	1.74	0.43
1:A:49:ILE:HG12	1:A:214:THR:HG23	2.00	0.43
2:C:93:LEU:HD22	2:C:97:GLN:HB2	2.00	0.43
2:C:175:PRO:HB2	2:C:190:ILE:HG13	2.01	0.43
1:B:125:GLY:HA3	1:B:157:VAL:HG22	2.00	0.43
1:A:135:SER:N	1:A:136:PRO:HD2	2.34	0.43
2:C:175:PRO:HB2	2:C:190:ILE:HB	2.01	0.43
2:C:134:LYS:HG2	2:C:156:GLN:HB3	2.01	0.42
1:B:96:VAL:HG12	1:B:97:ILE:H	1.82	0.42
1:A:6:VAL:HG23	1:A:120:VAL:HB	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:102:ARG:CZ	2:D:138:ILE:HD11	2.50	0.42
1:A:36:LEU:HD12	1:A:50:LEU:HD21	2.01	0.41
1:B:169:TYR:C	1:B:170:LYS:HG2	2.40	0.41
1:B:42:GLY:O	1:B:44:ARG:HG3	2.20	0.41
1:B:176:ILE:CG1	1:B:186:MET:HE1	2.51	0.41
2:C:98:ARG:NH1	2:C:116:GLU:OE2	2.53	0.41
2:D:120:GLY:O	2:D:136:GLY:HA2	2.21	0.41
2:D:118:LEU:HD22	2:D:134:LYS:HD2	2.03	0.40
1:B:46:LEU:HD22	1:B:50:LEU:HD22	2.03	0.40
1:B:109:ILE:O	1:B:112:SER:HB3	2.21	0.40
2:D:105:ILE:HD12	2:D:117:VAL:HG12	2.03	0.40
2:D:93:LEU:HD23	2:D:93:LEU:HA	1.95	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	240/258 (93%)	231 (96%)	8 (3%)	1 (0%)	39	69
1	B	238/258 (92%)	228 (96%)	9 (4%)	1 (0%)	39	69
2	C	120/128 (94%)	113 (94%)	7 (6%)	0	100	100
2	D	115/128 (90%)	110 (96%)	5 (4%)	0	100	100
All	All	713/772 (92%)	682 (96%)	29 (4%)	2 (0%)	46	75

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	41	ILE
1	B	41	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	207/220 (94%)	192 (93%)	15 (7%)	18	41
1	B	205/220 (93%)	188 (92%)	17 (8%)	14	31
2	C	97/103 (94%)	82 (84%)	15 (16%)	3	8
2	D	94/103 (91%)	78 (83%)	16 (17%)	2	6
All	All	603/646 (93%)	540 (90%)	63 (10%)	9	20

All (63) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	GLU
1	A	6	VAL
1	A	43	LEU
1	A	46	LEU
1	A	50	LEU
1	A	68	VAL
1	A	100	GLU
1	A	128	LYS
1	A	147	GLU
1	A	148	VAL
1	A	152	ARG
1	A	170	LYS
1	A	183	ARG
1	A	221	VAL
1	A	244	ILE
1	B	3	GLU
1	B	6	VAL
1	B	43	LEU
1	B	46	LEU
1	B	48	MET
1	B	50	LEU
1	B	53	GLU
1	B	55	ARG
1	B	67	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	112	SER
1	B	148	VAL
1	B	149	SER
1	B	152	ARG
1	B	170	LYS
1	B	171	VAL
1	B	183	ARG
1	B	201	ILE
2	C	85	ARG
2	C	86	LEU
2	C	87	LEU
2	C	93	LEU
2	C	102	ARG
2	C	106	LEU
2	C	129	LEU
2	C	138	ILE
2	C	140	ASP
2	C	148	LEU
2	C	160	LYS
2	C	181	GLU
2	C	191	GLU
2	C	195	ARG
2	C	200	LEU
2	D	85	ARG
2	D	86	LEU
2	D	87	LEU
2	D	93	LEU
2	D	102	ARG
2	D	106	LEU
2	D	129	LEU
2	D	138	ILE
2	D	148	LEU
2	D	160	LYS
2	D	177	VAL
2	D	181	GLU
2	D	191	GLU
2	D	195	ARG
2	D	200	LEU
2	D	201	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	ATP	A	800	3	24,33,33	1.02	1 (4%)	31,52,52	1.98	6 (19%)
4	ATP	B	800	3	24,33,33	1.00	1 (4%)	31,52,52	1.69	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	A	800	3	-	0/18/38/38	0/3/3/3
4	ATP	B	800	3	-	0/18/38/38	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	800	ATP	C5-C4	2.44	1.46	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	800	ATP	C5-C4	2.69	1.46	1.40

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	800	ATP	N3-C2-N1	-6.96	123.56	128.89
4	B	800	ATP	N3-C2-N1	-5.09	125.00	128.89
4	A	800	ATP	PA-O3A-PB	-4.03	121.42	132.73
4	B	800	ATP	PA-O3A-PB	-3.28	123.51	132.73
4	B	800	ATP	C2'-C1'-N9	-2.75	110.09	114.29
4	B	800	ATP	C4-C5-N7	-2.64	107.05	109.48
4	A	800	ATP	C1'-N9-C4	-2.19	123.64	126.94
4	B	800	ATP	PB-O3B-PG	-2.13	125.53	132.67
4	A	800	ATP	C2-N1-C6	2.01	122.37	118.77
4	A	800	ATP	N6-C6-N1	2.25	124.04	119.20
4	A	800	ATP	O3G-PG-O2G	3.75	121.66	107.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	244/258 (94%)	-0.44	1 (0%) 93 94	24, 41, 67, 127	0
1	B	242/258 (93%)	-0.33	0 100 100	31, 48, 73, 105	0
2	C	122/128 (95%)	-0.18	3 (2%) 61 61	30, 44, 92, 124	0
2	D	119/128 (92%)	-0.31	3 (2%) 61 61	29, 47, 91, 111	0
All	All	727/772 (94%)	-0.34	7 (0%) 84 85	24, 46, 77, 127	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	139	GLY	5.0
2	C	141	HIS	4.8
2	C	170	ASN	4.7
2	D	141	HIS	3.8
2	D	84	SER	2.7
1	A	182	LYS	2.2
2	D	205	HIS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	MG	A	700	1/1	0.82	0.35	16.52	34,34,34,34	0
3	MG	B	700	1/1	0.94	0.33	12.09	43,43,43,43	0
4	ATP	B	800	31/31	0.98	0.16	0.71	25,37,43,45	0
4	ATP	A	800	31/31	0.98	0.14	0.21	18,33,38,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.