wwPDB X-ray Structure Validation Summary Report (i)

Feb 1, 2016 – 01:54 PM GMT

PDB ID	:	3VEK
Title	:	Both Zn Fingers of GATA1 Bound to Palindromic DNA Recognition Site, P1 Crystal Form
Authors	:	Jacques, D.A.; Ripin, N.; Wilkinson-White, L.E.; Guss, J.M.; Matthews, J.M.
Deposited on	:	2012-01-09
Resolution	:	2.63 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3377 (2.68-2.60)
Clashscore	102246	3781 (2.68-2.60)
Ramachandran outliers	100387	3722 (2.68-2.60)
Sidechain outliers	100360	3722 (2.68-2.60)
RSRZ outliers	91569	3388 (2.68-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	20	95% 5%
1	D	20	90% 10%
2	B	20	90% 10%
2	E	20	90% 10%
3	C	119	8% 79% • 19%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	119	8% 80% 19%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3154 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*TP*TP*GP*TP*CP*TP*TP*AP*TP*CP*AP*GP*AP*TP*GP*GP*AP*CP*TP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	20	405	196	68	122	19	0	0	0
1	D	20	405	196	68	122	19	0	0	0

- Molecule 2 is a DNA chain called DNA (5'-D(*AP*AP*GP*AP*GP*TP*CP*CP*AP*TP*CP*TP*GP*AP*TP*AP*AP*GP*AP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	20	409	196	80	114	19	0	0	0
2	E	20	409	196	80	114	19	0	0	0

- Molecule 3 is a protein called Erythroid transcription factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	96	756	456	155	135	10	0	0	0
3	F	96	756	456	155	135	10	0	0	0

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
4	D	2	2	2	0	0
4	E	1	1	1	0	0
4	B	1	1	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
4	C	3	3	3	0	0
			Total	Zn		
4	A	2	2	2	0	0
			Total	Zn		
4	F	5	5	5	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA (5'-D(*TP*TP*GP*TP*CP*TP*TP*AP*TP*CP*AP*GP*AP*TP*GP*GP*AP*CP*TP*C)-3')

Chain A:



• Molecule 1: DNA (5'-D(*TP*TP*GP*TP*CP*TP*TP*AP*TP*CP*AP*GP*AP*TP*GP*GP*AP*CP*TP*C)-3')

Chain D:



• Molecule 2: DNA (5'-D(*AP*AP*GP*AP*GP*TP*CP*CP*AP*TP*CP*TP*GP*AP*TP*AP*AP*GP*AP*C)-3')

Chain B:



• Molecule 2: DNA (5'-D(*AP*AP*GP*AP*GP*TP*CP*CP*AP*TP*CP*TP*GP*AP*TP*AP*AP*GP*AP*C)-3')

Chain E:



• Molecule 3: Erythroid transcription factor

Chain C:



• Molecule 3: Erythroid transcription factor

Chain F:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	35.20Å 62.41Å 66.05Å 72.39° 74.57° 73.63°	Depositor
Resolution (Å)	48.68 – 2.63 48.68 – 2.63	Depositor EDS
% Data completeness (in resolution range)	96.3 (48.68-2.63) 88.4 (48.68-2.63)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.14 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.235 , 0.275 0.233 , 0.280	Depositor DCC
R_{free} test set	738 reflections (5.37%)	DCC
Wilson B-factor (Å^2)	51.4	Xtriage
Anisotropy	0.877	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.25 , 11.7	EDS
Estimated twinning fraction	0.456 for h,h-k,h-l	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Outliers	0 of 14469 reflections	Xtriage
F_o,F_c correlation	0.92	EDS
Total number of atoms	3154	wwPDB-VP
Average B, all atoms (Å^2)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.03% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.58	0/452	1.20	1/696 (0.1%)
1	D	0.58	0/452	1.21	2/696 (0.3%)
2	B	0.58	0/460	1.17	2/708 (0.3%)
2	E	0.57	0/460	1.14	2/708 (0.3%)
3	C	0.28	0/769	0.47	0/1038
3	F	0.28	0/769	0.47	0/1038
All	All	0.47	0/3362	0.94	7/4884 (0.1%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	D	18	DC	O4'-C1'-N1	6.01	112.21	108.00
2	E	38	DG	O4'-C1'-N9	5.65	111.95	108.00
1	A	18	DC	O4'-C1'-N1	5.42	111.79	108.00
1	D	18	DC	O4'-C4'-C3'	-5.41	102.34	104.50
2	B	38	DG	O4'-C1'-N9	5.39	111.77	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	405	0	230	0	0
1	D	405	0	230	1	0
2	B	409	0	226	0	0
2	E	409	0	226	0	0
3	C	756	0	736	1	0
3	F	756	0	736	1	0
4	A	2	0	0	0	0
4	B	1	0	0	0	0
4	C	3	0	0	0	0
4	D	2	0	0	0	0
4	E	1	0	0	0	0
4	F	5	0	0	0	0
All	All	3154	0	2384	2	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (2) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:257:GLN:HB3	3:C:264:THR:HG22	1.93	0.51
1:D:6:DT:H3'	3:F:297:MET:HE1	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	92/119 (77%)	88 (96%)	4 (4%)	0	100	100
3	F	92/119 (77%)	88 (96%)	4 (4%)	0	100	100
All	All	184/238 (77%)	176 (96%)	8 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	82/100 (82%)	82 (100%)	0	100	100
3	F	82/100 (82%)	82 (100%)	0	100	100
All	All	164/200 (82%)	164 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
3	C	238	ASN
3	C	292	ASN
3	F	238	ASN
3	F	292	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 14 ligands modelled in this entry, 14 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	20/20 (100%)	-0.36	0	100	100	39, 60, 107, 111	0
1	D	20/20 (100%)	-0.51	0	100	100	40, 59, 112, 118	0
2	B	20/20 (100%)	-0.45	0	100	100	46, 55, 109, 114	0
2	E	20/20 (100%)	-0.53	0	100	100	49, 54, 110, 116	0
3	C	96/119 (80%)	0.48	10 (10%)	8	5	32, 61, 87, 92	0
3	F	96/119 (80%)	0.51	9 (9%)	11	7	35, 61, 84, 89	0
All	All	272/318 (85%)	0.21	19 (6%)	19	14	32, 61, 101, 118	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	310	SER	6.4
3	C	310	SER	6.4
3	F	274	SER	4.2
3	C	309	ALA	3.8
3	C	274	SER	3.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å^2)	Q<0.9
4	ZN	F	402	1/1	0.99	0.06	-1.97	47,47,47,47	0
4	ZN	F	401	1/1	0.99	0.08	-2.41	51,51,51,51	0
4	ZN	C	402	1/1	0.99	0.06	-3.21	42,42,42,42	0
4	ZN	C	401	1/1	0.99	0.08	-3.25	53,53,53,53	0
4	ZN	B	101	1/1	0.99	0.11	-	62,62,62,62	0
4	ZN	D	101	1/1	0.97	0.05	-	59,59,59,59	0
4	ZN	C	403	1/1	0.86	0.06	-	90,90,90,90	0
4	ZN	A	102	1/1	0.91	0.06	-	92,92,92,92	0
4	ZN	F	405	1/1	0.90	0.06	-	89,89,89,89	0
4	ZN	F	404	1/1	0.98	0.11	-	61,61,61,61	0
4	ZN	A	101	1/1	0.98	0.04	-	59,59,59,59	0
4	ZN	D	102	1/1	0.93	0.06	-	93,93,93,93	0
4	ZN	E	101	1/1	0.98	0.11	-	69,69,69,69	0
4	ZN	F	403	1/1	0.97	0.10	-	60,60,60,60	0

6.5 Other polymers (i)

There are no such residues in this entry.