



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:00 PM GMT

PDB ID : 3VRZ
Title : Crystal structure of HCK complexed with a pyrrolo-pyrimidine inhibitor 1-[4-(4-amino-7-cyclopentyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)phenyl]-3-benzylurea
Authors : Kuratani, M.; Tomabechei, Y.; Handa, N.; Yokoyama, S.
Deposited on : 2012-04-21
Resolution : 2.22 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4405 (2.24-2.20)
Clashscore	102246	5146 (2.24-2.20)
Ramachandran outliers	100387	5065 (2.24-2.20)
Sidechain outliers	100360	5066 (2.24-2.20)
RSRZ outliers	91569	4414 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	454	
1	B	454	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CA	A	601	-	-	-	X
2	CA	B	601	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7258 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein kinase HCK.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	430	Total	C	N	O	P	S	0	0	0
			3468	2217	583	647	1	20			
1	B	431	Total	C	N	O	P	S	0	0	0
			3479	2226	584	648	1	20			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	78	GLY	-	EXPRESSION TAG	UNP P08631
A	79	ALA	-	EXPRESSION TAG	UNP P08631
A	80	MET	-	EXPRESSION TAG	UNP P08631
A	81	GLY	-	EXPRESSION TAG	UNP P08631
A	82	SER	-	EXPRESSION TAG	UNP P08631
A	83	GLY	-	EXPRESSION TAG	UNP P08631
A	84	ILE	-	EXPRESSION TAG	UNP P08631
A	85	ARG	-	EXPRESSION TAG	UNP P08631
A	528	GLU	GLN	ENGINEERED MUTATION	UNP P08631
A	529	GLU	GLN	ENGINEERED MUTATION	UNP P08631
A	530	ILE	GLN	ENGINEERED MUTATION	UNP P08631
B	78	GLY	-	EXPRESSION TAG	UNP P08631
B	79	ALA	-	EXPRESSION TAG	UNP P08631
B	80	MET	-	EXPRESSION TAG	UNP P08631
B	81	GLY	-	EXPRESSION TAG	UNP P08631
B	82	SER	-	EXPRESSION TAG	UNP P08631
B	83	GLY	-	EXPRESSION TAG	UNP P08631
B	84	ILE	-	EXPRESSION TAG	UNP P08631
B	85	ARG	-	EXPRESSION TAG	UNP P08631
B	528	GLU	GLN	ENGINEERED MUTATION	UNP P08631
B	529	GLU	GLN	ENGINEERED MUTATION	UNP P08631
B	530	ILE	GLN	ENGINEERED MUTATION	UNP P08631


- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Ca 1 1	0	0
2	A	1	Total Ca 1 1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Cl 1 1	0	0
3	A	1	Total Cl 1 1	0	0

- Molecule 4 is 1-[4-(4-AMINO-7-CYCLOPENTYL-7H-PYRROLO[2,3-D]PYRIMIDIN-5-YL)PHENYL]-3-BENZYLUREA (three-letter code: VRZ) (formula: C₂₅H₂₆N₆O).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O 32 25 6 1	0	0
4	B	1	Total C N O 32 25 6 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	97	Total O 97 97	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	146	Total	O	0	0
			146	146		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tyrosine-protein kinase HCK



• Molecule 1: Tyrosine-protein kinase HCK



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.64 Å 96.49 Å 180.86 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.14 – 2.22 49.14 – 2.22	Depositor EDS
% Data completeness (in resolution range)	91.9 (49.14-2.22) 92.0 (49.14-2.22)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.40 (at 2.22 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.239 , 0.267 0.234 , 0.264	Depositor DCC
R_{free} test set	2000 reflections (3.48%)	DCC
Wilson B-factor (Å ²)	38.3	Xtriage
Anisotropy	0.661	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	3 of 59493 reflections (0.005%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7258	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.97 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.6065e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: VRZ, CA, PTR, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/3533	0.48	1/4767 (0.0%)
1	B	0.28	0/3545	0.47	0/4784
All	All	0.28	0/7078	0.48	1/9551 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	227	LEU	CA-CB-CG	5.64	128.26	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3468	0	3430	24	0
1	B	3479	0	3438	31	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	32	0	26	2	0
4	B	32	0	26	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	97	0	0	1	0
5	B	146	0	0	3	0
All	All	7258	0	6920	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:385:ARG:HD2	1:A:408:ALA:HB1	1.59	0.83
1:B:304:VAL:HG13	1:B:334:ILE:HD11	1.69	0.75
1:B:440:THR:HG23	1:B:442:LYS:H	1.60	0.66
1:B:223:THR:H	1:B:226:GLU:HG3	1.62	0.64
1:B:94:TYR:CE1	1:B:253:LYS:HE3	2.36	0.60
1:A:319:HIS:HB3	1:A:322:LEU:HG	1.83	0.60
1:B:163:LEU:O	1:B:190:ARG:NH1	2.36	0.59
1:B:432:GLU:OE2	1:B:506:ARG:NH1	2.31	0.58
1:B:401:LYS:NZ	5:B:835:HOH:O	2.30	0.57
1:B:180:THR:OG1	1:B:204:LYS:NZ	2.37	0.55
1:B:97:ILE:HG12	1:B:253:LYS:HE2	1.91	0.53
1:B:243:SER:OG	1:B:244:VAL:N	2.38	0.52
1:B:424:PHE:HB3	1:B:425:PRO:HD3	1.92	0.52
1:B:293:ALA:HB3	1:B:338:THR:HG22	1.92	0.52
1:A:264:ARG:NH2	1:A:331:LYS:O	2.40	0.52
1:B:253:LYS:NZ	5:B:726:HOH:O	2.42	0.51
1:B:172:SER:HA	1:B:244:VAL:O	2.11	0.51
1:B:440:THR:HG23	1:B:442:LYS:N	2.25	0.50
1:B:302:MET:HG3	1:B:307:PHE:HD1	1.76	0.50
1:B:221:PHE:HB3	1:B:226:GLU:HB2	1.93	0.50
1:B:302:MET:HG3	1:B:307:PHE:CD1	2.47	0.50
1:B:221:PHE:HZ	1:B:230:HIS:CD2	2.30	0.50
1:B:362:LYS:NZ	5:B:815:HOH:O	2.45	0.49
1:A:181:THR:OG1	1:A:204:LYS:NZ	2.45	0.49
1:A:172:SER:HA	1:A:244:VAL:O	2.12	0.49
1:A:430:ALA:HB1	1:A:432:GLU:OE2	2.13	0.48
1:A:428:TRP:NE1	1:A:454:GLU:OE1	2.40	0.48
1:A:456:VAL:HG11	1:A:487:CYS:HB2	1.96	0.48
1:A:484:PRO:HG2	1:A:487:CYS:HB2	1.96	0.47
1:B:222:SER:OG	1:B:226:GLU:OE2	2.32	0.47
1:A:307:PHE:CZ	4:A:603:VRZ:H23	2.50	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:309:ALA:O	1:B:313:VAL:HG23	2.16	0.46
1:B:168:ASN:HB2	1:B:190:ARG:NH1	2.31	0.46
1:A:314:MET:HG3	4:A:603:VRZ:H21	1.99	0.46
1:B:452:LEU:HD23	1:B:495:MET:HG2	1.99	0.45
1:A:468:ASN:O	1:A:472:ILE:HG13	2.17	0.45
1:A:530:ILE:HA	1:A:531:PRO:HD2	1.63	0.44
1:A:384:HIS:O	1:A:385:ARG:HB3	2.18	0.44
1:A:225:GLN:NE2	1:A:229:ASP:OD2	2.50	0.44
1:B:114:GLU:HB2	1:B:121:LYS:HB3	2.00	0.44
1:A:232:LYS:HB3	1:A:232:LYS:HE2	1.84	0.43
1:A:530:ILE:HD12	1:A:531:PRO:HD2	2.00	0.43
1:B:461:ILE:HG13	1:B:462:PRO:HD2	2.01	0.43
1:B:494:ILE:HD11	1:B:515:VAL:HG11	2.02	0.42
1:B:502:ARG:HA	1:B:503:PRO:HD3	1.92	0.42
1:A:474:ALA:HA	1:A:477:ARG:HD2	2.02	0.42
1:B:383:ILE:HD11	1:B:441:ILE:N	2.34	0.41
1:A:279:GLY:HA3	1:A:296:THR:O	2.20	0.41
1:A:88:VAL:HG23	1:A:110:MET:HG3	2.02	0.41
1:A:154:ILE:HA	1:A:154:ILE:HD12	1.87	0.41
1:A:149:TRP:HZ2	1:A:225:GLN:HG2	1.85	0.41
1:A:324:LYS:NZ	5:A:789:HOH:O	2.54	0.41
1:B:223:THR:OG1	1:B:226:GLU:HG2	2.21	0.41
1:A:334:ILE:H	1:A:334:ILE:HG12	1.74	0.40
1:B:157:LYS:HA	1:B:157:LYS:HD2	1.89	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	423/454 (93%)	408 (96%)	15 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	424/454 (93%)	412 (97%)	12 (3%)	0	100	100
All	All	847/908 (93%)	820 (97%)	27 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	374/393 (95%)	354 (95%)	20 (5%)	28	31
1	B	375/393 (95%)	354 (94%)	21 (6%)	26	29
All	All	749/786 (95%)	708 (94%)	41 (6%)	27	29

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	MET
1	A	154	ILE
1	A	193	ASP
1	A	195	ARG
1	A	206	ARG
1	A	222	SER
1	A	225	GLN
1	A	227	LEU
1	A	233	LYS
1	A	302	MET
1	A	304	VAL
1	A	305	GLU
1	A	313	VAL
1	A	320	ASP
1	A	325	LEU
1	A	331	LYS
1	A	432	GLU
1	A	461	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	476	GLU
1	A	485	GLU
1	B	116	SER
1	B	140	ARG
1	B	154	ILE
1	B	169	MET
1	B	196	GLN
1	B	198	ASP
1	B	218	ARG
1	B	253	LYS
1	B	271	LYS
1	B	305	GLU
1	B	307	PHE
1	B	324	LYS
1	B	325	LEU
1	B	328	VAL
1	B	338	THR
1	B	404	ASP
1	B	432	GLU
1	B	456	VAL
1	B	485	GLU
1	B	489	GLU
1	B	501	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PTR	A	527	1,2	14,16,17	1.14	1 (7%)	18,22,24	0.84	1 (5%)
1	PTR	B	527	1,2	14,16,17	1.18	1 (7%)	18,22,24	0.79	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	527	1,2	-	0/9/11/13	0/1/1/1
1	PTR	B	527	1,2	-	0/9/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	527	PTR	OH-CZ	-4.29	1.30	1.40
1	A	527	PTR	OH-CZ	-4.14	1.30	1.40

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	527	PTR	O-C-CA	-2.14	119.91	125.49
1	A	527	PTR	CG-CB-CA	-2.10	109.47	114.21

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	VRZ	A	603	-	34,36,36	0.79	2 (5%)	37,50,50	2.19	6 (16%)
4	VRZ	B	603	-	34,36,36	0.82	2 (5%)	37,50,50	2.02	4 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	VRZ	A	603	-	-	0/13/24/24	0/5/5/5
4	VRZ	B	603	-	-	0/13/24/24	0/5/5/5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	603	VRZ	CAM-NBF	-2.62	1.34	1.38
4	B	603	VRZ	CAW-NAV	-2.47	1.33	1.37
4	B	603	VRZ	CAM-NBF	-2.46	1.34	1.38
4	A	603	VRZ	CAW-NAV	-2.20	1.33	1.37

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	603	VRZ	N3-C2-N1	-10.41	120.92	128.89
4	A	603	VRZ	N3-C2-N1	-10.23	121.06	128.89
4	A	603	VRZ	CAM-CBB-CAZ	-2.86	119.94	125.37
4	B	603	VRZ	CAM-CBB-CAZ	-2.71	120.21	125.37
4	A	603	VRZ	OAB-CAW-NAU	-2.69	117.75	122.75
4	A	603	VRZ	CAK-CAZ-CBB	-2.35	117.06	120.87
4	A	603	VRZ	CAM-NBF-CBE	-2.20	123.60	125.44
4	B	603	VRZ	CAX-CAR-NAU	2.25	118.02	112.88
4	B	603	VRZ	NAV-CAW-NAU	2.65	118.31	114.00
4	A	603	VRZ	NAV-CAW-NAU	4.07	120.61	114.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	603	VRZ	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	429/454 (94%)	0.31	22 (5%) 32 31	25, 46, 77, 92	1 (0%)
1	B	430/454 (94%)	0.21	12 (2%) 56 56	22, 42, 66, 91	1 (0%)
All	All	859/908 (94%)	0.26	34 (3%) 42 41	22, 44, 75, 92	2 (0%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	469	PRO	5.6
1	A	213	PHE	5.1
1	A	473	ARG	4.7
1	A	214	TYR	3.9
1	A	207	THR	3.3
1	B	424	PHE	3.3
1	A	475	LEU	3.2
1	A	433	ALA	3.0
1	A	465	GLY	3.0
1	A	142	ASP	2.9
1	A	244	VAL	2.9
1	A	435	ASN	2.9
1	B	305	GLU	2.9
1	B	302	MET	2.8
1	A	86	ILE	2.7
1	A	471	VAL	2.7
1	B	213	PHE	2.6
1	A	437	GLY	2.5
1	B	307	PHE	2.5
1	B	304	VAL	2.4
1	B	465	GLY	2.4
1	B	86	ILE	2.4
1	B	195	ARG	2.4
1	A	305	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	475	LEU	2.3
1	A	206	ARG	2.3
1	B	383	ILE	2.3
1	A	485	GLU	2.2
1	A	468	ASN	2.2
1	B	410	VAL	2.2
1	A	277	GLN	2.1
1	A	434	ILE	2.1
1	A	472	ILE	2.0
1	A	308	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	PTR	B	527	16/17	0.97	0.14	-	30,37,42,42	0
1	PTR	A	527	16/17	0.96	0.14	-	32,38,42,43	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	CA	A	601	1/1	0.95	0.24	4.62	41,41,41,41	0
2	CA	B	601	1/1	0.82	0.30	2.10	45,45,45,45	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	VRZ	A	603	32/32	0.96	0.15	0.66	24,33,56,62	0
4	VRZ	B	603	32/32	0.96	0.14	-0.11	19,27,50,54	0
3	CL	B	602	1/1	0.86	0.10	-1.20	70,70,70,70	0
3	CL	A	602	1/1	0.85	0.11	-1.88	71,71,71,71	0

6.5 Other polymers [i](#)

There are no such residues in this entry.