



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:04 AM GMT

PDB ID : 2VUA
Title : CRYSTAL STRUCTURE OF THE BOTULINUM NEUROTOXIN
SEROTYPE A BINDING DOMAIN
Authors : Stenmark, P.; Dupuy, J.; Stevens, R.C.
Deposited on : 2008-05-22
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	444	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3825 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BOTULINUM NEUROTOXIN A HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	423	3514	2238	607	655	14	0	4	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	311	Total	O	0	0
			311	311		

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	73.00Å 114.50Å 105.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.73 – 1.70 19.73 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.0 (19.73-1.70) 99.0 (19.73-1.70)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.55 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.168 , 0.205 0.178 , 0.211	Depositor DCC
R_{free} test set	2469 reflections (5.36%)	DCC
Wilson B-factor (Å ²)	23.7	Xtrriage
Anisotropy	0.087	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 56.4	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Outliers	0 of 48513 reflections	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3825	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.66	1/3599 (0.0%)	0.77	1/4861 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1070	LYS	CE-NZ	5.18	1.62	1.49

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	996	ARG	NE-CZ-NH1	5.65	123.13	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1255	PHE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3514	0	3479	29	0
2	A	311	0	0	8	0
All	All	3825	0	3479	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1013[A]:ARG:HH12	1:A:1231:ILE:HD11	0.97	1.07
1:A:1013[A]:ARG:NH1	1:A:1231:ILE:HD11	1.80	0.97
1:A:875:ASP:OD2	2:A:2003:HOH:O	1.91	0.87
1:A:1015:ILE:HG22	2:A:2089:HOH:O	1.83	0.78
1:A:1013[A]:ARG:HH12	1:A:1231:ILE:CD1	1.90	0.76
1:A:1232:THR:HG21	2:A:2094:HOH:O	1.94	0.68
1:A:925[B]:GLU:CD	1:A:1052:ASN:HD22	1.98	0.66
1:A:1089:ASP:O	1:A:1093:ASN:N	2.30	0.64
1:A:1092:SER:O	2:A:2143:HOH:O	2.16	0.60
1:A:1070:LYS:HE2	1:A:1071:TYR:CE2	2.37	0.59
1:A:988:GLN:HE21	1:A:992:GLU:HA	1.69	0.58
1:A:1070:LYS:HD3	2:A:2125:HOH:O	2.03	0.58
1:A:898:ILE:HD12	1:A:926:VAL:HG22	1.87	0.56
1:A:1179:ARG:HG2	1:A:1222:VAL:HG22	1.88	0.55
1:A:1070:LYS:HE2	1:A:1071:TYR:CZ	2.42	0.53
1:A:1015:ILE:HD13	1:A:1031:ILE:HG23	1.92	0.51
1:A:920:GLU:HA	1:A:1056:LYS:NZ	2.27	0.50
1:A:905:ASN:HB3	1:A:915:GLN:HB3	1.97	0.47
1:A:1088:TYR:O	1:A:1092:SER:OG	2.15	0.46
1:A:1232:THR:CG2	2:A:2094:HOH:O	2.58	0.46
1:A:985:TRP:CD2	1:A:1019:ILE:HG21	2.51	0.45
1:A:937:MET:HG3	1:A:1046:ASN:HD22	1.82	0.45
1:A:1070:LYS:CD	2:A:2125:HOH:O	2.65	0.42
1:A:946:TRP:HB2	1:A:1070:LYS:HG2	2.02	0.42
1:A:1266:TRP:O	1:A:1270:GLN:HG2	2.19	0.42
1:A:908:PRO:HG2	1:A:909:ILE:HD12	2.02	0.41
1:A:1211:ILE:N	1:A:1212:PRO:CD	2.84	0.41
1:A:1224:LYS:NZ	2:A:2258:HOH:O	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	423/444 (95%)	405 (96%)	17 (4%)	1 (0%)	52 32

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1127	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	394/408 (97%)	389 (99%)	5 (1%)	76 62

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	935	ASN
1	A	1015	ILE
1	A	1065	ARG
1	A	1144	MET
1	A	1169	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	930	ASN
1	A	935	ASN
1	A	988	GLN
1	A	1046	ASN
1	A	1216	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	423/444 (95%)	1.05	64 (15%) 3 3	13, 22, 37, 46	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1271	ILE	10.2
1	A	1256	ASN	8.7
1	A	1231	ILE	8.3
1	A	1272	GLU	7.5
1	A	1166	ALA	5.6
1	A	1274	SER	5.5
1	A	1200	ALA	5.5
1	A	1093	ASN	5.5
1	A	1275	SER	4.8
1	A	1168	GLY	4.6
1	A	1154	LEU	4.4
1	A	875	ASP	4.4
1	A	1276	ARG	4.3
1	A	1294	ARG	4.2
1	A	1273[A]	ARG	4.2
1	A	1227	ASN	4.0
1	A	1198	SER	4.0
1	A	1266	TRP	4.0
1	A	1232	THR	4.0
1	A	1047	ILE	3.9
1	A	1144	MET	3.9
1	A	970	ASN	3.8
1	A	993	ILE	3.8
1	A	1216	ASN	3.7
1	A	920	GLU	3.4
1	A	969	GLU	3.2
1	A	1051	ASN	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1169	ASN	2.9
1	A	1228	ASP	2.9
1	A	954	ASN	2.9
1	A	1201	GLY	2.8
1	A	1081	GLU	2.8
1	A	1257	ASN	2.8
1	A	1092	SER	2.7
1	A	1188	ASN	2.7
1	A	1011	ILE	2.7
1	A	1064	HIS	2.6
1	A	1167	SER	2.5
1	A	1097	LEU	2.5
1	A	1277	THR	2.5
1	A	895	ALA	2.5
1	A	924	ILE	2.4
1	A	991	GLN	2.4
1	A	1206	LEU	2.4
1	A	893	ARG	2.4
1	A	919	LEU	2.3
1	A	1261	LEU	2.3
1	A	1270	GLN	2.3
1	A	1197	ALA	2.3
1	A	1267	TYR	2.3
1	A	971	ASN	2.3
1	A	1269	ARG	2.3
1	A	1262	VAL	2.3
1	A	1057	LEU	2.2
1	A	952	TYR	2.2
1	A	1212	PRO	2.2
1	A	1202	VAL	2.2
1	A	992	GLU	2.2
1	A	898	ILE	2.1
1	A	1086	ASP	2.1
1	A	949	ILE	2.1
1	A	1233	ASN	2.1
1	A	1180	VAL	2.0
1	A	1060	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.