



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 06:08 AM GMT

PDB ID : 2W1A
Title : Non-covalent complex between dahp synthase and chorismate mutase from Mycobacterium tuberculosis with bound tsa
Authors : Okvist, M.; Sasso, S.; Roderer, K.; Gamper, M.; Codoni, G.; Krengel, U.; Kast, P.
Deposited on : 2008-10-16
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	1352 (2.38-2.34)
Clashscore	102246	1456 (2.38-2.34)
Ramachandran outliers	100387	1435 (2.38-2.34)
Sidechain outliers	100360	1436 (2.38-2.34)
RSRZ outliers	91569	1358 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	CE1	A	1465	-	-	-	X
6	PG4	A	1466	-	-	-	X
7	GOL	B	1467	-	-	-	X

2 Entry composition [\(i\)](#)

There are 9 unique types of molecules in this entry. The entry contains 8446 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-DEOXY-D-ARABINO-HEPTULOSONATE 7-PHOSPHATE SYNTHASE AROG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	451	3466	2166	628	655	17	0	0	0
1	B	449	3454	2160	626	651	17	0	0	0


- Molecule 2 is a protein called CHORISMATE MUTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	78	597	370	117	108	2	0	0	0
2	D	79	610	378	121	109	2	0	0	0

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mn		
			1	1	0	0
3	A	1	Total	Mn		
			1	1	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0

- Molecule 5 is O-DODECANYL OCTAETHYLENE GLYCOL (three-letter code: CE1) (formula: C₂₈H₅₈O₉).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 25 20 5	0	0

- Molecule 6 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C₈H₁₈O₅).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 13 8 5	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 6 3 3	0	0
7	B	1	Total C O 6 3 3	0	0
7	B	1	Total C O 6 3 3	0	0

- Molecule 8 is 8-HYDROXY-2-OXA-BICYCLO[3.3.1]NON-6-ENE-3,5-DICARBOXYLIC ACID (three-letter code: TSA) (formula: C₁₀H₁₂O₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	C	1	Total C O 16 10 6	0	0
8	D	1	Total C O 16 10 6	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	94	Total O 94 94	0	0
9	B	105	Total O 105 105	0	0
9	C	6	Total O 6 6	0	0
9	D	9	Total O 9 9	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3-DEOXY-D-ARABINO-HEPTULOSONATE 7-PHOSPHATE SYNTHASE AROG




- Molecule 1: 3-DEOXY-D-ARABINO-HEPTULOSONATE 7-PHOSPHATE SYNTHASE AROG



- Molecule 2: CHORISMATE MUTASE



- Molecule 2: CHORISMATE MUTASE



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	205.91Å 205.91Å 67.24Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.84 – 2.35 39.84 – 2.35	Depositor EDS
% Data completeness (in resolution range)	98.7 (39.84-2.35) 95.2 (39.84-2.35)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.45 (at 2.34Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R , R_{free}	0.191 , 0.229 0.208 , 0.243	Depositor DCC
R_{free} test set	3291 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	28.3	Xtriage
Anisotropy	0.132	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.7	EDS
Estimated twinning fraction	0.028 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 64839 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8446	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MN, TSA, PG4, SO4, CE1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/3536	0.59	0/4809
1	B	0.47	0/3525	0.61	1/4794 (0.0%)
2	C	0.36	0/600	0.54	0/799
2	D	0.38	0/614	0.60	0/817
All	All	0.44	0/8275	0.60	1/11219 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	25	ARG	NE-CZ-NH1	5.74	123.17	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3466	0	3422	26	0
1	B	3454	0	3404	26	0
2	C	597	0	631	0	0
2	D	610	0	650	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	5	0	0	0	0
4	B	10	0	0	0	0
5	A	25	0	41	4	0
6	A	13	0	18	4	0
7	A	6	0	8	1	0
7	B	12	0	16	3	0
8	C	16	0	10	0	0
8	D	16	0	10	0	0
9	A	94	0	0	1	0
9	B	105	0	0	0	0
9	C	6	0	0	0	0
9	D	9	0	0	0	0
All	All	8446	0	8210	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:264:GLY:HA3	1:B:270:GLN:HE21	1.39	0.86
1:B:410:VAL:HG12	1:B:412:ILE:HG23	1.69	0.72
1:B:23:ARG:HE	7:B:1467:GOL:H2	1.56	0.71
1:A:410:VAL:HG13	1:A:412:ILE:HG23	1.73	0.70
1:A:187:THR:O	1:A:241:ALA:HB1	1.93	0.68
1:A:281:ILE:HD13	1:A:293:ILE:HD13	1.81	0.63
5:A:1465:CE1:H151	5:A:1465:CE1:H112	1.80	0.62
1:B:18:LEU:N	7:B:1467:GOL:O2	2.32	0.59
1:B:17:PRO:HA	7:B:1467:GOL:O2	2.04	0.57
1:A:133:LYS:NZ	1:A:440:CYS:SG	2.65	0.56
1:A:425:GLN:HE22	1:A:445:ASN:HD21	1.53	0.56
1:A:117:ALA:HB2	1:A:460:LEU:HD13	1.89	0.54
5:A:1465:CE1:H112	5:A:1465:CE1:C15	2.38	0.54
1:A:197:VAL:HG21	9:A:2059:HOH:O	2.08	0.53
1:A:5:VAL:HG11	1:B:48:MET:CE	2.38	0.53
1:A:5:VAL:HB	1:B:48:MET:HE3	1.92	0.51
1:B:392:GLN:HE22	2:D:54:LEU:CD1	2.24	0.51
1:B:83:GLN:HA	1:B:124:VAL:O	2.10	0.51
1:B:210:ALA:HB3	1:B:424:ALA:HB2	1.93	0.49
1:B:392:GLN:HE22	2:D:54:LEU:HD11	1.77	0.49
1:A:193:SER:O	1:A:197:VAL:HG23	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:9:ILE:HD11	1:B:47:ALA:HB1	1.95	0.48
1:B:410:VAL:CG1	1:B:412:ILE:HG23	2.43	0.48
1:B:133:LYS:NZ	1:B:440:CYS:SG	2.73	0.48
1:A:281:ILE:HD11	1:A:296:ALA:HB2	1.96	0.47
1:A:191:LEU:HD23	1:A:200:TRP:HH2	1.78	0.47
1:A:92:MET:N	6:A:1466:PG4:H32	2.30	0.47
1:A:148:ARG:HD3	7:A:1467:GOL:C3	2.45	0.47
1:B:52:LEU:O	1:B:55:VAL:HG22	2.15	0.46
6:A:1466:PG4:C3	6:A:1466:PG4:H82	2.45	0.46
1:B:260:ARG:HG3	1:B:274:LEU:HD12	1.98	0.45
1:A:96:GLU:HB3	1:A:97:PRO:HD3	1.99	0.45
1:A:410:VAL:HG13	1:A:412:ILE:CG2	2.43	0.45
6:A:1466:PG4:H82	6:A:1466:PG4:O2	2.16	0.45
1:B:84:GLY:HA2	1:B:410:VAL:O	2.17	0.45
1:B:264:GLY:HA3	1:B:270:GLN:NE2	2.20	0.45
1:A:103:VAL:HG13	1:A:243:ILE:HD12	1.99	0.44
1:A:5:VAL:HG21	1:B:48:MET:HE1	1.98	0.44
1:A:83:GLN:HA	1:A:124:VAL:O	2.17	0.43
1:A:92:MET:H	6:A:1466:PG4:H32	1.84	0.42
1:A:12:LEU:HD22	1:B:92:MET:HA	2.00	0.42
1:A:171:ARG:HH21	5:A:1465:CE1:H171	1.84	0.42
1:A:95:THR:HG22	5:A:1465:CE1:H231	2.01	0.42
1:B:381:THR:HA	1:B:442:PRO:HG3	2.01	0.42
1:A:5:VAL:CB	1:B:48:MET:HE3	2.50	0.41
1:B:45:ALA:O	1:B:49:ARG:HG3	2.21	0.41
1:B:410:VAL:HG12	1:B:412:ILE:CG2	2.45	0.41
1:A:191:LEU:CD2	1:A:200:TRP:HH2	2.34	0.41
1:A:84:GLY:HA2	1:A:410:VAL:O	2.21	0.40
1:B:446:THR:HG23	1:B:447:GLN:H	1.86	0.40
1:B:179:ALA:O	1:B:183:VAL:HG13	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	445/472 (94%)	434 (98%)	11 (2%)	0	100 100
1	B	443/472 (94%)	431 (97%)	12 (3%)	0	100 100
2	C	76/90 (84%)	75 (99%)	1 (1%)	0	100 100
2	D	77/90 (86%)	77 (100%)	0	0	100 100
All	All	1041/1124 (93%)	1017 (98%)	24 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	366/385 (95%)	353 (96%)	13 (4%)	42 55
1	B	363/385 (94%)	360 (99%)	3 (1%)	86 94
2	C	60/75 (80%)	59 (98%)	1 (2%)	68 82
2	D	62/75 (83%)	62 (100%)	0	100 100
All	All	851/920 (92%)	834 (98%)	17 (2%)	63 77

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	THR
1	A	40	TRP
1	A	65	VAL
1	A	104	ARG
1	A	138	ASP
1	A	177	SER
1	A	194	LEU
1	A	196	LEU
1	A	215	GLU
1	A	266	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	270	GLN
1	A	337	ARG
1	A	410	VAL
1	B	15	LEU
1	B	40	TRP
1	B	337	ARG
2	C	53	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	GLN
1	A	70	GLN
1	A	425	GLN
1	B	270	GLN
1	B	373	HIS
1	B	383	HIS
1	B	392	GLN
2	C	56	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	SO4	A	1464	-	4,4,4	0.28	0	6,6,6	0.38	0
5	CE1	A	1465	-	24,24,36	0.76	0	23,23,35	0.61	1 (4%)
6	PG4	A	1466	-	12,12,12	0.51	0	11,11,11	0.70	0
7	GOL	A	1467	-	5,5,5	0.39	0	5,5,5	0.41	0
4	SO4	B	1463	-	4,4,4	0.39	0	6,6,6	0.18	0
4	SO4	B	1464	-	4,4,4	0.44	0	6,6,6	0.53	0
7	GOL	B	1466	-	5,5,5	0.47	0	5,5,5	0.39	0
7	GOL	B	1467	-	5,5,5	0.72	0	5,5,5	0.99	0
8	TSA	C	1091	-	11,17,17	0.99	0	7,26,26	1.71	1 (14%)
8	TSA	D	1091	-	11,17,17	1.05	1 (9%)	7,26,26	1.82	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SO4	A	1464	-	-	0/0/0/0	0/0/0/0
5	CE1	A	1465	-	-	0/22/22/34	0/0/0/0
6	PG4	A	1466	-	-	0/10/10/10	0/0/0/0
7	GOL	A	1467	-	-	0/4/4/4	0/0/0/0
4	SO4	B	1463	-	-	0/0/0/0	0/0/0/0
4	SO4	B	1464	-	-	0/0/0/0	0/0/0/0
7	GOL	B	1466	-	-	0/4/4/4	0/0/0/0
7	GOL	B	1467	-	-	0/4/4/4	0/0/0/0
8	TSA	C	1091	-	-	0/0/34/34	0/0/2/2
8	TSA	D	1091	-	-	0/0/34/34	0/0/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	D	1091	TSA	C6-C1	-2.30	1.51	1.56

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1465	CE1	O16-C15-C14	2.06	119.54	110.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	D	1091	TSA	O7-C5-C6	2.13	112.58	109.81
8	D	1091	TSA	C5-O7-C8	3.83	120.03	112.28
8	C	1091	TSA	C5-O7-C8	3.90	120.18	112.28

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1465	CE1	4	0
6	A	1466	PG4	4	0
7	A	1467	GOL	1	0
7	B	1467	GOL	3	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	451/472 (95%)	0.20	24 (5%) 30 45	16, 26, 43, 51	0
1	B	449/472 (95%)	0.32	32 (7%) 19 28	16, 24, 50, 63	0
2	C	78/90 (86%)	0.83	13 (16%) 2 4	38, 42, 48, 51	0
2	D	79/90 (87%)	0.74	10 (12%) 5 9	35, 42, 50, 51	0
All	All	1057/1124 (94%)	0.33	79 (7%) 17 26	16, 27, 48, 63	0

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	12	PRO	6.8
1	B	377	THR	6.3
1	B	3	TRP	6.0
1	B	376	SER	5.6
1	B	379	PHE	5.2
1	B	10	ASP	4.7
2	D	16	THR	4.7
1	A	430	THR	4.6
1	B	430	THR	4.6
2	C	16	THR	4.5
1	A	11	GLN	4.2
2	C	14	ILE	4.2
1	B	375	SER	4.1
1	A	377	THR	4.1
1	B	432	LEU	4.1
1	B	429	GLU	3.5
1	A	379	PHE	3.4
2	C	47	MET	3.4
1	B	431	ASP	3.3
2	D	48	ALA	3.3
2	C	17	LEU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	233	VAL	3.2
1	B	434	GLY	3.2
1	B	433	ALA	3.2
1	B	426	ASP	3.1
1	B	378	GLY	3.1
1	B	233	VAL	3.1
1	A	268	GLU	3.1
1	B	428	SER	3.1
1	B	373	HIS	3.0
2	D	14	ILE	2.9
2	C	48	ALA	2.9
1	B	15	LEU	2.8
2	C	45	ALA	2.8
1	B	408	ILE	2.8
1	B	435	ARG	2.7
2	C	15	ASP	2.7
1	A	431	ASP	2.7
1	A	376	SER	2.7
1	A	410	VAL	2.7
2	D	49	SER	2.6
1	B	266	ASP	2.6
1	A	433	ALA	2.6
1	B	195	HIS	2.6
2	C	50	GLY	2.6
1	B	410	VAL	2.6
1	A	3	TRP	2.5
1	B	418	THR	2.5
1	B	416	ASN	2.4
1	A	429	GLU	2.4
2	C	49	SER	2.4
2	D	88	LEU	2.4
1	B	245	ALA	2.4
1	A	124	VAL	2.3
2	D	44	LYS	2.3
1	A	241	ALA	2.3
1	A	408	ILE	2.2
2	D	13	GLU	2.2
2	C	34	ARG	2.2
1	A	434	GLY	2.2
2	C	43	GLY	2.2
1	A	128	ALA	2.2
1	A	462	ASP	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	374	GLU	2.2
2	C	60	MET	2.2
1	A	424	ALA	2.2
1	B	128	ALA	2.2
1	B	279	VAL	2.1
1	A	195	HIS	2.1
2	D	15	ASP	2.1
1	A	125	ALA	2.1
2	C	27	GLU	2.1
1	B	82	LEU	2.1
1	B	125	ALA	2.1
1	B	267	GLY	2.1
1	A	373	HIS	2.1
1	A	356	ALA	2.1
1	A	206	ARG	2.0
2	D	17	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	PG4	A	1466	13/13	0.91	0.38	7.27	36,38,40,41	0
5	CE1	A	1465	25/37	0.78	0.26	5.61	41,52,54,55	0
7	GOL	B	1467	6/6	0.88	0.27	4.36	37,39,40,41	0
7	GOL	A	1467	6/6	0.83	0.24	1.75	38,40,43,45	0
7	GOL	B	1466	6/6	0.88	0.20	0.67	50,50,52,52	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	SO4	B	1464	5/5	0.99	0.09	-0.80	34,35,36,36	0
8	TSA	C	1091	16/16	0.94	0.12	-0.95	34,36,38,40	0
3	MN	A	1463	1/1	0.95	0.10	-0.97	33,33,33,33	0
8	TSA	D	1091	16/16	0.93	0.11	-1.33	35,37,38,38	0
4	SO4	A	1464	5/5	0.98	0.08	-1.97	32,33,35,37	0
4	SO4	B	1463	5/5	0.99	0.10	-2.13	26,26,27,31	0
3	MN	B	1465	1/1	0.97	0.04	-2.38	38,38,38,38	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.