



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:08 AM GMT

PDB ID : 2W1U  
Title : A FAMILY 32 CARBOHYDRATE-BINDING MODULE, FROM THE MU  
TOXIN PRODUCED BY CLOSTRIDIUM PERFRINGENS, IN COMPLEX  
WITH BETA-D-GLCNAC-BETA(1,3)GALNAC  
Authors : Ficko-Blean, E.; Boraston, A.B.  
Deposited on : 2008-10-20  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	192	<div> <div>%</div> <div> <div></div> <div>65%</div> <div>7%</div> <div>28%</div> </div> </div>
1	B	192	<div> <div>%</div> <div> <div></div> <div>67%</div> <div>8%</div> <div>24%</div> </div> </div>
1	C	192	<div> <div>%</div> <div> <div></div> <div>67%</div> <div>6%</div> <div>27%</div> </div> </div>
1	D	192	<div> <div>%</div> <div> <div></div> <div>65%</div> <div>9%</div> <div>25%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NGA	A	1947	-	-	-	X
4	ACT	B	1953	-	-	-	X
4	ACT	C	1951	-	-	-	X
4	ACT	D	1951	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5331 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HYALURONOGLUCOSAMINIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	139	Total	C	N	O	S	0	0	0
			1100	699	177	223	1			
1	B	146	Total	C	N	O	S	0	0	0
			1153	730	184	237	2			
1	C	140	Total	C	N	O	S	0	0	0
			1109	704	177	227	1			
1	D	144	Total	C	N	O	S	3	0	0
			1136	719	182	233	2			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	944	VAL	ILE	VARIANT	UNP P26831
B	944	VAL	ILE	VARIANT	UNP P26831
C	944	VAL	ILE	VARIANT	UNP P26831
D	944	VAL	ILE	VARIANT	UNP P26831

- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	2	Total	C	N	O	0	0
			29	16	2	11		
2	B	2	Total	C	N	O	0	0
			29	16	2	11		
2	D	2	Total	C	N	O	0	0
			29	16	2	11		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Ca	0	0
			1	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		
3	D	1	Total	Ca	0	0
			1	1		
3	C	1	Total	Ca	0	0
			1	1		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	2	Total	C	N	O	0	0
			29	16	2	11		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	154	Total 154	O 154	0	0
6	B	171	Total 171	O 171	0	0
6	C	197	Total 197	O 197	0	0
6	D	179	Total 179	O 179	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	91.17Å 91.17Å 132.61Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.00 19.89 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.6 (20.00-2.00) 98.5 (19.89-2.00)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.75 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, $R_{free}$	0.169 , 0.213 0.174 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	19.0	Xtriage
Anisotropy	0.031	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 54.5	EDS
Estimated twinning fraction	0.013 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Outliers	0 of 43004 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5331	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, A2G, NAG, NGA, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.74	0/1123	0.71	0/1518
1	B	0.80	0/1176	0.72	0/1589
1	C	0.75	0/1132	0.72	0/1529
1	D	0.82	1/1159 (0.1%)	1.01	2/1566 (0.1%)
All	All	0.78	1/4590 (0.0%)	0.80	2/6202 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	804	MET	CB-CG	-7.35	1.27	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	804	MET	CA-CB-CG	27.76	160.49	113.30
1	D	804	MET	CB-CG-SD	5.87	130.02	112.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1100	0	1058	8	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1153	0	1103	16	0
1	C	1109	0	1062	7	0
1	D	1136	0	1086	15	0
2	A	29	0	27	0	0
2	B	29	0	27	0	0
2	D	29	0	27	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	B	4	0	3	0	0
4	C	4	0	3	1	0
4	D	4	0	3	1	0
5	C	29	0	27	0	0
6	A	154	0	0	0	1
6	B	171	0	0	1	1
6	C	197	0	0	1	1
6	D	179	0	0	4	1
All	All	5331	0	4426	43	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:808:PRO:N	6:C:2001:HOH:O	2.05	0.89
1:C:822:ASN:HD22	1:C:824:ALA:H	1.29	0.80
1:B:807:ASN:H	1:B:807:ASN:HD22	1.33	0.77
1:A:826:LEU:HD21	1:A:835:VAL:HG12	1.74	0.69
1:B:871:ASN:ND2	1:B:908:LYS:HE2	2.11	0.64
1:B:807:ASN:ND2	1:D:812:ARG:H	1.95	0.64
1:C:810:LEU:HD11	1:C:851:ILE:HD11	1.79	0.64
1:D:932:ILE:HG12	1:D:934:LYS:HG3	1.80	0.62
1:B:812:ARG:H	1:D:807:ASN:ND2	1.99	0.61
1:C:822:ASN:ND2	1:C:824:ALA:H	1.96	0.60
1:D:822:ASN:HD22	1:D:822:ASN:C	2.06	0.59
1:A:870:LYS:HA	1:A:908:LYS:HG2	1.84	0.59
1:D:804:MET:N	6:D:2001:HOH:O	2.39	0.56
1:D:947:GLU:HG3	6:D:2163:HOH:O	2.07	0.54
1:A:871:ASN:HD21	1:A:908:LYS:HZ3	1.56	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:879:TRP:CE2	1:B:928:ASN:HB2	2.44	0.53
1:B:871:ASN:HD21	1:B:908:LYS:HE2	1.74	0.52
1:C:871:ASN:ND2	1:C:908:LYS:H	2.07	0.52
1:B:948:LEU:O	1:B:949:GLU:HB2	2.08	0.51
1:A:871:ASN:HD21	1:A:908:LYS:NZ	2.07	0.51
1:D:822:ASN:ND2	1:D:825:ASN:H	2.09	0.50
1:D:842:GLY:O	6:D:2063:HOH:O	2.20	0.50
1:B:807:ASN:N	1:B:807:ASN:HD22	2.06	0.50
1:D:908:LYS:HB2	1:D:908:LYS:HE2	1.68	0.48
1:B:855:LEU:HB3	1:B:859:ILE:HD12	1.96	0.46
1:B:807:ASN:H	1:B:807:ASN:ND2	2.07	0.46
1:A:887:SER:HB2	1:A:893:TRP:CE3	2.49	0.46
1:B:807:ASN:HD21	1:D:812:ARG:H	1.61	0.46
1:B:880:ASN:ND2	6:B:2090:HOH:O	2.49	0.46
1:D:826:LEU:HD21	1:D:835:VAL:HG12	1.99	0.44
1:B:811:ILE:HG22	1:B:850:PHE:HE2	1.82	0.44
1:A:879:TRP:CE2	1:A:928:ASN:HB2	2.53	0.44
1:C:817:GLN:HB3	4:C:1951:ACT:H2	2.00	0.43
1:B:807:ASN:N	1:B:807:ASN:ND2	2.66	0.43
1:A:822:ASN:O	1:A:825:ASN:HB2	2.19	0.43
4:D:1951:ACT:H1	6:D:2178:HOH:O	2.19	0.42
1:A:869:GLY:HA2	1:A:937:THR:OG1	2.20	0.42
1:B:807:ASN:HD21	1:D:811:ILE:HA	1.83	0.42
1:D:946:ASP:O	1:D:947:GLU:HB2	2.20	0.41
1:D:879:TRP:CE2	1:D:928:ASN:HB2	2.56	0.41
1:B:916:GLU:HA	1:B:916:GLU:OE1	2.21	0.40
1:C:860:LYS:HE2	1:C:947:GLU:OE2	2.21	0.40
1:D:933:ASN:O	1:D:934:LYS:HG2	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:2099:HOH:O	6:D:2136:HOH:O[2_445]	1.89	0.31
6:B:2107:HOH:O	6:C:2112:HOH:O[2_545]	2.05	0.15

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	137/192 (71%)	132 (96%)	5 (4%)	0	100	100
1	B	144/192 (75%)	141 (98%)	3 (2%)	0	100	100
1	C	138/192 (72%)	133 (96%)	5 (4%)	0	100	100
1	D	142/192 (74%)	139 (98%)	2 (1%)	1 (1%)	26	19
All	All	561/768 (73%)	545 (97%)	15 (3%)	1 (0%)	52	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	935	TRP

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	119/165 (72%)	118 (99%)	1 (1%)	86	89
1	B	125/165 (76%)	121 (97%)	4 (3%)	46	44
1	C	120/165 (73%)	117 (98%)	3 (2%)	55	55
1	D	123/165 (74%)	120 (98%)	3 (2%)	57	58
All	All	487/660 (74%)	476 (98%)	11 (2%)	58	60

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	916	GLU
1	B	807	ASN
1	B	822	ASN
1	B	860	LYS
1	B	888	LEU
1	C	822	ASN
1	C	888	LEU
1	C	916	GLU
1	D	804	MET
1	D	822	ASN
1	D	888	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	871	ASN
1	B	807	ASN
1	B	871	ASN
1	B	880	ASN
1	C	822	ASN
1	C	871	ASN
1	C	880	ASN
1	D	807	ASN
1	D	822	ASN
1	D	880	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

8 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	A	1946	2	14,14,15	0.52	0	15,19,21	0.69	0
2	NGA	A	1947	2	15,15,15	0.52	0	17,21,21	1.19	1 (5%)
2	NAG	B	1950	2	14,14,15	0.63	0	15,19,21	1.07	1 (6%)
2	NGA	B	1951	2	15,15,15	0.50	0	17,21,21	0.98	1 (5%)
5	NAG	C	1948	5	14,14,15	0.60	0	15,19,21	1.19	2 (13%)
5	A2G	C	1949	5	15,15,15	0.59	0	17,21,21	0.82	1 (5%)
2	NAG	D	1948	2	14,14,15	0.81	0	15,19,21	1.03	0
2	NGA	D	1949	2	15,15,15	0.55	0	17,21,21	0.91	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1946	2	-	0/6/23/26	0/1/1/1
2	NGA	A	1947	2	-	0/6/26/26	0/1/1/1
2	NAG	B	1950	2	-	0/6/23/26	0/1/1/1
2	NGA	B	1951	2	-	0/6/26/26	0/1/1/1
5	NAG	C	1948	5	-	0/6/23/26	0/1/1/1
5	A2G	C	1949	5	-	0/6/26/26	0/1/1/1
2	NAG	D	1948	2	-	0/6/23/26	0/1/1/1
2	NGA	D	1949	2	-	0/6/26/26	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1951	NGA	O3-C3-C2	-2.45	104.66	109.66
2	B	1950	NAG	O7-C7-C8	-2.20	118.03	122.06
5	C	1948	NAG	C2-N2-C7	-2.15	120.28	123.04
5	C	1949	A2G	C1-O-C5	-2.13	109.53	113.47
5	C	1948	NAG	C1-O5-C5	2.34	115.22	112.25
2	A	1947	NGA	C3-C2-N2	3.02	116.92	110.66

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	ACT	B	1953	-	1,3,3	1.50	0	0,3,3	0.00	-
4	ACT	C	1951	-	1,3,3	1.15	0	0,3,3	0.00	-
4	ACT	D	1951	-	1,3,3	1.05	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ACT	B	1953	-	-	0/0/0/0	0/0/0/0
4	ACT	C	1951	-	-	0/0/0/0	0/0/0/0
4	ACT	D	1951	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	1951	ACT	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	1951	ACT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	139/192 (72%)	-0.33	2 (1%) 78 78	12, 20, 35, 39	0
1	B	146/192 (76%)	-0.56	2 (1%) 78 78	10, 15, 28, 53	0
1	C	140/192 (72%)	-0.56	2 (1%) 78 78	11, 15, 32, 44	0
1	D	144/192 (75%)	-0.62	2 (1%) 78 78	11, 16, 25, 43	1 (0%)
All	All	569/768 (74%)	-0.52	8 (1%) 78 78	10, 16, 32, 53	1 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	916	GLU	3.5
1	A	856	GLY	2.9
1	A	916	GLU	2.8
1	C	946	ASP	2.4
1	C	916	GLU	2.4
1	B	949	GLU	2.4
1	D	946	ASP	2.3
1	B	946	ASP	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NGA	A	1947	15/15	0.90	0.18	5.09	30,36,42,44	0
5	A2G	C	1949	15/15	0.92	0.09	1.34	13,16,19,22	0
2	NAG	A	1946	14/15	0.96	0.07	-0.71	19,23,27,28	0
2	NAG	D	1948	14/15	0.96	0.07	-0.81	12,17,19,19	0
5	NAG	C	1948	14/15	0.97	0.07	-1.08	11,13,17,20	0
2	NAG	B	1950	14/15	0.96	0.07	-1.14	13,16,17,18	0
2	NGA	D	1949	15/15	0.88	0.17	-	24,30,36,37	0
2	NGA	B	1951	15/15	0.90	0.14	-	19,22,28,32	0

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	ACT	C	1951	4/4	0.76	0.28	8.39	19,21,22,22	0
4	ACT	D	1951	4/4	0.88	0.22	7.78	27,27,27,29	0
4	ACT	B	1953	4/4	0.53	0.28	5.12	47,47,47,48	0
3	CA	A	1948	1/1	0.98	0.03	-1.68	19,19,19,19	0
3	CA	C	1950	1/1	1.00	0.03	-2.03	14,14,14,14	0
3	CA	B	1952	1/1	0.99	0.04	-2.21	14,14,14,14	0
3	CA	D	1950	1/1	1.00	0.04	-2.90	17,17,17,17	0

## 6.5 Other polymers

There are no such residues in this entry.