



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:14 AM GMT

PDB ID : 2WG1
Title : TERNARY COMPLEX OF THE AGED CONJUGATE OF TORPEDO CALIFORNICA ACETYLCHELYNOLINESTERASE WITH SOMAN AND 2-PAM
Authors : Sanson, B.; Nachon, F.; Colletier, J.P.; Froment, M.T.; Toker, L.; Greenblatt, H.M.; Sussman, J.L.; Ashani, Y.; Masson, P.; Silman, I.; Weik, M.
Deposited on : 2009-04-15
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	537	<div> <div>2%</div> <div>86%</div> <div>13%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	FP1	A	2540	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	MES	A	2541	-	-	-	X
6	MES	A	2542	-	-	X	-
7	PG4	A	2545	-	-	-	X
7	PG4	A	2546	-	-	-	X
7	PG4	A	2547	-	-	-	X
7	PG4	A	2548	-	-	-	X
8	PGE	A	2549	-	-	-	X
8	PGE	A	2550	-	-	-	X
8	PGE	A	2551	-	-	-	X
9	PEG	A	2555	-	-	-	X

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 5221 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACETYLCHOLINESTERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	532	Total	C	N	O	S	0	33	0
			4449	2864	752	810	23			

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			24	14	1	9		

- Molecule 4 is METHYLPHOSPHONIC ACID ESTER GROUP (three-letter code: GB) (formula: $\text{CH}_5\text{O}_3\text{P}$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	0
			4	1	2	1		

- Molecule 5 is N-HYDROXY-1-(1-METHYLPYRIDIN-2(1H)-YLIDENE)METHANAMINE (three-letter code: FP1) (formula: $\text{C}_7\text{H}_{10}\text{N}_2\text{O}$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			10	7	2	1		

- Molecule 6 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: $C_6H_{13}NO_4S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
6	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
6	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
6	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 7 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			13	8	5		
7	A	1	Total	C	O	0	0
			13	8	5		
7	A	1	Total	C	O	0	0
			13	8	5		
7	A	1	Total	C	O	0	0
			13	8	5		

- Molecule 8 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			10	6	4		
8	A	1	Total	C	O	0	0
			10	6	4		
8	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 9 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	O	0	0
			7	4	3		
9	A	1	Total	C	O	0	0
			7	4	3		
9	A	1	Total	C	O	0	0
			7	4	3		
9	A	1	Total	C	O	0	0
			7	4	3		
9	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	555	Total	O	0	0
			555	555		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ACETYLCHOLINESTERASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	111.91Å 111.91Å 137.17Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.20 20.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.6 (20.00-2.20) 98.8 (20.00-2.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.82 (at 2.19Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.166 , 0.212 0.162 , 0.208	Depositor DCC
R_{free} test set	2462 reflections (5.15%)	DCC
Wilson B-factor (Å ²)	30.7	Xtriage
Anisotropy	0.616	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 51.9	EDS
Estimated twinning fraction	0.028 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 50238 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5221	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, NAG, PG4, FP1, FUC, MES, PEG, GB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	0/4663	0.65	3/6323 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	494	LEU	CA-CB-CG	5.80	128.63	115.30
1	A	388	ARG	NE-CZ-NH2	-5.75	117.42	120.30
1	A	388	ARG	NE-CZ-NH1	5.31	122.95	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4449	0	4389	63	0
2	A	14	0	13	0	0
3	A	24	0	22	0	0
4	A	4	0	3	0	0
5	A	10	0	8	2	0
6	A	48	0	52	8	0
7	A	52	0	72	8	0
8	A	30	0	42	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	A	35	0	50	1	0
10	A	555	0	0	21	0
All	All	5221	0	4651	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 66 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:220[B]:ARG:NH1	10:A:1258:HOH:O	1.62	1.19
1:A:181[B]:HIS:ND1	10:A:1230:HOH:O	1.90	1.04
1:A:353[A]:MET:HE2	10:A:1362:HOH:O	1.70	0.90
1:A:19[B]:ARG:NH1	10:A:1016:HOH:O	1.99	0.89
1:A:504[B]:ASP:OD2	10:A:1488:HOH:O	1.91	0.87

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	563/537 (105%)	537 (95%)	26 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	498/469 (106%)	486 (98%)	12 (2%)	57	69

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	430	LEU
1	A	454[A]	LYS
1	A	487	SER
1	A	288	PHE
1	A	473	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	2537	1,3	14,14,15	0.62	0	15,19,21	1.00	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FUC	A	2538	3	10,10,11	0.66	0	14,14,16	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	2537	1,3	-	0/6/23/26	0/1/1/1
3	FUC	A	2538	3	-	0/0/17/20	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	2536	1	14,14,15	0.58	0	15,19,21	0.85	0
4	GB	A	2539	1	0,3,4	0.00	-	0,3,6	0.00	-
5	FP1	A	2540	-	8,10,10	3.60	2 (25%)	8,12,12	3.50	3 (37%)
6	MES	A	2541	-	11,12,12	0.96	0	14,16,16	1.50	4 (28%)
6	MES	A	2542	-	11,12,12	1.15	0	14,16,16	2.06	2 (14%)
6	MES	A	2543	-	11,12,12	1.10	0	14,16,16	1.42	3 (21%)
6	MES	A	2544	-	11,12,12	1.07	0	14,16,16	1.36	2 (14%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	PG4	A	2545	-	12,12,12	0.74	0	11,11,11	0.76	0
7	PG4	A	2546	-	12,12,12	0.69	0	11,11,11	1.01	0
7	PG4	A	2547	-	12,12,12	0.72	0	11,11,11	0.50	0
7	PG4	A	2548	-	12,12,12	0.70	0	11,11,11	0.44	0
8	PGE	A	2549	-	9,9,9	0.70	0	8,8,8	0.49	0
8	PGE	A	2550	-	9,9,9	0.77	0	8,8,8	0.73	0
8	PGE	A	2551	-	9,9,9	0.60	0	8,8,8	0.40	0
9	PEG	A	2552	-	6,6,6	0.60	0	5,5,5	0.47	0
9	PEG	A	2553	-	6,6,6	0.60	0	5,5,5	0.31	0
9	PEG	A	2554	-	6,6,6	0.62	0	5,5,5	0.33	0
9	PEG	A	2555	-	6,6,6	0.49	0	5,5,5	1.06	1 (20%)
9	PEG	A	2556	-	6,6,6	0.83	0	5,5,5	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	2536	1	-	0/6/23/26	0/1/1/1
4	GB	A	2539	1	-	0/0/0/0	0/0/0/0
5	FP1	A	2540	-	-	0/0/3/3	0/1/1/1
6	MES	A	2541	-	-	0/6/14/14	0/1/1/1
6	MES	A	2542	-	-	0/6/14/14	0/1/1/1
6	MES	A	2543	-	-	0/6/14/14	0/1/1/1
6	MES	A	2544	-	-	0/6/14/14	0/1/1/1
7	PG4	A	2545	-	-	0/10/10/10	0/0/0/0
7	PG4	A	2546	-	-	0/10/10/10	0/0/0/0
7	PG4	A	2547	-	-	0/10/10/10	0/0/0/0
7	PG4	A	2548	-	-	0/10/10/10	0/0/0/0
8	PGE	A	2549	-	-	0/7/7/7	0/0/0/0
8	PGE	A	2550	-	-	0/7/7/7	0/0/0/0
8	PGE	A	2551	-	-	0/7/7/7	0/0/0/0
9	PEG	A	2552	-	-	0/4/4/4	0/0/0/0
9	PEG	A	2553	-	-	0/4/4/4	0/0/0/0
9	PEG	A	2554	-	-	0/4/4/4	0/0/0/0
9	PEG	A	2555	-	-	0/4/4/4	0/0/0/0
9	PEG	A	2556	-	-	0/4/4/4	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	2540	FP1	C6-N2	-9.42	1.18	1.30
5	A	2540	FP1	C1-N1	2.99	1.40	1.33

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	2540	FP1	C3-C4-C5	-7.27	118.94	121.21
5	A	2540	FP1	C1-N1-C5	-2.90	118.57	121.74
6	A	2541	MES	C2-C3-N4	-2.68	106.06	110.12
6	A	2544	MES	O3S-S-O1S	-2.54	105.71	111.61
6	A	2541	MES	O1-C6-C5	-2.32	106.51	111.84

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	2540	FP1	2	0
6	A	2542	MES	6	0
6	A	2543	MES	1	0
6	A	2544	MES	1	0
7	A	2546	PG4	6	0
7	A	2547	PG4	1	0
7	A	2548	PG4	1	0
8	A	2551	PGE	2	0
9	A	2555	PEG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	532/537 (99%)	-0.65	9 (1%) 73 72	17, 27, 41, 61	5 (0%)

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	486	HIS	5.3
1	A	485	PRO	4.8
1	A	488	GLN	4.5
1	A	55	SER	3.5
1	A	535	THR	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	FUC	A	2538	10/11	0.85	0.36	-	85,87,88,89	0
3	NAG	A	2537	14/15	0.88	0.26	-	59,66,68,75	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
9	PEG	A	2555	7/7	0.90	0.27	13.82	58,59,60,60	0
8	PGE	A	2551	10/10	0.71	0.29	10.18	70,78,82,82	0
7	PG4	A	2545	13/13	0.79	0.21	6.97	60,62,64,65	0
8	PGE	A	2550	10/10	0.89	0.22	6.78	56,57,58,58	0
7	PG4	A	2546	13/13	0.72	0.32	6.12	61,64,69,69	0
7	PG4	A	2547	13/13	0.74	0.27	5.31	69,70,75,75	0
6	MES	A	2541	12/12	0.77	0.24	4.84	56,59,62,63	12
8	PGE	A	2549	10/10	0.68	0.29	3.45	60,63,69,69	0
7	PG4	A	2548	13/13	0.73	0.25	3.07	85,88,91,91	0
5	FP1	A	2540	10/10	0.95	0.13	2.94	28,30,43,45	0
6	MES	A	2543	12/12	0.96	0.12	1.73	41,46,48,49	12
6	MES	A	2544	12/12	0.91	0.17	1.70	52,57,59,60	12
4	GB	A	2539	4/5	0.99	0.08	1.64	19,20,22,22	0
9	PEG	A	2552	7/7	0.71	0.26	1.61	68,69,71,71	0
6	MES	A	2542	12/12	0.87	0.27	0.96	54,62,67,67	12
9	PEG	A	2553	7/7	0.88	0.20	-	72,72,74,75	0
2	NAG	A	2536	14/15	0.90	0.26	-	40,44,49,50	0
9	PEG	A	2554	7/7	0.76	0.24	-	97,98,99,99	0
9	PEG	A	2556	7/7	0.69	0.28	-	64,68,73,74	0

6.5 Other polymers [i](#)

There are no such residues in this entry.