



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:16 AM GMT

PDB ID : 2WG6  
Title : PROTEASOME-ACTIVATING NUCLEOTIDASE (PAN) N-DOMAIN (57-134) FROM ARCHAEOGLOBUS FULGIDUS FUSED TO GCN4, P61A MUTANT  
Authors : Hartmann, M.D.; Djuranovic, S.; Ursinus, A.; Zeth, K.; Lupas, A.N.  
Deposited on : 2009-04-15  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	109	<div> <div>3%</div> <div>76%</div> <div>20%</div> </div>
1	B	109	<div> <div>2%</div> <div>66%</div> <div>14%</div> <div>20%</div> </div>
1	C	109	<div> <div>0%</div> <div>75%</div> <div>5%</div> <div>20%</div> </div>
1	D	109	<div> <div>68%</div> <div>12%</div> <div>20%</div> </div>
1	E	109	<div> <div>75%</div> <div>5%</div> <div>20%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	109	<div><div><div></div><div></div><div></div></div><div><div></div><div></div><div></div></div><div>%65%15%20%</div></div>
1	G	109	<div><div><div></div><div></div><div></div></div><div><div></div><div></div><div></div></div><div>2%72%6%20%</div></div>
1	H	109	<div><div><div></div><div></div><div></div></div><div><div></div><div></div><div></div></div><div>%67%12%20%</div></div>
1	I	109	<div><div><div></div><div></div><div></div></div><div><div></div><div></div><div></div></div><div>2%69%9%20%</div></div>
1	J	109	<div><div><div></div><div></div><div></div></div><div><div></div><div></div><div></div></div><div>%65%14%20%</div></div>
1	K	109	<div><div><div></div><div></div><div></div></div><div><div></div><div></div><div></div></div><div>3%70%8%20%</div></div>
1	L	109	<div><div><div></div><div></div><div></div></div><div><div></div><div></div><div></div></div><div>%66%13%20%</div></div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8199 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GENERAL CONTROL PROTEIN GCN4, PROTEASOME-ACTIVATING NUCLEOTIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	87	Total	C	N	O	S	0	0	0
			666	422	117	126	1			
1	B	87	Total	C	N	O	S	0	0	0
			664	420	115	128	1			
1	C	87	Total	C	N	O	S	0	0	0
			666	422	117	126	1			
1	D	87	Total	C	N	O	S	0	0	0
			668	423	116	128	1			
1	E	87	Total	C	N	O	S	0	0	0
			666	422	117	126	1			
1	F	87	Total	C	N	O	S	0	0	0
			668	423	116	128	1			
1	G	87	Total	C	N	O	S	0	0	0
			674	425	116	132	1			
1	H	87	Total	C	N	O	S	0	0	0
			656	416	114	125	1			
1	I	87	Total	C	N	O	S	0	0	0
			674	425	116	132	1			
1	J	87	Total	C	N	O	S	0	0	0
			656	416	114	125	1			
1	K	87	Total	C	N	O	S	0	0	0
			674	425	116	132	1			
1	L	87	Total	C	N	O	S	0	0	0
			656	416	114	125	1			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	61	ALA	PRO	ENGINEERED MUTATION	UNP O28303
B	61	ALA	PRO	ENGINEERED MUTATION	UNP O28303
C	61	ALA	PRO	ENGINEERED MUTATION	UNP O28303
D	61	ALA	PRO	ENGINEERED MUTATION	UNP O28303

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	61	ALA	PRO	ENGINEERED MUTATION	UNP O28303
F	61	ALA	PRO	ENGINEERED MUTATION	UNP O28303
G	61	ALA	PRO	ENGINEERED MUTATION	UNP O28303
H	61	ALA	PRO	ENGINEERED MUTATION	UNP O28303
I	61	ALA	PRO	ENGINEERED MUTATION	UNP O28303
J	61	ALA	PRO	ENGINEERED MUTATION	UNP O28303
K	61	ALA	PRO	ENGINEERED MUTATION	UNP O28303
L	61	ALA	PRO	ENGINEERED MUTATION	UNP O28303


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	21	Total O 21 21	0	0
2	B	22	Total O 22 22	0	0
2	C	18	Total O 18 18	0	0
2	D	19	Total O 19 19	0	0
2	E	16	Total O 16 16	0	0
2	F	20	Total O 20 20	0	0
2	G	20	Total O 20 20	0	0
2	H	15	Total O 15 15	0	0
2	I	16	Total O 16 16	0	0
2	J	17	Total O 17 17	0	0
2	K	16	Total O 16 16	0	0
2	L	11	Total O 11 11	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PROTEASOME-ACTIVATING NUCLEOTIDASE




- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PROTEASOME-ACTIVATING NUCLEOTIDASE



- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PROTEASOME-ACTIVATING NUCLEOTIDASE



- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PROTEASOME-ACTIVATING NUCLEOTIDASE



- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PROTEASOME-ACTIVATING NUCLEOTIDASE





- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PROTEASOME-ACTIVATING NUCLEOTIDASE




- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PROTEASOME-ACTIVATING NUCLEOTIDASE




- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PROTEASOME-ACTIVATING NUCLEOTIDASE



- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PROTEASOME-ACTIVATING NUCLEOTIDASE




- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PROTEASOME-ACTIVATING NUCLEOTIDASE




- Molecule 1: GENERAL CONTROL PROTEIN GCN4, PROTEASOME-ACTIVATING NUCLEOTIDASE





● Molecule 1: GENERAL CONTROL PROTEIN GCN4, PROTEASOME-ACTIVATING NUCLEOTIDASE





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	103.35Å 91.38Å 103.36Å 90.00° 119.97° 90.00°	Depositor
Resolution (Å)	34.24 – 2.50 34.24 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (34.24-2.50) 99.5 (34.24-2.50)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.21 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.196 , 0.222 0.198 , 0.220	Depositor DCC
$R_{free}$ test set	2889 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	57.9	Xtriage
Anisotropy	0.270	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 36.0	EDS
Estimated twinning fraction	0.480 for -h-l,k,h 0.480 for l,k,-h-l 0.056 for h,-k,-h-l 0.057 for -h-l,-k,l 0.057 for l,-k,h	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 57981 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8199	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 29.56 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.5166e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.84	0/673	0.84	0/913
1	B	0.86	0/671	0.84	0/912
1	C	0.85	0/673	0.84	0/913
1	D	0.86	0/675	0.84	0/916
1	E	0.84	0/673	0.84	0/913
1	F	0.86	0/675	0.85	0/916
1	G	0.84	0/681	0.84	1/924 (0.1%)
1	H	0.85	0/663	0.84	0/902
1	I	0.83	0/681	0.84	1/924 (0.1%)
1	J	0.83	0/663	0.86	0/902
1	K	0.82	0/681	0.84	1/924 (0.1%)
1	L	0.82	0/663	0.86	0/902
All	All	0.84	0/8072	0.84	3/10961 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	I	64	LEU	CB-CG-CD2	-5.90	100.97	111.00
1	G	64	LEU	CB-CG-CD2	-5.70	101.31	111.00
1	K	64	LEU	CB-CG-CD2	-5.35	101.90	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	666	0	694	1	0
1	B	664	0	680	7	0
1	C	666	0	694	1	0
1	D	668	0	691	6	0
1	E	666	0	694	1	0
1	F	668	0	691	7	0
1	G	674	0	695	7	0
1	H	656	0	670	11	0
1	I	674	0	695	9	0
1	J	656	0	670	12	0
1	K	674	0	695	9	0
1	L	656	0	670	11	0
2	A	21	0	0	0	0
2	B	22	0	0	0	0
2	C	18	0	0	0	0
2	D	19	0	0	0	0
2	E	16	0	0	0	0
2	F	20	0	0	0	0
2	G	20	0	0	0	0
2	H	15	0	0	0	0
2	I	16	0	0	0	0
2	J	17	0	0	0	0
2	K	16	0	0	0	0
2	L	11	0	0	0	0
All	All	8199	0	8239	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:64:LEU:HD11	1:I:83:THR:HG23	1.78	0.64
1:G:64:LEU:HD11	1:G:83:THR:HG23	1.79	0.63
1:F:95:ILE:HD11	1:F:115:ILE:HG22	1.81	0.62
1:H:96:ASN:HB3	1:H:99:GLU:OE2	1.99	0.61
1:J:96:ASN:HB3	1:J:99:GLU:OE2	2.00	0.61
1:L:96:ASN:HB3	1:L:99:GLU:OE2	2.02	0.60
1:K:64:LEU:HD11	1:K:83:THR:HG23	1.83	0.60
1:B:95:ILE:HD11	1:B:115:ILE:HG22	1.86	0.58
1:G:34:MET:HE2	1:H:34:MET:HA	1.88	0.56
1:D:95:ILE:HD11	1:D:115:ILE:HG22	1.88	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:64:LEU:HD21	1:D:83:THR:HG23	1.90	0.54
1:H:64:LEU:HD11	1:H:83:THR:HG23	1.90	0.53
1:K:34:MET:HE2	1:L:34:MET:HA	1.90	0.53
1:I:34:MET:HE2	1:J:34:MET:HA	1.90	0.53
1:B:64:LEU:HD21	1:B:83:THR:HG23	1.90	0.53
1:F:64:LEU:HD21	1:F:83:THR:HG23	1.91	0.53
1:G:116:VAL:O	1:G:117:ASN:ND2	2.42	0.53
1:B:45:LEU:HD13	1:H:45:LEU:CD1	2.40	0.52
1:J:64:LEU:HD11	1:J:83:THR:HG23	1.91	0.52
1:L:64:LEU:HD11	1:L:83:THR:HG23	1.92	0.51
1:F:45:LEU:HD13	1:J:45:LEU:CD1	2.40	0.51
1:K:116:VAL:O	1:K:117:ASN:ND2	2.44	0.51
1:G:54:GLU:OE1	1:G:54:GLU:HA	2.12	0.49
1:K:54:GLU:OE1	1:K:54:GLU:HA	2.12	0.49
1:J:96:ASN:HD22	1:J:99:GLU:HG3	1.79	0.48
1:I:116:VAL:O	1:I:117:ASN:ND2	2.46	0.47
1:H:96:ASN:HD22	1:H:99:GLU:HG3	1.78	0.47
1:D:45:LEU:HD13	1:L:45:LEU:CD1	2.43	0.47
1:I:54:GLU:OE1	1:I:54:GLU:HA	2.15	0.47
1:L:96:ASN:HD22	1:L:99:GLU:HG3	1.80	0.46
1:G:63:LEU:HB2	1:H:88:VAL:HB	1.99	0.45
1:B:45:LEU:HD13	1:H:45:LEU:HD12	1.99	0.45
1:K:63:LEU:HB2	1:L:88:VAL:HB	1.98	0.45
1:F:45:LEU:HD13	1:J:45:LEU:HD12	1.98	0.45
1:I:80:LYS:NZ	1:I:84:GLY:O	2.45	0.45
1:L:95:ILE:CD1	1:L:115:ILE:HG22	2.48	0.44
1:H:95:ILE:CD1	1:H:115:ILE:HG22	2.48	0.44
1:F:87:PHE:CD1	1:F:113:LEU:HD13	2.53	0.44
1:I:108:LEU:HD22	1:I:113:LEU:O	2.18	0.43
1:F:78:VAL:HG22	1:F:88:VAL:HG22	2.01	0.43
1:I:63:LEU:HB2	1:J:88:VAL:HB	2.01	0.43
1:B:87:PHE:CD1	1:B:113:LEU:HD13	2.52	0.43
1:K:83:THR:HG22	1:L:85:PRO:CG	2.48	0.42
1:K:83:THR:HG22	1:L:85:PRO:HG3	2.00	0.42
1:J:109:ASN:O	1:J:113:LEU:HA	2.19	0.42
1:D:87:PHE:CD1	1:D:113:LEU:HD13	2.54	0.42
1:L:109:ASN:O	1:L:113:LEU:HA	2.19	0.42
1:B:78:VAL:HG22	1:B:88:VAL:HG22	2.01	0.42
1:B:79:VAL:HG23	1:B:79:VAL:O	2.20	0.42
1:H:109:ASN:O	1:H:113:LEU:HA	2.19	0.42
1:A:58:LEU:HA	1:A:58:LEU:HD23	1.87	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:64:LEU:HD11	1:C:83:THR:HG23	2.01	0.41
1:D:64:LEU:HD11	1:D:83:THR:HG23	2.02	0.41
1:F:79:VAL:HG23	1:F:79:VAL:O	2.21	0.41
1:K:80:LYS:NZ	1:K:84:GLY:O	2.45	0.41
1:D:45:LEU:HD13	1:L:45:LEU:HD12	2.02	0.41
1:E:64:LEU:HD11	1:E:83:THR:HG23	2.01	0.41
1:G:83:THR:HG22	1:H:85:PRO:CG	2.50	0.41
1:I:83:THR:HG22	1:J:85:PRO:CG	2.51	0.41
1:I:83:THR:HG22	1:J:85:PRO:HG3	2.02	0.40
1:G:83:THR:HG22	1:H:85:PRO:HG3	2.02	0.40
1:J:95:ILE:CD1	1:J:115:ILE:HG22	2.51	0.40
1:K:95:ILE:O	1:K:95:ILE:HG22	2.21	0.40
1:J:71:ILE:HG22	1:J:72:LEU:N	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	85/109 (78%)	80 (94%)	5 (6%)	0	100	100
1	B	85/109 (78%)	83 (98%)	2 (2%)	0	100	100
1	C	85/109 (78%)	81 (95%)	4 (5%)	0	100	100
1	D	85/109 (78%)	83 (98%)	2 (2%)	0	100	100
1	E	85/109 (78%)	81 (95%)	4 (5%)	0	100	100
1	F	85/109 (78%)	84 (99%)	1 (1%)	0	100	100
1	G	85/109 (78%)	83 (98%)	2 (2%)	0	100	100
1	H	85/109 (78%)	83 (98%)	2 (2%)	0	100	100
1	I	85/109 (78%)	83 (98%)	2 (2%)	0	100	100
1	J	85/109 (78%)	82 (96%)	3 (4%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	K	85/109 (78%)	83 (98%)	2 (2%)	0	100	100
1	L	85/109 (78%)	82 (96%)	3 (4%)	0	100	100
All	All	1020/1308 (78%)	988 (97%)	32 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	75/99 (76%)	72 (96%)	3 (4%)	38	64
1	B	74/99 (75%)	69 (93%)	5 (7%)	20	36
1	C	75/99 (76%)	72 (96%)	3 (4%)	38	64
1	D	75/99 (76%)	69 (92%)	6 (8%)	15	28
1	E	75/99 (76%)	72 (96%)	3 (4%)	38	64
1	F	75/99 (76%)	69 (92%)	6 (8%)	15	28
1	G	77/99 (78%)	75 (97%)	2 (3%)	54	81
1	H	72/99 (73%)	69 (96%)	3 (4%)	36	62
1	I	77/99 (78%)	75 (97%)	2 (3%)	54	81
1	J	72/99 (73%)	69 (96%)	3 (4%)	36	62
1	K	77/99 (78%)	75 (97%)	2 (3%)	54	81
1	L	72/99 (73%)	68 (94%)	4 (6%)	26	47
All	All	896/1188 (75%)	854 (95%)	42 (5%)	32	56

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	MET
1	A	76	ARG
1	A	95	ILE
1	B	60	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	71	ILE
1	B	100	LEU
1	B	110	GLN
1	B	119	LEU
1	C	34	MET
1	C	76	ARG
1	C	95	ILE
1	D	35	LYS
1	D	60	SER
1	D	71	ILE
1	D	100	LEU
1	D	110	GLN
1	D	119	LEU
1	E	34	MET
1	E	76	ARG
1	E	95	ILE
1	F	35	LYS
1	F	60	SER
1	F	71	ILE
1	F	100	LEU
1	F	110	GLN
1	F	119	LEU
1	G	34	MET
1	G	70	ASP
1	H	47	LYS
1	H	95	ILE
1	H	119	LEU
1	I	34	MET
1	I	70	ASP
1	J	47	LYS
1	J	95	ILE
1	J	119	LEU
1	K	34	MET
1	K	70	ASP
1	L	47	LYS
1	L	71	ILE
1	L	95	ILE
1	L	119	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	ASN
1	B	48	ASN
1	B	110	GLN
1	C	53	ASN
1	D	48	ASN
1	D	110	GLN
1	E	53	ASN
1	F	48	ASN
1	F	110	GLN
1	G	53	ASN
1	G	117	ASN
1	H	96	ASN
1	I	53	ASN
1	I	117	ASN
1	J	96	ASN
1	K	53	ASN
1	K	117	ASN
1	L	96	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.



## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	87/109 (79%)	0.25	3 (3%) 49 54	37, 53, 83, 99	0
1	B	87/109 (79%)	0.19	2 (2%) 64 67	36, 52, 83, 91	0
1	C	87/109 (79%)	0.24	1 (1%) 82 84	37, 53, 83, 99	0
1	D	87/109 (79%)	0.18	0 100 100	36, 52, 83, 92	0
1	E	87/109 (79%)	0.25	0 100 100	37, 53, 83, 99	0
1	F	87/109 (79%)	0.23	1 (1%) 82 84	36, 52, 83, 92	0
1	G	87/109 (79%)	0.17	2 (2%) 64 67	42, 57, 92, 109	0
1	H	87/109 (79%)	0.20	1 (1%) 82 84	40, 57, 90, 111	0
1	I	87/109 (79%)	0.22	2 (2%) 64 67	42, 57, 92, 109	0
1	J	87/109 (79%)	0.16	1 (1%) 82 84	40, 57, 90, 111	0
1	K	87/109 (79%)	0.27	3 (3%) 49 54	42, 58, 92, 109	0
1	L	87/109 (79%)	0.21	1 (1%) 82 84	40, 57, 90, 111	0
All	All	1044/1308 (79%)	0.21	17 (1%) 74 78	36, 56, 90, 111	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	37	LEU	4.7
1	K	37	LEU	4.2
1	K	34	MET	4.0
1	G	37	LEU	3.4
1	I	34	MET	3.1
1	B	37	LEU	3.0
1	H	37	LEU	3.0
1	I	37	LEU	3.0
1	J	37	LEU	2.8
1	K	35	LYS	2.7
1	B	34	MET	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	34	MET	2.7
1	C	34	MET	2.4
1	G	34	MET	2.4
1	A	37	LEU	2.4
1	F	34	MET	2.2
1	A	100	LEU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.