



wwPDB X-ray Structure Validation Summary Report i

Feb 19, 2016 – 08:13 PM GMT

PDB ID : 4WGL
Title : Crystal structure of a GroEL D83A/R197A double mutant
Authors : Yang, D.; Fei, X.; LaRonde, N.A.; Beckett, D.; Lund, P.A.; Lorimer, G.H.
Deposited on : 2014-09-19
Resolution : 3.13 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026982
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	1095 (3.18-3.10)
Clashscore	102246	1202 (3.18-3.10)
Ramachandran outliers	100387	1162 (3.18-3.10)
Sidechain outliers	100360	1162 (3.18-3.10)
RSRZ outliers	91569	1097 (3.18-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain			
1	F	548	%	67%	26%	• •
1	G	548	4%	74%	21%	• •
1	H	548		74%	20%	• •
1	I	548		70%	24%	• •
1	J	548		70%	24%	• •
1	K	548	9%	71%	23%	• •
1	L	548	%	72%	22%	• •
1	M	548	%	72%	22%	• •
1	N	548	%	77%	16%	• •

2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 53844 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 60 kDa chaperonin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	524	Total	C	N	O	S	0	0	0
			3846	2393	662	771	20			
1	B	524	Total	C	N	O	S	0	0	0
			3846	2393	662	771	20			
1	C	524	Total	C	N	O	S	0	0	0
			3846	2393	662	771	20			
1	D	524	Total	C	N	O	S	0	0	0
			3846	2393	662	771	20			
1	E	524	Total	C	N	O	S	0	0	0
			3846	2393	662	771	20			
1	F	524	Total	C	N	O	S	0	0	0
			3846	2393	662	771	20			
1	G	524	Total	C	N	O	S	0	0	0
			3846	2393	662	771	20			
1	H	524	Total	C	N	O	S	0	0	0
			3846	2393	662	771	20			
1	I	524	Total	C	N	O	S	0	0	0
			3846	2393	662	771	20			
1	J	524	Total	C	N	O	S	0	0	0
			3846	2393	662	771	20			
1	K	524	Total	C	N	O	S	0	0	0
			3846	2393	662	771	20			
1	L	524	Total	C	N	O	S	0	0	0
			3846	2393	662	771	20			
1	M	524	Total	C	N	O	S	0	0	0
			3846	2393	662	771	20			
1	N	524	Total	C	N	O	S	0	0	0
			3846	2393	662	771	20			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	83	ALA	ASP	engineered mutation	UNP P0A6F5

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	197	ALA	ARG	engineered mutation	UNP P0A6F5
B	83	ALA	ASP	engineered mutation	UNP P0A6F5
B	197	ALA	ARG	engineered mutation	UNP P0A6F5
C	83	ALA	ASP	engineered mutation	UNP P0A6F5
C	197	ALA	ARG	engineered mutation	UNP P0A6F5
D	83	ALA	ASP	engineered mutation	UNP P0A6F5
D	197	ALA	ARG	engineered mutation	UNP P0A6F5
E	83	ALA	ASP	engineered mutation	UNP P0A6F5
E	197	ALA	ARG	engineered mutation	UNP P0A6F5
F	83	ALA	ASP	engineered mutation	UNP P0A6F5
F	197	ALA	ARG	engineered mutation	UNP P0A6F5
G	83	ALA	ASP	engineered mutation	UNP P0A6F5
G	197	ALA	ARG	engineered mutation	UNP P0A6F5
H	83	ALA	ASP	engineered mutation	UNP P0A6F5
H	197	ALA	ARG	engineered mutation	UNP P0A6F5
I	83	ALA	ASP	engineered mutation	UNP P0A6F5
I	197	ALA	ARG	engineered mutation	UNP P0A6F5
J	83	ALA	ASP	engineered mutation	UNP P0A6F5
J	197	ALA	ARG	engineered mutation	UNP P0A6F5
K	83	ALA	ASP	engineered mutation	UNP P0A6F5
K	197	ALA	ARG	engineered mutation	UNP P0A6F5
L	83	ALA	ASP	engineered mutation	UNP P0A6F5
L	197	ALA	ARG	engineered mutation	UNP P0A6F5
M	83	ALA	ASP	engineered mutation	UNP P0A6F5
M	197	ALA	ARG	engineered mutation	UNP P0A6F5
N	83	ALA	ASP	engineered mutation	UNP P0A6F5
N	197	ALA	ARG	engineered mutation	UNP P0A6F5




3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 60 kDa chaperonin




- Molecule 1: 60 kDa chaperonin




- Molecule 1: 60 kDa chaperonin






- Molecule 1: 60 kDa chaperonin

A horizontal bar chart illustrating the distribution of Chain D. The total length of the bar is divided into three segments: a small red segment at the left end labeled '2%', a long green segment in the middle labeled '72%', and a yellow segment on the right labeled '22%'. The segments are separated by thin black lines.



- Molecule 1: 60 kDa chaperonin


Chain E: 75% 18% • •




- Molecule 1: 60 kDa chaperonin

A horizontal bar chart titled "Chain F:" at the top left. The bar is divided into three segments: a red segment at the far left labeled with a percentage symbol (%), a long green segment labeled "67%", and a yellow segment on the right labeled "26%". The total length of the bar represents 100%.


- Molecule 1: 60 kDa chaperonin




- Molecule 1: 60 kDa chaperonin




- Molecule 1: 60 kDa chaperonin






- Molecule 1: 60 kDa chaperonin

Chain L: 72% 22% • •



- Molecule 1: 60 kDa chaperonin


Chain M: 72% 22% • •



- Molecule 1: 60 kDa chaperonin

Chain N: 77% 16% • •





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	135.62 Å 259.71 Å 280.85 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	123.52 – 3.13 123.52 – 3.13	Depositor EDS
% Data completeness (in resolution range)	99.6 (123.52-3.13) 99.6 (123.52-3.13)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle^1$	2.65 (at 3.13 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R , R_{free}	0.166 , 0.233 0.166 , 0.230	Depositor DCC
R_{free} test set	2001 reflections (1.15%)	DCC
Wilson B-factor (Å ²)	69.5	Xtriage
Anisotropy	0.402	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 70.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 174049 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	53844	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.43	0/3874	0.61	0/5232
1	B	0.43	0/3874	0.62	0/5232
1	C	0.42	0/3874	0.60	0/5232
1	D	0.42	0/3874	0.59	1/5232 (0.0%)
1	E	0.42	0/3874	0.61	0/5232
1	F	0.41	0/3874	0.59	0/5232
1	G	0.41	0/3874	0.58	0/5232
1	H	0.45	0/3874	0.62	0/5232
1	I	0.46	0/3874	0.66	1/5232 (0.0%)
1	J	0.46	0/3874	0.63	1/5232 (0.0%)
1	K	0.43	0/3874	0.61	1/5232 (0.0%)
1	L	0.47	0/3874	0.64	0/5232
1	M	0.41	0/3874	0.59	1/5232 (0.0%)
1	N	0.43	0/3874	0.61	0/5232
All	All	0.43	0/54236	0.61	5/73248 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1
1	J	0	1
All	All	0	2

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	I	295	LEU	CA-CB-CG	6.30	129.78	115.30
1	M	183	LEU	CA-CB-CG	5.83	128.70	115.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	J	131	LEU	CA-CB-CG	5.14	127.11	115.30
1	K	182	GLY	N-CA-C	5.13	125.94	113.10
1	D	383	ALA	N-CA-C	5.11	124.78	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	383	ALA	Peptide
1	J	31	LEU	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3846	0	3969	59	0
1	B	3846	0	3969	83	0
1	C	3846	0	3969	73	0
1	D	3846	0	3969	78	0
1	E	3846	0	3969	64	0
1	F	3846	0	3969	89	0
1	G	3846	0	3969	68	0
1	H	3846	0	3969	71	0
1	I	3846	0	3969	88	0
1	J	3846	0	3969	79	0
1	K	3846	0	3969	87	0
1	L	3846	0	3969	72	0
1	M	3846	0	3969	70	0
1	N	3846	0	3969	54	0
All	All	53844	0	55566	986	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 986 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:232:GLU:HB3	1:K:309:LEU:HD11	1.46	0.95
1:F:241:ALA:HA	1:F:271:VAL:HG11	1.51	0.92
1:L:130:GLU:HB3	1:L:422:VAL:HG22	1.51	0.91
1:E:173:GLY:O	1:E:404:ARG:NH2	2.08	0.86
1:E:524:LEU:HD12	1:E:525:PRO:HD2	1.56	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	522/548 (95%)	512 (98%)	8 (2%)	2 (0%)	39 77
1	B	522/548 (95%)	510 (98%)	10 (2%)	2 (0%)	39 77
1	C	522/548 (95%)	510 (98%)	10 (2%)	2 (0%)	39 77
1	D	522/548 (95%)	511 (98%)	10 (2%)	1 (0%)	52 85
1	E	522/548 (95%)	510 (98%)	10 (2%)	2 (0%)	39 77
1	F	522/548 (95%)	511 (98%)	10 (2%)	1 (0%)	52 85
1	G	522/548 (95%)	512 (98%)	9 (2%)	1 (0%)	52 85
1	H	522/548 (95%)	514 (98%)	8 (2%)	0	100 100
1	I	522/548 (95%)	507 (97%)	13 (2%)	2 (0%)	39 77
1	J	522/548 (95%)	514 (98%)	7 (1%)	1 (0%)	52 85
1	K	522/548 (95%)	506 (97%)	14 (3%)	2 (0%)	39 77
1	L	522/548 (95%)	511 (98%)	10 (2%)	1 (0%)	52 85
1	M	522/548 (95%)	507 (97%)	14 (3%)	1 (0%)	52 85
1	N	522/548 (95%)	513 (98%)	8 (2%)	1 (0%)	52 85
All	All	7308/7672 (95%)	7148 (98%)	141 (2%)	19 (0%)	46 81

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	271	VAL
1	J	271	VAL
1	M	271	VAL
1	N	271	VAL
1	A	183	LEU

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	402/413 (97%)	379 (94%)	23 (6%)	25 62
1	B	402/413 (97%)	382 (95%)	20 (5%)	30 67
1	C	402/413 (97%)	382 (95%)	20 (5%)	30 67
1	D	402/413 (97%)	380 (94%)	22 (6%)	27 63
1	E	402/413 (97%)	381 (95%)	21 (5%)	29 65
1	F	402/413 (97%)	380 (94%)	22 (6%)	27 63
1	G	402/413 (97%)	385 (96%)	17 (4%)	36 73
1	H	402/413 (97%)	383 (95%)	19 (5%)	32 70
1	I	402/413 (97%)	376 (94%)	26 (6%)	21 57
1	J	402/413 (97%)	377 (94%)	25 (6%)	23 59
1	K	402/413 (97%)	383 (95%)	19 (5%)	32 70
1	L	402/413 (97%)	376 (94%)	26 (6%)	21 57
1	M	402/413 (97%)	379 (94%)	23 (6%)	25 62
1	N	402/413 (97%)	375 (93%)	27 (7%)	20 55
All	All	5628/5782 (97%)	5318 (94%)	310 (6%)	27 63

5 of 310 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	331	THR
1	I	138	CYS
1	N	43	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	G	494	LEU
1	H	328	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	326	ASN
1	M	433	ASN
1	H	479	ASN
1	B	206	ASN
1	J	146	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	524/548 (95%)	-0.18	2 (0%)	93	86	22, 60, 120, 174	0
1	B	524/548 (95%)	0.17	30 (5%)	27	11	24, 65, 207, 301	0
1	C	524/548 (95%)	-0.12	5 (0%)	84	70	25, 70, 166, 204	0
1	D	524/548 (95%)	-0.02	12 (2%)	64	42	28, 73, 161, 232	0
1	E	524/548 (95%)	-0.15	1 (0%)	95	91	28, 69, 136, 190	0
1	F	524/548 (95%)	-0.07	4 (0%)	87	76	31, 77, 135, 190	0
1	G	524/548 (95%)	0.09	21 (4%)	42	21	27, 80, 167, 208	0
1	H	524/548 (95%)	-0.25	0	100	100	25, 63, 118, 168	0
1	I	524/548 (95%)	-0.13	2 (0%)	93	86	21, 54, 146, 227	0
1	J	524/548 (95%)	-0.16	1 (0%)	95	91	18, 63, 134, 189	0
1	K	524/548 (95%)	0.29	47 (8%)	12	4	22, 66, 239, 292	0
1	L	524/548 (95%)	-0.23	1 (0%)	95	91	25, 58, 125, 177	0
1	M	524/548 (95%)	-0.04	7 (1%)	79	63	28, 77, 172, 233	0
1	N	524/548 (95%)	-0.12	3 (0%)	90	81	22, 64, 152, 207	0
All	All	7336/7672 (95%)	-0.07	136 (1%)	70	49	18, 66, 167, 301	0

The worst 5 of 136 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	K	223	ALA	12.0
1	B	251	ALA	8.8
1	K	301	ILE	7.6
1	K	203	TYR	7.3
1	K	251	ALA	7.2

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.