



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 06:51 PM GMT

PDB ID : 3WUT  
Title : Structure basis of inactivating cell abscission  
Authors : Kim, H.J.; Matsuura, A.; Lee, H.H.  
Deposited on : 2014-05-05  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	63	
1	B	63	
1	D	63	
1	E	63	
1	G	63	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	H	63	 65% 14% 21%
1	J	63	 13% 59% 11% 30%
1	K	63	 11% 62% 6% 32%
2	C	14	 7% 86% 7% 7%
2	F	14	 93% 7%
2	I	14	 86% 7% 7%
2	L	14	 93% 7%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3906 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Centrosomal protein of 55 kDa.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	53	Total 446	C 284	N 74	O 87	S 1	0	0	0
1	B	44	Total 377	C 242	N 62	O 72	S 1	0	0	0
1	D	54	Total 447	C 285	N 75	O 85	S 2	0	0	0
1	E	52	Total 434	C 277	N 73	O 83	S 1	0	0	0
1	G	55	Total 456	C 290	N 76	O 88	S 2	0	0	0
1	H	50	Total 423	C 271	N 71	O 80	S 1	0	0	0
1	J	44	Total 377	C 242	N 62	O 72	S 1	0	0	0
1	K	43	Total 370	C 239	N 62	O 68	S 1	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	155	GLY	-	EXPRESSION TAG	UNP Q53EZ4
A	156	ALA	-	EXPRESSION TAG	UNP Q53EZ4
A	157	MET	-	EXPRESSION TAG	UNP Q53EZ4
A	158	GLY	-	EXPRESSION TAG	UNP Q53EZ4
A	159	SER	-	EXPRESSION TAG	UNP Q53EZ4
B	155	GLY	-	EXPRESSION TAG	UNP Q53EZ4
B	156	ALA	-	EXPRESSION TAG	UNP Q53EZ4
B	157	MET	-	EXPRESSION TAG	UNP Q53EZ4
B	158	GLY	-	EXPRESSION TAG	UNP Q53EZ4
B	159	SER	-	EXPRESSION TAG	UNP Q53EZ4
D	155	GLY	-	EXPRESSION TAG	UNP Q53EZ4
D	156	ALA	-	EXPRESSION TAG	UNP Q53EZ4
D	157	MET	-	EXPRESSION TAG	UNP Q53EZ4

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	158	GLY	-	EXPRESSION TAG	UNP Q53EZ4
D	159	SER	-	EXPRESSION TAG	UNP Q53EZ4
E	155	GLY	-	EXPRESSION TAG	UNP Q53EZ4
E	156	ALA	-	EXPRESSION TAG	UNP Q53EZ4
E	157	MET	-	EXPRESSION TAG	UNP Q53EZ4
E	158	GLY	-	EXPRESSION TAG	UNP Q53EZ4
E	159	SER	-	EXPRESSION TAG	UNP Q53EZ4
G	155	GLY	-	EXPRESSION TAG	UNP Q53EZ4
G	156	ALA	-	EXPRESSION TAG	UNP Q53EZ4
G	157	MET	-	EXPRESSION TAG	UNP Q53EZ4
G	158	GLY	-	EXPRESSION TAG	UNP Q53EZ4
G	159	SER	-	EXPRESSION TAG	UNP Q53EZ4
H	155	GLY	-	EXPRESSION TAG	UNP Q53EZ4
H	156	ALA	-	EXPRESSION TAG	UNP Q53EZ4
H	157	MET	-	EXPRESSION TAG	UNP Q53EZ4
H	158	GLY	-	EXPRESSION TAG	UNP Q53EZ4
H	159	SER	-	EXPRESSION TAG	UNP Q53EZ4
J	155	GLY	-	EXPRESSION TAG	UNP Q53EZ4
J	156	ALA	-	EXPRESSION TAG	UNP Q53EZ4
J	157	MET	-	EXPRESSION TAG	UNP Q53EZ4
J	158	GLY	-	EXPRESSION TAG	UNP Q53EZ4
J	159	SER	-	EXPRESSION TAG	UNP Q53EZ4
K	155	GLY	-	EXPRESSION TAG	UNP Q53EZ4
K	156	ALA	-	EXPRESSION TAG	UNP Q53EZ4
K	157	MET	-	EXPRESSION TAG	UNP Q53EZ4
K	158	GLY	-	EXPRESSION TAG	UNP Q53EZ4
K	159	SER	-	EXPRESSION TAG	UNP Q53EZ4


- Molecule 2 is a protein called Inactive serine/threonine-protein kinase TEX14.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	13	Total	C	N	O	0	0	0
			94	64	14	16			
2	F	13	Total	C	N	O	0	0	0
			94	64	14	16			
2	I	13	Total	C	N	O	0	0	0
			94	64	14	16			
2	L	13	Total	C	N	O	0	0	0
			94	64	14	16			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	791	ASP	-	EXPRESSION TAG	UNP Q8IWB6
F	791	ASP	-	EXPRESSION TAG	UNP Q8IWB6
I	791	ASP	-	EXPRESSION TAG	UNP Q8IWB6
L	791	ASP	-	EXPRESSION TAG	UNP Q8IWB6

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O 6 3 3	0	0
3	F	1	Total C O 6 3 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	18	Total O 18 18	0	0
4	B	30	Total O 30 30	0	0
4	C	3	Total O 3 3	0	0
4	D	20	Total O 20 20	0	0
4	E	34	Total O 34 34	0	0
4	F	4	Total O 4 4	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	G	11	Total 11	O 11	0	0
4	H	26	Total 26	O 26	0	0
4	I	9	Total 9	O 9	0	0
4	J	13	Total 13	O 13	0	0
4	K	16	Total 16	O 16	0	0
4	L	4	Total 4	O 4	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Centrosomal protein of 55 kDa




- Molecule 1: Centrosomal protein of 55 kDa




- Molecule 1: Centrosomal protein of 55 kDa



- Molecule 1: Centrosomal protein of 55 kDa



- Molecule 1: Centrosomal protein of 55 kDa



- Molecule 1: Centrosomal protein of 55 kDa

Chain H:  65% 14% 21%



- Molecule 1: Centrosomal protein of 55 kDa

Chain J:  13% 59% 11% 30%




- Molecule 1: Centrosomal protein of 55 kDa

Chain K:  11% 62% 6% 32%



- Molecule 2: Inactive serine/threonine-protein kinase TEX14

Chain C:  7% 86% 7% 7%



- Molecule 2: Inactive serine/threonine-protein kinase TEX14

Chain F:  93% 7%



- Molecule 2: Inactive serine/threonine-protein kinase TEX14

Chain I:  86% 7% 7%



- Molecule 2: Inactive serine/threonine-protein kinase TEX14

Chain L:  93% 7%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.91Å 102.62Å 132.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.88 – 2.30 28.88 – 2.30	Depositor EDS
% Data completeness (in resolution range)	94.3 (28.88-2.30) 94.3 (28.88-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.22 (at 2.31Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.220 , 0.259 0.218 , 0.258	Depositor DCC
$R_{free}$ test set	1588 reflections (5.04%)	DCC
Wilson B-factor (Å <sup>2</sup> )	43.1	Xtrriage
Anisotropy	0.014	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 52.4	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Outliers	0 of 31522 reflections	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3906	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.36	0/452	0.47	0/606
1	B	0.33	0/382	0.50	0/511
1	D	0.32	0/453	0.45	0/606
1	E	0.35	0/440	0.44	0/589
1	G	0.33	0/462	0.42	0/618
1	H	0.31	0/429	0.46	0/574
1	J	0.37	0/382	0.46	0/511
1	K	0.44	0/375	0.55	0/501
2	C	0.29	0/98	0.57	0/137
2	F	0.32	0/98	0.65	0/137
2	I	0.33	0/98	0.63	0/137
2	L	0.28	0/98	0.56	0/137
All	All	0.35	0/3767	0.48	0/5064

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	446	0	445	6	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	377	0	381	7	0
1	D	447	0	449	2	0
1	E	434	0	435	2	0
1	G	456	0	455	9	0
1	H	423	0	425	8	0
1	J	377	0	381	4	0
1	K	370	0	378	3	0
2	C	94	0	97	0	0
2	F	94	0	97	0	0
2	I	94	0	97	1	0
2	L	94	0	97	0	0
3	B	6	0	8	2	0
3	F	6	0	8	0	0
4	A	18	0	0	1	0
4	B	30	0	0	0	0
4	C	3	0	0	0	0
4	D	20	0	0	0	0
4	E	34	0	0	0	0
4	F	4	0	0	0	0
4	G	11	0	0	0	0
4	H	26	0	0	1	0
4	I	9	0	0	0	0
4	J	13	0	0	0	0
4	K	16	0	0	1	0
4	L	4	0	0	0	0
All	All	3906	0	3753	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:202:ILE:HG23	1:H:201:LYS:HZ1	1.60	0.67
4:A:313:HOH:O	1:D:182:GLN:HG3	1.95	0.67
1:G:157:MET:HG3	1:G:159:SER:H	1.63	0.63
1:A:182:GLN:NE2	1:G:208:LYS:HG3	2.17	0.60
3:B:301:GOL:H11	2:I:803:PRO:HB2	1.85	0.59
1:K:171:GLU:O	1:K:175:LYS:HG2	2.03	0.59
1:D:206:GLU:O	1:D:209:THR:HG22	2.04	0.58
1:B:182:GLN:HA	1:B:185:LEU:HD22	1.88	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:202:ILE:HG12	1:H:201:LYS:NZ	2.21	0.55
1:B:209:THR:HA	1:B:210:GLU:C	2.28	0.54
1:B:187:TYR:CD1	1:G:197:GLY:HA3	2.43	0.54
1:G:202:ILE:CG2	1:H:201:LYS:HZ1	2.21	0.53
1:A:196:LYS:NZ	3:B:301:GOL:H31	2.25	0.52
1:J:208:LYS:O	1:J:210:GLU:HG2	2.11	0.51
1:A:174:LEU:HD12	1:B:170:MET:HE3	1.93	0.51
1:G:160:PHE:O	1:G:164:ILE:HG23	2.11	0.51
1:G:167:ILE:HD13	1:H:167:ILE:HD12	1.92	0.50
1:E:160:PHE:O	1:E:164:ILE:HG13	2.12	0.49
1:A:170:MET:HG2	1:B:170:MET:HE1	1.94	0.49
1:J:168:HIS:O	1:J:168:HIS:ND1	2.46	0.49
1:K:171:GLU:HB2	4:K:314:HOH:O	2.13	0.47
1:A:182:GLN:HE22	1:G:208:LYS:HG3	1.80	0.46
1:A:209:THR:HG22	1:B:209:THR:OG1	2.16	0.45
1:J:169:GLU:HB3	1:J:172:ILE:HD12	1.99	0.44
1:E:168:HIS:O	1:E:172:ILE:HG12	2.18	0.44
1:H:183:GLN:OE1	4:H:316:HOH:O	2.21	0.43
1:H:163:SER:O	1:H:167:ILE:HG12	2.20	0.42
1:H:168:HIS:O	1:H:172:ILE:HG12	2.20	0.41
1:B:210:GLU:HB3	1:B:211:THR:H	1.53	0.41
1:H:187:TYR:CD1	1:J:197:GLY:HA3	2.56	0.41
1:K:182:GLN:O	1:K:186:VAL:HG23	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	51/63 (81%)	51 (100%)	0	0	100	100
1	B	42/63 (67%)	41 (98%)	1 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	52/63 (82%)	52 (100%)	0	0	100	100
1	E	50/63 (79%)	50 (100%)	0	0	100	100
1	G	53/63 (84%)	53 (100%)	0	0	100	100
1	H	48/63 (76%)	48 (100%)	0	0	100	100
1	J	42/63 (67%)	40 (95%)	1 (2%)	1 (2%)	7	5
1	K	41/63 (65%)	39 (95%)	2 (5%)	0	100	100
2	C	11/14 (79%)	11 (100%)	0	0	100	100
2	F	11/14 (79%)	11 (100%)	0	0	100	100
2	I	11/14 (79%)	11 (100%)	0	0	100	100
2	L	11/14 (79%)	11 (100%)	0	0	100	100
All	All	423/560 (76%)	418 (99%)	4 (1%)	1 (0%)	52	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	J	209	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	50/55 (91%)	48 (96%)	2 (4%)	38	52
1	B	41/55 (74%)	38 (93%)	3 (7%)	17	22
1	D	49/55 (89%)	48 (98%)	1 (2%)	63	79
1	E	48/55 (87%)	47 (98%)	1 (2%)	61	78
1	G	50/55 (91%)	48 (96%)	2 (4%)	38	52
1	H	47/55 (86%)	45 (96%)	2 (4%)	35	47
1	J	41/55 (74%)	41 (100%)	0	100	100
1	K	40/55 (73%)	40 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	C	11/12 (92%)	10 (91%)	1 (9%)	12	13
2	F	11/12 (92%)	11 (100%)	0	100	100
2	I	11/12 (92%)	11 (100%)	0	100	100
2	L	11/12 (92%)	11 (100%)	0	100	100
All	All	410/488 (84%)	398 (97%)	12 (3%)	50	66

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	174	LEU
1	A	195	VAL
1	B	170	MET
1	B	185	LEU
1	B	211	THR
2	C	798	SER
1	D	162	SER
1	E	208	LYS
1	G	164	ILE
1	G	167	ILE
1	H	159	SER
1	H	208	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	K	166	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	B	301	-	5,5,5	0.41	0	5,5,5	0.38	0
3	GOL	F	901	-	5,5,5	0.28	0	5,5,5	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	301	-	-	0/4/4/4	0/0/0/0
3	GOL	F	901	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	301	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	53/63 (84%)	0.97	9 (16%) <span style="border: 1px solid red; padding: 2px;">2</span> <span style="border: 1px solid red; padding: 2px;">3</span>	37, 65, 86, 97	0
1	B	44/63 (69%)	1.11	15 (34%) <span style="border: 1px solid red; padding: 2px;">0</span> <span style="border: 1px solid red; padding: 2px;">0</span>	41, 62, 103, 107	0
1	D	54/63 (85%)	0.41	1 (1%) <span style="border: 1px solid blue; padding: 2px;">70</span> <span style="border: 1px solid blue; padding: 2px;">76</span>	40, 54, 78, 97	0
1	E	52/63 (82%)	0.09	1 (1%) <span style="border: 1px solid blue; padding: 2px;">70</span> <span style="border: 1px solid blue; padding: 2px;">76</span>	39, 55, 84, 102	0
1	G	55/63 (87%)	0.24	1 (1%) <span style="border: 1px solid blue; padding: 2px;">71</span> <span style="border: 1px solid blue; padding: 2px;">78</span>	46, 62, 82, 111	0
1	H	50/63 (79%)	-0.03	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	45, 58, 76, 86	0
1	J	44/63 (69%)	0.88	8 (18%) <span style="border: 1px solid red; padding: 2px;">2</span> <span style="border: 1px solid red; padding: 2px;">3</span>	41, 58, 100, 115	0
1	K	43/63 (68%)	1.00	7 (16%) <span style="border: 1px solid red; padding: 2px;">2</span> <span style="border: 1px solid red; padding: 2px;">4</span>	42, 59, 111, 129	0
2	C	13/14 (92%)	0.29	1 (7%) <span style="border: 1px solid red; padding: 2px;">16</span> <span style="border: 1px solid red; padding: 2px;">23</span>	46, 61, 80, 91	0
2	F	13/14 (92%)	0.02	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	43, 48, 61, 63	0
2	I	13/14 (92%)	-0.28	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	46, 53, 60, 67	0
2	L	13/14 (92%)	0.17	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	50, 60, 73, 86	0
All	All	447/560 (79%)	0.50	43 (9%) <span style="border: 1px solid red; padding: 2px;">10</span> <span style="border: 1px solid red; padding: 2px;">15</span>	37, 59, 97, 129	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	211	THR	7.7
1	K	166	ASN	6.5
1	B	208	LYS	5.5
1	B	210	GLU	5.2
1	K	170	MET	5.1
1	A	172	ILE	5.1
1	D	209	THR	4.7
1	B	172	ILE	4.5
1	B	209	THR	4.4
1	J	209	THR	4.1
1	B	211	THR	4.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	K	169	GLU	4.0
1	B	170	MET	4.0
1	E	209	THR	3.8
1	A	167	ILE	3.7
1	K	172	ILE	3.7
1	A	211	THR	3.7
1	K	168	HIS	3.5
1	G	210	GLU	3.4
1	A	209	THR	3.3
1	J	210	GLU	3.2
1	B	169	GLU	3.1
1	B	173	GLN	3.0
1	B	174	LEU	3.0
1	B	168	HIS	2.9
1	K	167	ILE	2.8
1	B	177	ALA	2.7
2	C	792	LEU	2.7
1	A	168	HIS	2.7
1	A	210	GLU	2.6
1	A	169	GLU	2.6
1	J	170	MET	2.6
1	B	207	LYS	2.6
1	B	185	LEU	2.5
1	A	173	GLN	2.5
1	J	172	ILE	2.4
1	B	171	GLU	2.3
1	K	171	GLU	2.3
1	J	173	GLN	2.2
1	A	159	SER	2.1
1	B	175	LYS	2.1
1	J	198	LEU	2.0
1	J	171	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	301	6/6	0.93	0.19	1.03	39,49,52,63	0
3	GOL	F	901	6/6	0.95	0.19	0.01	43,49,58,59	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.