



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:25 PM GMT

PDB ID : 3WVL
Title : Crystal structure of the football-shaped GroEL-GroES complex (GroEL: GroES2:ATP14) from Escherichia coli
Authors : Koike-Takeshita, A.; Arakawa, T.; Taguchi, H.; Shimamura, T.
Deposited on : 2014-05-23
Resolution : 3.79 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.


























Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1273 (4.06-3.50)
Clashscore	102246	1412 (4.06-3.50)
Ramachandran outliers	100387	1351 (4.06-3.50)
Sidechain outliers	100360	1347 (4.06-3.50)
RSRZ outliers	91569	1281 (4.06-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	548	<div> <div>2%</div> <div>68% 26% . .</div> </div>
1	B	548	<div> <div>%</div> <div>68% 26% . .</div> </div>
1	C	548	<div> <div>%</div> <div>67% 26% . .</div> </div>
1	D	548	<div> <div>%</div> <div>62% 30% . .</div> </div>
1	E	548	<div> <div>%</div> <div>65% 28% . .</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	548	% 
1	G	548	% 
1	H	548	% 
1	I	548	% 
1	J	548	% 
1	K	548	% 
1	L	548	% 
1	M	548	% 
1	N	548	% 
2	O	97	9% 
2	P	97	% 
2	Q	97	14% 
2	R	97	16% 
2	S	97	14% 
2	T	97	23% 
2	U	97	10% 
2	V	97	7% 
2	W	97	% 
2	X	97	13% 
2	Y	97	20% 
2	Z	97	10% 
2	a	97	13% 
2	b	97	13% 

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 64540 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 60 kDa chaperonin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	524	Total	C	N	O	S	0	0	0
			3849	2395	665	769	20			
1	B	524	Total	C	N	O	S	0	0	0
			3849	2395	665	769	20			
1	C	524	Total	C	N	O	S	0	0	0
			3849	2395	665	769	20			
1	D	524	Total	C	N	O	S	0	0	0
			3849	2395	665	769	20			
1	E	524	Total	C	N	O	S	0	0	0
			3849	2395	665	769	20			
1	F	524	Total	C	N	O	S	0	0	0
			3849	2395	665	769	20			
1	G	524	Total	C	N	O	S	0	0	0
			3849	2395	665	769	20			
1	H	524	Total	C	N	O	S	0	0	0
			3849	2395	665	769	20			
1	I	524	Total	C	N	O	S	0	0	0
			3849	2395	665	769	20			
1	J	524	Total	C	N	O	S	0	0	0
			3849	2395	665	769	20			
1	K	524	Total	C	N	O	S	0	0	0
			3849	2395	665	769	20			
1	L	524	Total	C	N	O	S	0	0	0
			3849	2395	665	769	20			
1	M	524	Total	C	N	O	S	0	0	0
			3849	2395	665	769	20			
1	N	524	Total	C	N	O	S	0	0	0
			3849	2395	665	769	20			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	52	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	398	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
B	52	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
B	398	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
C	52	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
C	398	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
D	52	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
D	398	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
E	52	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
E	398	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
F	52	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
F	398	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
G	52	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
G	398	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
H	52	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
H	398	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
I	52	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
I	398	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
J	52	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
J	398	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
K	52	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
K	398	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
L	52	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
L	398	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
M	52	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
M	398	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
N	52	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5
N	398	ALA	ASP	ENGINEERED MUTATION	UNP P0A6F5

- Molecule 2 is a protein called 10 kDa chaperonin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	O	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	P	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	Q	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	R	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	S	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	T	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	U	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	V	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	W	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	X	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	Y	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	Z	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	a	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			
2	b	97	Total	C	N	O	S	0	0	0
			728	454	127	145	2			

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	D	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	E	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	F	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	G	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	H	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	I	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	J	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	K	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	L	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	M	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	N	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	G	1	Total	Mg	0	0
			1	1		
4	J	1	Total	Mg	0	0
			1	1		
4	D	1	Total	Mg	0	0
			1	1		
4	K	1	Total	Mg	0	0
			1	1		
4	E	1	Total	Mg	0	0
			1	1		
4	H	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		
4	I	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	1	Total 1	Mg 1	0	0
4	A	1	Total 1	Mg 1	0	0
4	N	1	Total 1	Mg 1	0	0
4	L	1	Total 1	Mg 1	0	0
4	F	1	Total 1	Mg 1	0	0
4	M	1	Total 1	Mg 1	0	0

- Molecule 5 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	G	1	Total 1	K 1	0	0
5	J	1	Total 1	K 1	0	0
5	D	1	Total 1	K 1	0	0
5	K	1	Total 1	K 1	0	0
5	E	1	Total 1	K 1	0	0
5	H	1	Total 1	K 1	0	0
5	B	1	Total 1	K 1	0	0
5	I	1	Total 1	K 1	0	0
5	C	1	Total 1	K 1	0	0
5	A	1	Total 1	K 1	0	0
5	N	1	Total 1	K 1	0	0
5	L	1	Total 1	K 1	0	0
5	F	1	Total 1	K 1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	M	1	Total	K	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 60 kDa chaperonin




- Molecule 1: 60 kDa chaperonin




- Molecule 1: 60 kDa chaperonin




- Molecule 1: 60 kDa chaperonin




- Molecule 1: 60 kDa chaperonin








• Molecule 1: 60 kDa chaperonin




• Molecule 1: 60 kDa chaperonin







- Molecule 2: 10 kDa chaperonin




- Molecule 2: 10 kDa chaperonin




- Molecule 2: 10 kDa chaperonin




- Molecule 2: 10 kDa chaperonin





- Molecule 2: 10 kDa chaperonin





- Molecule 2: 10 kDa chaperonin



- Molecule 2: 10 kDa chaperonin




- Molecule 2: 10 kDa chaperonin




- Molecule 2: 10 kDa chaperonin



- Molecule 2: 10 kDa chaperonin









- Molecule 2: 10 kDa chaperonin



- Molecule 2: 10 kDa chaperonin



- Molecule 2: 10 kDa chaperonin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	160.28Å 217.12Å 198.48Å 90.00° 98.02° 90.00°	Depositor
Resolution (Å)	49.56 – 3.79 49.56 – 3.79	Depositor EDS
% Data completeness (in resolution range)	91.0 (49.56-3.79) 83.4 (49.56-3.79)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.45 (at 3.77Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1589)	Depositor
R, R_{free}	0.211 , 0.250 0.213 , 0.223	Depositor DCC
R_{free} test set	5551 reflections (5.02%)	DCC
Wilson B-factor (Å ²)	112.4	Xtriage
Anisotropy	0.294	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 148.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	4 of 122258 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	64540	wwPDB-VP
Average B, all atoms (Å ²)	175.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 64.02 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 8.8390e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K, MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/3877	0.43	0/5235
1	B	0.25	0/3877	0.45	0/5235
1	C	0.25	0/3877	0.44	0/5235
1	D	0.25	0/3877	0.46	1/5235 (0.0%)
1	E	0.25	0/3877	0.44	0/5235
1	F	0.25	0/3877	0.44	0/5235
1	G	0.25	0/3877	0.45	0/5235
1	H	0.25	0/3877	0.44	0/5235
1	I	0.25	0/3877	0.45	0/5235
1	J	0.25	0/3877	0.44	0/5235
1	K	0.25	0/3877	0.43	0/5235
1	L	0.25	0/3877	0.44	0/5235
1	M	0.25	0/3877	0.45	0/5235
1	N	0.25	0/3877	0.44	0/5235
2	O	0.19	0/732	0.38	0/983
2	P	0.19	0/732	0.38	0/983
2	Q	0.20	0/732	0.39	0/983
2	R	0.19	0/732	0.39	0/983
2	S	0.19	0/732	0.40	0/983
2	T	0.19	0/732	0.37	0/983
2	U	0.20	0/732	0.39	0/983
2	V	0.19	0/732	0.40	0/983
2	W	0.19	0/732	0.41	0/983
2	X	0.19	0/732	0.39	0/983
2	Y	0.20	0/732	0.38	0/983
2	Z	0.19	0/732	0.38	0/983
2	a	0.19	0/732	0.37	0/983
2	b	0.19	0/732	0.37	0/983
All	All	0.24	0/64526	0.44	1/87052 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	17	LEU	CA-CB-CG	5.21	127.28	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3849	0	3978	108	0
1	B	3849	0	3978	107	0
1	C	3849	0	3978	109	0
1	D	3849	0	3977	133	0
1	E	3849	0	3978	111	0
1	F	3849	0	3978	128	0
1	G	3849	0	3978	120	0
1	H	3849	0	3978	109	0
1	I	3849	0	3978	143	0
1	J	3849	0	3978	106	0
1	K	3849	0	3978	126	0
1	L	3849	0	3978	113	0
1	M	3849	0	3978	123	0
1	N	3849	0	3978	126	0
2	O	728	0	762	27	0
2	P	728	0	762	35	0
2	Q	728	0	762	33	0
2	R	728	0	762	23	0
2	S	728	0	762	37	0
2	T	728	0	762	22	0
2	U	728	0	762	44	0
2	V	728	0	762	35	0
2	W	728	0	762	40	0
2	X	728	0	762	24	0
2	Y	728	0	762	26	0
2	Z	728	0	762	22	0
2	a	728	0	762	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	b	728	0	762	0	0
3	A	31	0	12	1	0
3	B	31	0	12	3	0
3	C	31	0	12	0	0
3	D	31	0	12	1	0
3	E	31	0	12	0	0
3	F	31	0	12	1	0
3	G	31	0	12	1	0
3	H	31	0	12	1	0
3	I	31	0	12	4	0
3	J	31	0	12	1	0
3	K	31	0	12	1	0
3	L	31	0	12	1	0
3	M	31	0	12	0	0
3	N	31	0	12	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
4	G	1	0	0	0	0
4	H	1	0	0	0	0
4	I	1	0	0	0	0
4	J	1	0	0	0	0
4	K	1	0	0	0	0
4	L	1	0	0	0	0
4	M	1	0	0	0	0
4	N	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
5	G	1	0	0	0	0
5	H	1	0	0	0	0
5	I	1	0	0	0	0
5	J	1	0	0	0	0
5	K	1	0	0	0	0
5	L	1	0	0	0	0
5	M	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	N	1	0	0	0	0
All	All	64540	0	66527	1902	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 1902 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:10:ASN:HD21	1:H:10:ASN:HD21	1.11	0.98
1:B:10:ASN:HD21	1:I:10:ASN:HD21	1.11	0.97
1:D:226:LYS:HZ3	1:D:252:GLU:H	1.12	0.92
1:G:10:ASN:HD21	1:K:10:ASN:HD21	1.19	0.90
1:G:77:VAL:HG22	1:G:510:VAL:HG21	1.54	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	522/548 (95%)	503 (96%)	17 (3%)	2 (0%)	39	80
1	B	522/548 (95%)	504 (97%)	16 (3%)	2 (0%)	39	80
1	C	522/548 (95%)	503 (96%)	15 (3%)	4 (1%)	24	70
1	D	522/548 (95%)	500 (96%)	18 (3%)	4 (1%)	24	70
1	E	522/548 (95%)	500 (96%)	19 (4%)	3 (1%)	30	74
1	F	522/548 (95%)	498 (95%)	19 (4%)	5 (1%)	19	66
1	G	522/548 (95%)	500 (96%)	18 (3%)	4 (1%)	24	70
1	H	522/548 (95%)	502 (96%)	17 (3%)	3 (1%)	30	74

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	I	522/548 (95%)	502 (96%)	17 (3%)	3 (1%)	30	74
1	J	522/548 (95%)	497 (95%)	20 (4%)	5 (1%)	19	66
1	K	522/548 (95%)	503 (96%)	16 (3%)	3 (1%)	30	74
1	L	522/548 (95%)	501 (96%)	18 (3%)	3 (1%)	30	74
1	M	522/548 (95%)	501 (96%)	19 (4%)	2 (0%)	39	80
1	N	522/548 (95%)	500 (96%)	18 (3%)	4 (1%)	24	70
2	O	95/97 (98%)	83 (87%)	10 (10%)	2 (2%)	9	53
2	P	95/97 (98%)	84 (88%)	9 (10%)	2 (2%)	9	53
2	Q	95/97 (98%)	83 (87%)	10 (10%)	2 (2%)	9	53
2	R	95/97 (98%)	82 (86%)	10 (10%)	3 (3%)	5	44
2	S	95/97 (98%)	83 (87%)	10 (10%)	2 (2%)	9	53
2	T	95/97 (98%)	81 (85%)	12 (13%)	2 (2%)	9	53
2	U	95/97 (98%)	83 (87%)	9 (10%)	3 (3%)	5	44
2	V	95/97 (98%)	82 (86%)	11 (12%)	2 (2%)	9	53
2	W	95/97 (98%)	83 (87%)	10 (10%)	2 (2%)	9	53
2	X	95/97 (98%)	83 (87%)	10 (10%)	2 (2%)	9	53
2	Y	95/97 (98%)	83 (87%)	10 (10%)	2 (2%)	9	53
2	Z	95/97 (98%)	82 (86%)	10 (10%)	3 (3%)	5	44
2	a	95/97 (98%)	83 (87%)	9 (10%)	3 (3%)	5	44
2	b	95/97 (98%)	83 (87%)	10 (10%)	2 (2%)	9	53
All	All	8638/9030 (96%)	8172 (95%)	387 (4%)	79 (1%)	21	68

5 of 79 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	358	SER
1	C	193	MET
1	C	356	ALA
1	D	360	TYR
1	G	193	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	402/413 (97%)	393 (98%)	9 (2%)	60	85
1	B	402/413 (97%)	381 (95%)	21 (5%)	29	69
1	C	402/413 (97%)	381 (95%)	21 (5%)	29	69
1	D	402/413 (97%)	372 (92%)	30 (8%)	17	57
1	E	402/413 (97%)	375 (93%)	27 (7%)	20	62
1	F	402/413 (97%)	382 (95%)	20 (5%)	30	70
1	G	402/413 (97%)	388 (96%)	14 (4%)	43	78
1	H	402/413 (97%)	386 (96%)	16 (4%)	38	75
1	I	402/413 (97%)	378 (94%)	24 (6%)	24	65
1	J	402/413 (97%)	379 (94%)	23 (6%)	25	67
1	K	402/413 (97%)	378 (94%)	24 (6%)	24	65
1	L	402/413 (97%)	377 (94%)	25 (6%)	23	64
1	M	402/413 (97%)	384 (96%)	18 (4%)	34	73
1	N	402/413 (97%)	377 (94%)	25 (6%)	23	64
2	O	80/80 (100%)	77 (96%)	3 (4%)	40	76
2	P	80/80 (100%)	75 (94%)	5 (6%)	22	64
2	Q	80/80 (100%)	68 (85%)	12 (15%)	3	26
2	R	80/80 (100%)	73 (91%)	7 (9%)	12	50
2	S	80/80 (100%)	71 (89%)	9 (11%)	7	38
2	T	80/80 (100%)	70 (88%)	10 (12%)	6	33
2	U	80/80 (100%)	68 (85%)	12 (15%)	3	26
2	V	80/80 (100%)	67 (84%)	13 (16%)	3	21
2	W	80/80 (100%)	71 (89%)	9 (11%)	7	38
2	X	80/80 (100%)	71 (89%)	9 (11%)	7	38
2	Y	80/80 (100%)	67 (84%)	13 (16%)	3	21
2	Z	80/80 (100%)	75 (94%)	5 (6%)	22	64
2	a	80/80 (100%)	76 (95%)	4 (5%)	30	70
2	b	80/80 (100%)	72 (90%)	8 (10%)	9	43
All	All	6748/6902 (98%)	6332 (94%)	416 (6%)	23	64

5 of 416 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	204	PHE
1	L	204	PHE
2	X	85	ILE
1	J	257	GLU
1	K	236	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	10	ASN
1	H	10	ASN
1	J	352	GLN
1	F	479	ASN
1	I	432	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 42 ligands modelled in this entry, 28 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ATP	A	601	5,4	24,33,33	0.96	1 (4%)	31,52,52	1.85	5 (16%)
3	ATP	B	601	5,4	24,33,33	0.97	1 (4%)	31,52,52	1.81	5 (16%)
3	ATP	C	601	5,4	24,33,33	0.96	1 (4%)	31,52,52	1.89	5 (16%)
3	ATP	D	601	5,4	24,33,33	0.93	1 (4%)	31,52,52	1.91	5 (16%)
3	ATP	E	601	5,4	24,33,33	0.96	1 (4%)	31,52,52	1.99	5 (16%)
3	ATP	F	601	5,4	24,33,33	0.94	1 (4%)	31,52,52	1.85	5 (16%)
3	ATP	G	601	5,4	24,33,33	0.93	1 (4%)	31,52,52	1.96	6 (19%)
3	ATP	H	601	5,4	24,33,33	0.97	1 (4%)	31,52,52	1.81	5 (16%)
3	ATP	I	601	5,4	24,33,33	0.96	1 (4%)	31,52,52	1.89	5 (16%)
3	ATP	J	601	5,4	24,33,33	0.98	1 (4%)	31,52,52	2.00	5 (16%)
3	ATP	K	601	5,4	24,33,33	0.94	1 (4%)	31,52,52	1.89	5 (16%)
3	ATP	L	601	5,4	24,33,33	0.94	1 (4%)	31,52,52	1.98	6 (19%)
3	ATP	M	601	5,4	24,33,33	0.94	1 (4%)	31,52,52	1.90	6 (19%)
3	ATP	N	601	5,4	24,33,33	0.95	1 (4%)	31,52,52	1.84	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATP	A	601	5,4	-	0/18/38/38	0/3/3/3
3	ATP	B	601	5,4	-	0/18/38/38	0/3/3/3
3	ATP	C	601	5,4	-	0/18/38/38	0/3/3/3
3	ATP	D	601	5,4	-	0/18/38/38	0/3/3/3
3	ATP	E	601	5,4	-	0/18/38/38	0/3/3/3
3	ATP	F	601	5,4	-	0/18/38/38	0/3/3/3
3	ATP	G	601	5,4	-	0/18/38/38	0/3/3/3
3	ATP	H	601	5,4	-	0/18/38/38	0/3/3/3
3	ATP	I	601	5,4	-	0/18/38/38	0/3/3/3
3	ATP	J	601	5,4	-	0/18/38/38	0/3/3/3
3	ATP	K	601	5,4	-	0/18/38/38	0/3/3/3
3	ATP	L	601	5,4	-	0/18/38/38	0/3/3/3
3	ATP	M	601	5,4	-	0/18/38/38	0/3/3/3
3	ATP	N	601	5,4	-	0/18/38/38	0/3/3/3

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	601	ATP	C5-C4	2.97	1.47	1.40
3	M	601	ATP	C5-C4	2.99	1.47	1.40
3	F	601	ATP	C5-C4	3.02	1.47	1.40
3	E	601	ATP	C5-C4	3.02	1.47	1.40
3	G	601	ATP	C5-C4	3.04	1.47	1.40

The worst 5 of 73 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	601	ATP	N3-C2-N1	-7.02	123.52	128.89
3	L	601	ATP	N3-C2-N1	-6.96	123.57	128.89
3	M	601	ATP	N3-C2-N1	-6.88	123.62	128.89
3	E	601	ATP	N3-C2-N1	-6.85	123.65	128.89
3	K	601	ATP	N3-C2-N1	-6.68	123.78	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

11 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	601	ATP	1	0
3	B	601	ATP	3	0
3	D	601	ATP	1	0
3	F	601	ATP	1	0
3	G	601	ATP	1	0
3	H	601	ATP	1	0
3	I	601	ATP	4	0
3	J	601	ATP	1	0
3	K	601	ATP	1	0
3	L	601	ATP	1	0
3	N	601	ATP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	524/548 (95%)	-0.22	9 (1%) 73 58	68, 134, 281, 335	0
1	B	524/548 (95%)	-0.23	6 (1%) 82 69	71, 134, 286, 336	0
1	C	524/548 (95%)	-0.29	7 (1%) 79 65	65, 132, 275, 331	0
1	D	524/548 (95%)	-0.30	8 (1%) 76 62	68, 130, 277, 327	0
1	E	524/548 (95%)	-0.29	4 (0%) 87 77	65, 133, 275, 330	0
1	F	524/548 (95%)	-0.35	6 (1%) 82 69	67, 133, 274, 324	0
1	G	524/548 (95%)	-0.25	8 (1%) 76 62	68, 135, 282, 344	0
1	H	524/548 (95%)	-0.23	7 (1%) 79 65	64, 134, 288, 342	0
1	I	524/548 (95%)	-0.20	8 (1%) 76 62	66, 135, 292, 349	0
1	J	524/548 (95%)	-0.31	6 (1%) 82 69	66, 132, 274, 332	0
1	K	524/548 (95%)	-0.30	6 (1%) 82 69	64, 130, 275, 326	0
1	L	524/548 (95%)	-0.29	6 (1%) 82 69	69, 132, 276, 328	0
1	M	524/548 (95%)	-0.33	4 (0%) 87 77	63, 134, 276, 334	0
1	N	524/548 (95%)	-0.25	8 (1%) 76 62	73, 132, 297, 355	0
2	O	97/97 (100%)	0.31	9 (9%) 11 7	183, 233, 312, 338	0
2	P	97/97 (100%)	0.17	1 (1%) 84 72	189, 233, 313, 346	0
2	Q	97/97 (100%)	0.94	14 (14%) 3 3	194, 240, 324, 352	0
2	R	97/97 (100%)	0.76	16 (16%) 2 2	205, 243, 313, 341	0
2	S	97/97 (100%)	0.71	14 (14%) 3 3	207, 246, 314, 354	0
2	T	97/97 (100%)	1.15	22 (22%) 1 1	205, 244, 315, 346	0
2	U	97/97 (100%)	0.77	10 (10%) 9 6	188, 234, 314, 348	0
2	V	97/97 (100%)	0.33	7 (7%) 18 11	190, 236, 312, 348	0
2	W	97/97 (100%)	0.25	1 (1%) 84 72	195, 237, 315, 347	0
2	X	97/97 (100%)	0.79	13 (13%) 4 4	186, 232, 316, 346	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å²)	Q<0.9
2	Y	97/97 (100%)	0.98	19 (19%)	1 1	191, 232, 313, 344	0
2	Z	97/97 (100%)	0.61	10 (10%)	9 6	197, 236, 316, 355	0
2	a	97/97 (100%)	0.80	13 (13%)	4 4	189, 233, 316, 342	0
2	b	97/97 (100%)	0.80	13 (13%)	4 4	193, 237, 315, 344	0
All	All	8694/9030 (96%)	-0.13	255 (2%)	55 38	63, 157, 291, 355	0

The worst 5 of 255 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	Z	32	ALA	9.3
2	Z	31	ALA	8.4
2	X	32	ALA	7.9
2	Y	32	ALA	7.5
2	R	32	ALA	7.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	K	M	603	1/1	0.92	0.20	1.00	192,192,192,192	0
3	ATP	I	601	31/31	0.96	0.20	0.51	58,120,159,197	0
3	ATP	A	601	31/31	0.95	0.23	0.42	74,112,169,195	0
3	ATP	C	601	31/31	0.94	0.22	0.39	65,116,176,192	0
5	K	I	603	1/1	0.89	0.20	0.39	138,138,138,138	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	ATP	N	601	31/31	0.95	0.23	0.37	75,110,162,207	0
3	ATP	J	601	31/31	0.95	0.20	0.34	76,112,176,190	0
3	ATP	E	601	31/31	0.94	0.23	0.33	61,95,148,174	0
3	ATP	H	601	31/31	0.94	0.23	0.22	67,95,170,203	0
5	K	F	603	1/1	0.94	0.19	0.13	132,132,132,132	0
3	ATP	K	601	31/31	0.96	0.20	0.12	55,109,148,207	0
3	ATP	B	601	31/31	0.94	0.20	0.02	70,113,189,206	0
3	ATP	D	601	31/31	0.97	0.20	0.02	67,95,205,263	0
3	ATP	F	601	31/31	0.97	0.19	-0.01	76,101,160,203	0
3	ATP	M	601	31/31	0.97	0.17	-0.07	71,98,158,206	0
3	ATP	G	601	31/31	0.96	0.20	-0.11	68,99,162,188	0
3	ATP	L	601	31/31	0.96	0.20	-0.13	66,97,157,198	0
5	K	A	603	1/1	0.86	0.20	-0.35	200,200,200,200	0
5	K	D	603	1/1	0.88	0.14	-1.59	139,139,139,139	0
5	K	H	603	1/1	0.99	0.14	-3.04	168,168,168,168	0
4	MG	A	602	1/1	0.96	0.24	-	76,76,76,76	0
4	MG	M	602	1/1	1.00	0.19	-	81,81,81,81	0
5	K	G	603	1/1	0.95	0.25	-	197,197,197,197	0
4	MG	D	602	1/1	0.99	0.17	-	87,87,87,87	0
4	MG	C	602	1/1	0.97	0.25	-	71,71,71,71	0
4	MG	F	602	1/1	1.00	0.20	-	77,77,77,77	0
4	MG	E	602	1/1	0.99	0.26	-	94,94,94,94	0
4	MG	B	602	1/1	0.98	0.27	-	68,68,68,68	0
5	K	E	603	1/1	0.72	0.21	-	140,140,140,140	0
4	MG	L	602	1/1	0.98	0.20	-	87,87,87,87	0
4	MG	H	602	1/1	0.97	0.32	-	74,74,74,74	0
4	MG	J	602	1/1	0.99	0.24	-	112,112,112,112	0
5	K	K	603	1/1	0.93	0.17	-	149,149,149,149	0
5	K	J	603	1/1	0.98	0.23	-	251,251,251,251	0
4	MG	N	602	1/1	0.98	0.21	-	103,103,103,103	0
5	K	L	603	1/1	0.93	0.25	-	185,185,185,185	0
5	K	N	603	1/1	0.96	0.24	-	165,165,165,165	0
5	K	B	603	1/1	0.97	0.19	-	200,200,200,200	0
4	MG	K	602	1/1	1.00	0.21	-	71,71,71,71	0
4	MG	I	602	1/1	0.98	0.26	-	106,106,106,106	0
5	K	C	603	1/1	0.92	0.34	-	291,291,291,291	0
4	MG	G	602	1/1	0.99	0.24	-	88,88,88,88	0

6.5 Other polymers ⓘ

There are no such residues in this entry.