



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:11 PM GMT

PDB ID : 4WYV
Title : Crystal Structure of Human Translin in Open Conformation
Authors : Dvir, H.; Eliahoo, E.; Alian, A.
Deposited on : 2014-11-18
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	240	 71% 16% • 11%
1	B	240	 73% 14% • 12%
1	C	240	 68% 17% • 13%
1	D	240	 71% 15% • 11%
1	E	240	 68% 19% • 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	240	 64% 23% • 12%
1	G	240	 68% 19% • 10%
1	H	240	 73% 13% • 12%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 13935 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Translin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	213	Total	C	N	O	S	0	0	0
			1741	1116	300	323	2			
1	B	212	Total	C	N	O	S	0	0	0
			1737	1114	299	322	2			
1	C	210	Total	C	N	O	S	0	0	0
			1717	1100	296	319	2			
1	D	214	Total	C	N	O	S	0	0	0
			1741	1113	301	325	2			
1	E	213	Total	C	N	O	S	0	0	0
			1743	1117	300	324	2			
1	F	212	Total	C	N	O	S	0	0	0
			1737	1114	299	322	2			
1	G	216	Total	C	N	O	S	0	0	0
			1767	1133	304	328	2			
1	H	212	Total	C	N	O	S	0	0	0
			1734	1112	298	322	2			

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	initiating methionine	UNP Q15631
A	-10	ARG	-	expression tag	UNP Q15631
A	-9	GLY	-	expression tag	UNP Q15631
A	-8	SER	-	expression tag	UNP Q15631
A	-7	HIS	-	expression tag	UNP Q15631
A	-6	HIS	-	expression tag	UNP Q15631
A	-5	HIS	-	expression tag	UNP Q15631
A	-4	HIS	-	expression tag	UNP Q15631
A	-3	HIS	-	expression tag	UNP Q15631
A	-2	HIS	-	expression tag	UNP Q15631
A	-1	GLY	-	expression tag	UNP Q15631
A	0	SER	-	expression tag	UNP Q15631
B	-11	MET	-	initiating methionine	UNP Q15631

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-10	ARG	-	expression tag	UNP Q15631
B	-9	GLY	-	expression tag	UNP Q15631
B	-8	SER	-	expression tag	UNP Q15631
B	-7	HIS	-	expression tag	UNP Q15631
B	-6	HIS	-	expression tag	UNP Q15631
B	-5	HIS	-	expression tag	UNP Q15631
B	-4	HIS	-	expression tag	UNP Q15631
B	-3	HIS	-	expression tag	UNP Q15631
B	-2	HIS	-	expression tag	UNP Q15631
B	-1	GLY	-	expression tag	UNP Q15631
B	0	SER	-	expression tag	UNP Q15631
C	-11	MET	-	initiating methionine	UNP Q15631
C	-10	ARG	-	expression tag	UNP Q15631
C	-9	GLY	-	expression tag	UNP Q15631
C	-8	SER	-	expression tag	UNP Q15631
C	-7	HIS	-	expression tag	UNP Q15631
C	-6	HIS	-	expression tag	UNP Q15631
C	-5	HIS	-	expression tag	UNP Q15631
C	-4	HIS	-	expression tag	UNP Q15631
C	-3	HIS	-	expression tag	UNP Q15631
C	-2	HIS	-	expression tag	UNP Q15631
C	-1	GLY	-	expression tag	UNP Q15631
C	0	SER	-	expression tag	UNP Q15631
D	-11	MET	-	initiating methionine	UNP Q15631
D	-10	ARG	-	expression tag	UNP Q15631
D	-9	GLY	-	expression tag	UNP Q15631
D	-8	SER	-	expression tag	UNP Q15631
D	-7	HIS	-	expression tag	UNP Q15631
D	-6	HIS	-	expression tag	UNP Q15631
D	-5	HIS	-	expression tag	UNP Q15631
D	-4	HIS	-	expression tag	UNP Q15631
D	-3	HIS	-	expression tag	UNP Q15631
D	-2	HIS	-	expression tag	UNP Q15631
D	-1	GLY	-	expression tag	UNP Q15631
D	0	SER	-	expression tag	UNP Q15631
E	-11	MET	-	initiating methionine	UNP Q15631
E	-10	ARG	-	expression tag	UNP Q15631
E	-9	GLY	-	expression tag	UNP Q15631
E	-8	SER	-	expression tag	UNP Q15631
E	-7	HIS	-	expression tag	UNP Q15631
E	-6	HIS	-	expression tag	UNP Q15631
E	-5	HIS	-	expression tag	UNP Q15631

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-4	HIS	-	expression tag	UNP Q15631
E	-3	HIS	-	expression tag	UNP Q15631
E	-2	HIS	-	expression tag	UNP Q15631
E	-1	GLY	-	expression tag	UNP Q15631
E	0	SER	-	expression tag	UNP Q15631
F	-11	MET	-	initiating methionine	UNP Q15631
F	-10	ARG	-	expression tag	UNP Q15631
F	-9	GLY	-	expression tag	UNP Q15631
F	-8	SER	-	expression tag	UNP Q15631
F	-7	HIS	-	expression tag	UNP Q15631
F	-6	HIS	-	expression tag	UNP Q15631
F	-5	HIS	-	expression tag	UNP Q15631
F	-4	HIS	-	expression tag	UNP Q15631
F	-3	HIS	-	expression tag	UNP Q15631
F	-2	HIS	-	expression tag	UNP Q15631
F	-1	GLY	-	expression tag	UNP Q15631
F	0	SER	-	expression tag	UNP Q15631
G	-11	MET	-	initiating methionine	UNP Q15631
G	-10	ARG	-	expression tag	UNP Q15631
G	-9	GLY	-	expression tag	UNP Q15631
G	-8	SER	-	expression tag	UNP Q15631
G	-7	HIS	-	expression tag	UNP Q15631
G	-6	HIS	-	expression tag	UNP Q15631
G	-5	HIS	-	expression tag	UNP Q15631
G	-4	HIS	-	expression tag	UNP Q15631
G	-3	HIS	-	expression tag	UNP Q15631
G	-2	HIS	-	expression tag	UNP Q15631
G	-1	GLY	-	expression tag	UNP Q15631
G	0	SER	-	expression tag	UNP Q15631
H	-11	MET	-	initiating methionine	UNP Q15631
H	-10	ARG	-	expression tag	UNP Q15631
H	-9	GLY	-	expression tag	UNP Q15631
H	-8	SER	-	expression tag	UNP Q15631
H	-7	HIS	-	expression tag	UNP Q15631
H	-6	HIS	-	expression tag	UNP Q15631
H	-5	HIS	-	expression tag	UNP Q15631
H	-4	HIS	-	expression tag	UNP Q15631
H	-3	HIS	-	expression tag	UNP Q15631
H	-2	HIS	-	expression tag	UNP Q15631
H	-1	GLY	-	expression tag	UNP Q15631
H	0	SER	-	expression tag	UNP Q15631


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	7	Total 7	O 7	0	0
2	B	2	Total 2	O 2	0	0
2	C	2	Total 2	O 2	0	0
2	D	2	Total 2	O 2	0	0
2	E	2	Total 2	O 2	0	0
2	F	2	Total 2	O 2	0	0
2	G	1	Total 1	O 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Translin




• Molecule 1: Translin



• Molecule 1: Translin




• Molecule 1: Translin




- Molecule 1: Translin

Chain E: 




- Molecule 1: Translin

Chain F: 




- Molecule 1: Translin

Chain G: 



- Molecule 1: Translin

Chain H: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	82.31Å 82.31Å 635.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.00 – 3.00 57.96 – 3.00	Depositor EDS
% Data completeness (in resolution range)	98.5 (58.00-3.00) 98.6 (57.96-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.93 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.226 , 0.272 0.215 , 0.259	Depositor DCC
R_{free} test set	2272 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	76.8	Xtriage
Anisotropy	0.031	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 47.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtriage
Outliers	1 of 45031 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13935	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.91	3/1772 (0.2%)	0.76	1/2390 (0.0%)
1	B	0.88	2/1768 (0.1%)	0.79	2/2385 (0.1%)
1	C	0.94	5/1747 (0.3%)	0.75	2/2357 (0.1%)
1	D	0.83	0/1771	0.73	0/2389
1	E	0.76	0/1774	0.68	0/2393
1	F	0.84	3/1768 (0.2%)	0.72	1/2385 (0.0%)
1	G	0.93	4/1799 (0.2%)	0.77	3/2426 (0.1%)
1	H	0.83	2/1765 (0.1%)	0.71	0/2381
All	All	0.87	19/14164 (0.1%)	0.74	9/19106 (0.0%)

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	46	HIS	C-O	12.28	1.46	1.23
1	A	9	GLU	CD-OE1	12.14	1.39	1.25
1	G	69	LYS	CG-CD	10.11	1.86	1.52
1	H	4	SER	CB-OG	9.42	1.54	1.42
1	B	194	ARG	CZ-NH1	7.98	1.43	1.33

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	194	ARG	NE-CZ-NH2	-9.94	115.33	120.30
1	G	129	ARG	NE-CZ-NH1	9.67	125.13	120.30
1	G	129	ARG	NE-CZ-NH2	-6.67	116.97	120.30
1	B	183	LEU	CA-CB-CG	6.52	130.29	115.30
1	G	21	ARG	NE-CZ-NH1	5.42	123.01	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1741	0	1761	25	0
1	B	1737	0	1758	26	0
1	C	1717	0	1741	37	0
1	D	1741	0	1762	24	0
1	E	1743	0	1763	30	0
1	F	1737	0	1758	40	0
1	G	1767	0	1783	38	0
1	H	1734	0	1755	28	0
2	A	7	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
2	G	1	0	0	0	0
All	All	13935	0	14081	218	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 218 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:69:LYS:CD	1:G:69:LYS:CG	1.86	1.50
1:F:110:THR:CG2	1:F:112:THR:HG22	1.81	1.09
1:F:110:THR:HG23	1:F:112:THR:HG22	1.37	1.03
1:E:72:LEU:HD23	1:E:125:ILE:HD11	1.52	0.91
1:C:39:LEU:HD11	1:C:43:GLN:HE21	1.43	0.83

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	209/240 (87%)	201 (96%)	4 (2%)	4 (2%)	10	43
1	B	208/240 (87%)	202 (97%)	4 (2%)	2 (1%)	19	61
1	C	206/240 (86%)	198 (96%)	6 (3%)	2 (1%)	19	61
1	D	210/240 (88%)	202 (96%)	4 (2%)	4 (2%)	10	43
1	E	209/240 (87%)	199 (95%)	5 (2%)	5 (2%)	7	35
1	F	208/240 (87%)	201 (97%)	4 (2%)	3 (1%)	14	51
1	G	212/240 (88%)	204 (96%)	5 (2%)	3 (1%)	14	51
1	H	208/240 (87%)	199 (96%)	5 (2%)	4 (2%)	10	43
All	All	1670/1920 (87%)	1606 (96%)	37 (2%)	27 (2%)	12	48

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	131	LYS
1	A	133	PHE
1	B	133	PHE
1	C	133	PHE
1	D	133	PHE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/212 (91%)	187 (97%)	5 (3%)	54	85

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	192/212 (91%)	183 (95%)	9 (5%)	32	72
1	C	190/212 (90%)	180 (95%)	10 (5%)	28	67
1	D	192/212 (91%)	185 (96%)	7 (4%)	42	79
1	E	193/212 (91%)	184 (95%)	9 (5%)	32	72
1	F	192/212 (91%)	182 (95%)	10 (5%)	29	68
1	G	195/212 (92%)	189 (97%)	6 (3%)	47	83
1	H	192/212 (91%)	184 (96%)	8 (4%)	36	76
All	All	1538/1696 (91%)	1474 (96%)	64 (4%)	36	76

5 of 64 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	112	THR
1	E	112	THR
1	H	53	ASP
1	D	190	SER
1	E	31	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	47	GLN
1	E	43	GLN
1	F	134	HIS
1	D	11	GLN
1	F	90	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	213/240 (88%)	-0.34	0 100 100	40, 72, 106, 127	0
1	B	212/240 (88%)	-0.38	0 100 100	44, 74, 104, 128	0
1	C	210/240 (87%)	-0.20	1 (0%) 91 76	46, 80, 112, 125	0
1	D	214/240 (89%)	-0.31	0 100 100	47, 73, 109, 131	0
1	E	213/240 (88%)	-0.32	1 (0%) 91 76	59, 85, 114, 134	0
1	F	212/240 (88%)	-0.19	0 100 100	60, 91, 122, 136	0
1	G	216/240 (90%)	-0.33	0 100 100	50, 79, 110, 132	0
1	H	212/240 (88%)	-0.16	1 (0%) 91 76	59, 85, 105, 121	0
All	All	1702/1920 (88%)	-0.28	3 (0%) 95 87	40, 81, 112, 136	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	129	ARG	3.3
1	C	129	ARG	2.7
1	E	47	GLN	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.