



Full wwPDB X-ray Structure Validation Report i

Feb 27, 2016 – 11:06 AM GMT

PDB ID : 3X3E
Title : Crystal structure of LysK from Thermus thermophilus complex with Lysine
Authors : Fujita, S.; Hasebe, F.; Cho, S.H.; Tomita, T.; Kuzuyama, T.; Nishiyama, M.
Deposited on : 2015-01-19
Resolution : 2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026982
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
			8%	73%	17%	•	7%
1	A	372					
1	B	372	5%	76%	16%	•	6%

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 5615 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called N-acetyl-lysine deacetylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	346	Total	C 2682	N 1717	O 476	S 483	6	0	0
1	B	350	Total	C 2742	N 1758	O 489	S 489	6	0	2

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	EXPRESSION TAG	UNP Q8VUS5
A	-9	ALA	-	EXPRESSION TAG	UNP Q8VUS5
A	-8	SER	-	EXPRESSION TAG	UNP Q8VUS5
A	-7	TRP	-	EXPRESSION TAG	UNP Q8VUS5
A	-6	SER	-	EXPRESSION TAG	UNP Q8VUS5
A	-5	HIS	-	EXPRESSION TAG	UNP Q8VUS5
A	-4	PRO	-	EXPRESSION TAG	UNP Q8VUS5
A	-3	GLN	-	EXPRESSION TAG	UNP Q8VUS5
A	-2	PHE	-	EXPRESSION TAG	UNP Q8VUS5
A	-1	GLU	-	EXPRESSION TAG	UNP Q8VUS5
A	0	GLY	-	EXPRESSION TAG	UNP Q8VUS5
A	1	GLY	-	EXPRESSION TAG	UNP Q8VUS5
B	-10	MET	-	EXPRESSION TAG	UNP Q8VUS5
B	-9	ALA	-	EXPRESSION TAG	UNP Q8VUS5
B	-8	SER	-	EXPRESSION TAG	UNP Q8VUS5
B	-7	TRP	-	EXPRESSION TAG	UNP Q8VUS5
B	-6	SER	-	EXPRESSION TAG	UNP Q8VUS5
B	-5	HIS	-	EXPRESSION TAG	UNP Q8VUS5
B	-4	PRO	-	EXPRESSION TAG	UNP Q8VUS5
B	-3	GLN	-	EXPRESSION TAG	UNP Q8VUS5
B	-2	PHE	-	EXPRESSION TAG	UNP Q8VUS5
B	-1	GLU	-	EXPRESSION TAG	UNP Q8VUS5
B	0	GLY	-	EXPRESSION TAG	UNP Q8VUS5
B	1	GLY	-	EXPRESSION TAG	UNP Q8VUS5

- Molecule 2 is LYSINE (three-letter code: LYS) (formula: C₆H₁₅N₂O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	10	6	2	2	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	O	S			
3	A	1	5	4	1		0	0
3	B	1	5	4	1		0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total O S 5 4 1	0	0



- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	91	Total O 91 91	0	0
4	B	75	Total O 75 75	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N-acetyl-lysine deacetylase



- Molecule 1: N-acetyl-lysine deacetylase



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	132.78 Å 69.93 Å 118.09 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.65 – 2.39 35.65 – 2.39	Depositor EDS
% Data completeness (in resolution range)	99.3 (35.65-2.39) 99.3 (35.65-2.39)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	3.32 (at 2.39 Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R , R_{free}	0.197 , 0.239 0.195 , 0.237	Depositor DCC
R_{free} test set	2224 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	48.3	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 35.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Outliers	0 of 44081 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5615	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.61	2/2750 (0.1%)	0.76	2/3730 (0.1%)
1	B	0.61	3/2822 (0.1%)	0.70	1/3829 (0.0%)
All	All	0.61	5/5572 (0.1%)	0.73	3/7559 (0.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	152	TRP	CD2-CE2	5.55	1.48	1.41
1	B	310	TRP	CD2-CE2	5.43	1.47	1.41
1	A	152	TRP	CD2-CE2	5.25	1.47	1.41
1	B	197	TRP	CD2-CE2	5.17	1.47	1.41
1	A	197	TRP	CD2-CE2	5.13	1.47	1.41

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	236	ARG	NE-CZ-NH2	-6.68	116.96	120.30
1	A	236	ARG	NE-CZ-NH1	6.38	123.49	120.30
1	B	236	ARG	NE-CZ-NH2	5.25	122.92	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2682	0	2689	43	0
1	B	2742	0	2744	33	0
2	A	10	0	12	2	0
3	A	5	0	0	1	0
3	B	10	0	0	0	0
4	A	91	0	0	1	0
4	B	75	0	0	2	0
All	All	5615	0	5445	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:TYR:HD2	1:A:303:MET:HE2	1.32	0.94
1:A:273:LYS:O	1:A:278:THR:HG21	1.69	0.93
1:B:175:HIS:HD2	1:B:177:ALA:H	1.16	0.92
1:A:247:ILE:O	1:A:251:THR:HG23	1.74	0.88
1:A:170:GLU:HG3	1:A:228:VAL:HG22	1.58	0.83
1:B:20:SER:HB3	1:B:28:VAL:HG21	1.65	0.77
1:A:112:LYS:HE2	1:A:113:ARG:HE	1.52	0.74
1:B:197:TRP:HE1	1:B:249:HIS:HD2	1.39	0.71
1:A:159:TYR:HD2	1:A:303:MET:CE	2.05	0.70
1:B:20:SER:HB3	1:B:28:VAL:CG2	2.24	0.67
1:A:275:THR:HB	1:A:278:THR:HB	1.77	0.67
1:A:255:PRO:HG2	1:A:257:THR:HG22	1.76	0.66
1:B:275:THR:H	1:B:278:THR:HG22	1.61	0.66
1:A:159:TYR:CD2	1:A:303:MET:HE2	2.23	0.65
1:A:249:HIS:HD2	1:A:253:TYR:OH	1.80	0.64
1:B:273:LYS:O	1:B:278:THR:HG21	1.97	0.63
1:B:73:GLY:O	1:B:87:ARG:HD2	2.01	0.61
1:A:20:SER:HB3	1:A:28:VAL:CG2	2.29	0.61
1:A:182:ASN:HA	1:A:227:GLN:HE22	1.65	0.60
1:B:190:TYR:CZ	1:B:194:ILE:HD11	2.36	0.60
1:B:275:THR:H	1:B:278:THR:CG2	2.15	0.60
1:B:274:ASP:HA	1:B:278:THR:HG21	1.84	0.60
1:A:265:ARG:NH1	3:A:402:SO4:O2	2.30	0.58
1:A:20:SER:HB3	1:A:28:VAL:HG22	1.86	0.58
1:B:21:PRO:HG2	1:B:24:LYS:HD3	1.86	0.58
1:A:236:ARG:HH22	2:A:401:LYS:C	2.09	0.56
1:A:278:THR:O	1:A:282:ARG:HG3	2.05	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:354:THR:HA	1:B:355:HIS:HB3	1.87	0.56
1:B:175:HIS:CD2	1:B:177:ALA:H	2.09	0.55
1:A:267:VAL:HG12	1:A:268:PRO:O	2.09	0.52
1:A:155:ILE:HD11	1:A:315:VAL:HG11	1.92	0.51
1:A:236:ARG:NH2	2:A:401:LYS:O	2.42	0.51
1:A:105:GLY:O	1:A:345:ARG:HD2	2.10	0.51
1:B:282:ARG:NH1	4:B:561:HOH:O	2.44	0.50
1:A:106:LEU:HD21	1:A:348:LEU:HB3	1.94	0.50
1:A:108:GLU:HA	1:A:111:ARG:HD3	1.93	0.50
1:A:158:GLY:HA2	1:A:303:MET:HE3	1.94	0.50
1:B:190:TYR:CE2	1:B:194:ILE:HD11	2.48	0.48
1:A:275:THR:HG22	1:A:278:THR:H	1.79	0.48
1:B:335:ALA:O	1:B:339:LYS:HG2	2.14	0.48
1:B:58:GLY:HA3	1:B:113:ARG:C	2.35	0.47
1:A:170:GLU:HG3	1:A:228:VAL:CG2	2.40	0.47
1:A:90:VAL:HG22	1:A:320:GLY:HA3	1.95	0.47
1:B:237:LEU:HG	1:B:241:LEU:O	2.15	0.47
1:B:122:THR:HG23	1:B:123:GLU:HG3	1.96	0.46
1:B:166:LYS:HG2	4:B:520:HOH:O	2.15	0.46
1:B:43:GLY:HA2	1:B:52:ARG:O	2.15	0.46
1:B:122:THR:CG2	1:B:130:LYS:HE2	2.46	0.46
1:B:87:ARG:NH2	1:B:327:THR:O	2.45	0.46
1:B:20:SER:CB	1:B:28:VAL:HG21	2.43	0.45
1:A:220:VAL:O	1:A:222:PRO:HD3	2.16	0.45
1:A:212:GLN:HE22	1:B:179:HIS:HA	1.81	0.45
1:A:170:GLU:HG2	1:A:226:ARG:HD3	1.98	0.45
1:A:18:ILE:HB	1:A:28:VAL:HG13	1.98	0.45
1:A:106:LEU:O	1:A:111:ARG:HD2	2.16	0.45
1:A:309:HIS:HD2	4:A:539:HOH:O	2.00	0.44
1:A:187:LEU:HA	1:A:187:LEU:HD12	1.84	0.44
1:A:20:SER:HB3	1:A:28:VAL:HG21	1.98	0.44
1:B:212:GLN:O	1:B:235:LEU:HA	2.17	0.44
1:A:225:LEU:H	1:A:225:LEU:HD12	1.82	0.44
1:B:327:THR:HB	1:B:328:PRO:HD2	2.01	0.43
1:A:83:ARG:HG2	1:A:331:HIS:CD2	2.54	0.42
1:A:136:ALA:HB3	1:A:137:PRO:HD3	1.99	0.42
1:A:249:HIS:CD2	1:A:253:TYR:OH	2.66	0.42
1:A:43:GLY:HA2	1:A:52:ARG:O	2.19	0.42
1:B:206:ARG:O	1:B:210:GLN:HG3	2.20	0.41
1:B:197:TRP:HE1	1:B:249:HIS:CD2	2.27	0.41
1:A:90:VAL:HG13	1:A:149:PRO:HG3	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:162:ARG:HD2	1:B:236:ARG:NH1	2.35	0.41
1:A:169:ARG:NH2	1:A:171:LYS:HG3	2.36	0.41
1:A:171:LYS:HE3	1:A:181:PRO:O	2.20	0.41
1:A:159:TYR:CD2	1:A:303:MET:CE	2.94	0.40
1:B:307:ALA:HB3	1:B:308:PRO:HD3	2.03	0.40
1:B:63:VAL:HG23	1:B:141:PRO:HB3	2.02	0.40
1:B:197:TRP:NE1	1:B:249:HIS:HD2	2.12	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	342/372 (92%)	328 (96%)	14 (4%)	0	100 100
1	B	350/372 (94%)	343 (98%)	7 (2%)	0	100 100
All	All	692/744 (93%)	671 (97%)	21 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	273/293 (93%)	247 (90%)	26 (10%)	11 15

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	280/293 (96%)	251 (90%)	29 (10%)	9 12
All	All	553/586 (94%)	498 (90%)	55 (10%)	9 14

All (55) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	VAL
1	A	27	LEU
1	A	28	VAL
1	A	30	GLU
1	A	32	LEU
1	A	60	VAL
1	A	78	ARG
1	A	90	VAL
1	A	122	THR
1	A	129	SER
1	A	155	ILE
1	A	157	LEU
1	A	173	HIS
1	A	187	LEU
1	A	192	VAL
1	A	216	ARG
1	A	236	ARG
1	A	241	LEU
1	A	251	THR
1	A	278	THR
1	A	279	ARG
1	A	298	THR
1	A	306	LEU
1	A	348	LEU
1	A	351	LEU
1	A	353	GLN
1	B	9	VAL
1	B	28	VAL
1	B	32	LEU
1	B	34	GLU
1	B	37	GLN
1	B	57	GLU
1	B	62	VAL
1	B	75	ILE
1	B	78	ARG
1	B	106	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	109	GLU
1	B	112	LYS
1	B	122	THR
1	B	129	SER
1	B	135	VAL
1	B	156	THR
1	B	162	ARG
1	B	163	LEU
1	B	187	LEU
1	B	220	VAL
1	B	225	LEU
1	B	237	LEU
1	B	273	LYS
1	B	278	THR
1	B	286	ARG
1	B	306	LEU
1	B	336	GLU
1	B	344	LEU
1	B	353	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	GLN
1	A	221	HIS
1	A	227	GLN
1	A	249	HIS
1	A	326	HIS
1	A	355	HIS
1	B	61	GLN
1	B	175	HIS
1	B	210	GLN
1	B	249	HIS
1	B	326	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	LYS	A	401	-	6,9,9	0.50	0	4,10,10	0.62	0
3	SO4	A	402	-	4,4,4	0.45	0	6,6,6	0.67	0
3	SO4	B	401	-	4,4,4	0.39	0	6,6,6	0.35	0
3	SO4	B	402	-	4,4,4	0.37	0	6,6,6	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LYS	A	401	-	-	0/5/9/9	0/0/0/0
3	SO4	A	402	-	-	0/0/0/0	0/0/0/0
3	SO4	B	401	-	-	0/0/0/0	0/0/0/0
3	SO4	B	402	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	LYS	2	0
3	A	402	SO4	1	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	346/372 (93%)	0.34	31 (8%) 12 11	30, 44, 81, 137	0
1	B	350/372 (94%)	0.21	20 (5%) 27 27	28, 47, 71, 127	0
All	All	696/744 (93%)	0.28	51 (7%) 18 18	28, 46, 72, 137	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	357	GLY	8.8
1	A	225	LEU	7.3
1	A	356	ALA	6.8
1	A	224	GLU	6.6
1	B	355	HIS	6.0
1	A	226	ARG	4.9
1	A	355	HIS	4.4
1	B	192	VAL	4.3
1	A	192	VAL	4.0
1	B	64	LEU	3.9
1	A	354	THR	3.7
1	A	172	ASP	3.7
1	B	354	THR	3.6
1	B	291	ARG	3.4
1	A	223	ALA	3.3
1	A	173	HIS	3.1
1	B	329[A]	TYR	3.0
1	A	222	PRO	3.0
1	A	206	ARG	3.0
1	A	113	ARG	3.0
1	A	147	GLY	3.0
1	A	6	LEU	2.9
1	B	193	ALA	2.8
1	A	108	GLU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	112	LYS	2.6
1	A	353	GLN	2.5
1	B	81	GLY	2.4
1	B	194	ILE	2.4
1	A	291	ARG	2.4
1	B	98	ALA	2.4
1	B	108	GLU	2.4
1	B	109	GLU	2.3
1	A	204	GLY	2.3
1	B	99	MET	2.3
1	B	145	VAL	2.2
1	A	146	ILE	2.2
1	A	109	GLU	2.2
1	B	196	ALA	2.2
1	B	146	ILE	2.2
1	A	329	TYR	2.1
1	A	228	VAL	2.1
1	B	96	PHE	2.1
1	A	98	ALA	2.1
1	A	188	ILE	2.1
1	B	179	HIS	2.1
1	A	145	VAL	2.1
1	A	59	PRO	2.1
1	B	65	LEU	2.1
1	A	191	PHE	2.0
1	B	6	LEU	2.0
1	A	64	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	LYS	A	401	10/10	0.95	0.15	0.29	33,38,39,40	0
3	SO4	B	401	5/5	0.98	0.13	-0.56	60,61,66,66	0
3	SO4	A	402	5/5	0.93	0.25	-	64,64,73,73	0
3	SO4	B	402	5/5	0.96	0.29	-	60,67,76,79	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.