



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2016 – 10:34 PM EDT

PDB ID : 4X3G
Title : Crystal structure of SIAH1 SINA domain in complex with a USP19 peptide
Authors : Walker, J.R.; Dong, A.; Zhang, Q.; Huang, X.; Li, Y.; Bountra, C.; Edwards, A.M.; Arrowsmith, C.H.; Tong, Y.; Structural Genomics Consortium (SGC)
Deposited on : 2014-11-28
Resolution : 2.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027257
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027257


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1406 (2.36-2.32)
Clashscore	102246	1509 (2.36-2.32)
Ramachandran outliers	100387	1490 (2.36-2.32)
Sidechain outliers	100360	1491 (2.36-2.32)
RSRZ outliers	91569	1412 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	193	<div> <div>3%</div> <div> <div></div> <div>90%</div> <div>7%</div> <div>.</div> </div> </div>
1	B	193	<div> <div>3%</div> <div> <div></div> <div>88%</div> <div>10%</div> <div>.</div> </div> </div>
2	C	14	<div> <div>57%</div> <div>21%</div> <div>21%</div> </div>
2	D	14	<div> <div>64%</div> <div>7%</div> <div>29%</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3173 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase SIAH1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	188	Total	C	N	O	S	0	3	0
			1486	939	253	277	17			
1	B	190	Total	C	N	O	S	0	1	0
			1472	933	250	273	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	90	GLY	-	expression tag	UNP Q8IUQ4
B	90	GLY	-	expression tag	UNP Q8IUQ4

- Molecule 2 is a protein called Ubiquitin carboxyl-terminal hydrolase 19.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	11	Total	C	N	O	S	0	0	0
			80	53	12	12	3			
2	D	10	Total	C	N	O	S	0	0	0
			73	48	11	11	3			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Zn	0	0
			2	2		
3	A	2	Total	Zn	0	0
			2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	25	Total	O	0	1
			26	26		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	26	Total 27	O 27	0	1
4	C	1	Total 1	O 1	0	0
4	D	4	Total 4	O 4	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: E3 ubiquitin-protein ligase SIAH1

Chain A: 



- Molecule 1: E3 ubiquitin-protein ligase SIAH1

Chain B: 



- Molecule 2: Ubiquitin carboxyl-terminal hydrolase 19

Chain C: 



- Molecule 2: Ubiquitin carboxyl-terminal hydrolase 19

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	41.34Å 88.09Å 59.59Å 90.00° 103.24° 90.00°	Depositor
Resolution (Å)	48.45 – 2.34 48.45 – 2.34	Depositor EDS
% Data completeness (in resolution range)	89.7 (48.45-2.34) 89.8 (48.45-2.34)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 2.34Å)	Xtriage
Refinement program	BUSTER-TNT	Depositor
R, R_{free}	0.222 , 0.251 0.234 , 0.274	Depositor DCC
R_{free} test set	780 reflections (5.19%)	DCC
Wilson B-factor (Å ²)	60.6	Xtriage
Anisotropy	0.611	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 59.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3173	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/1519	0.58	0/2057
1	B	0.39	0/1507	0.57	0/2043
2	C	0.41	0/84	0.51	0/115
2	D	0.40	0/76	0.45	0/104
All	All	0.40	0/3186	0.57	0/4319

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1486	0	1406	8	0
1	B	1472	0	1399	9	0
2	C	80	0	87	2	0
2	D	73	0	79	1	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	26	0	0	0	0
4	B	27	0	0	0	0
4	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	4	0	0	0	0
All	All	3173	0	2971	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:226:GLU:HG3	1:A:276:ASN:HB2	1.80	0.64
1:A:163:ILE:HG22	2:C:464:PRO:HG2	1.87	0.57
1:B:166:LEU:HD23	2:D:467:MET:HG2	1.90	0.52
1:B:260:ASP:H	1:B:263:ILE:HD12	1.74	0.52
1:A:208:ILE:HD12	1:A:256:CYS:HB3	1.91	0.52
1:A:231:ARG:HH22	1:B:219:GLU:HG2	1.75	0.51
1:B:208:ILE:HD12	1:B:256:CYS:HB3	1.93	0.51
1:A:241:ARG:HE	1:A:250:ALA:HB2	1.75	0.49
1:B:227:LEU:HD12	1:B:234:LEU:HD23	1.95	0.47
1:B:247:ILE:HD12	1:B:251:ILE:HD11	1.96	0.46
1:A:166:LEU:HD23	2:C:467:MET:HG2	1.97	0.46
1:B:191:LEU:HD13	1:B:277:VAL:HG21	1.99	0.44
1:B:263:ILE:HA	1:B:266:LEU:HD12	1.99	0.43
1:A:178:TRP:HB2	1:A:193:LEU:HB3	2.02	0.42
1:B:155:ILE:HG12	1:B:179:VAL:HG13	2.03	0.40
1:A:207:ALA:O	1:A:256:CYS:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	185/193 (96%)	181 (98%)	4 (2%)	0	100	100
1	B	189/193 (98%)	186 (98%)	3 (2%)	0	100	100
2	C	9/14 (64%)	9 (100%)	0	0	100	100
2	D	8/14 (57%)	8 (100%)	0	0	100	100
All	All	391/414 (94%)	384 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/165 (97%)	158 (99%)	2 (1%)	76	86
1	B	157/165 (95%)	151 (96%)	6 (4%)	40	51
2	C	11/14 (79%)	10 (91%)	1 (9%)	12	11
2	D	10/14 (71%)	10 (100%)	0	100	100
All	All	338/358 (94%)	329 (97%)	9 (3%)	57	65

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	165	PHE
1	A	176	VAL
1	B	158	LEU
1	B	165	PHE
1	B	170	ILE
1	B	203	GLN
1	B	228[A]	ASN
1	B	228[B]	ASN
2	C	471	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	253	ASN
1	B	171	ASN
1	B	265	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	188/193 (97%)	0.40	5 (2%) 58 69	54, 78, 104, 118	0
1	B	190/193 (98%)	0.38	5 (2%) 59 70	53, 77, 145, 158	0
2	C	11/14 (78%)	0.41	0 100 100	80, 84, 92, 98	0
2	D	10/14 (71%)	0.55	0 100 100	66, 74, 90, 100	0
All	All	399/414 (96%)	0.40	10 (2%) 61 72	53, 79, 134, 158	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	96	PHE	4.2
1	A	174	GLY	3.0
1	B	149	MET	3.0
1	A	175	ALA	2.9
1	B	211	LEU	2.6
1	A	109	LEU	2.5
1	A	96	PHE	2.5
1	B	95	LEU	2.4
1	B	243	ILE	2.3
1	A	199	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	ZN	A	602	1/1	0.99	0.16	0.12	82,82,82,82	0
3	ZN	B	602	1/1	0.97	0.16	-0.10	108,108,108,108	0
3	ZN	B	601	1/1	0.99	0.13	-1.38	90,90,90,90	0
3	ZN	A	601	1/1	0.99	0.09	-2.55	89,89,89,89	0

6.5 Other polymers [i](#)

There are no such residues in this entry.