



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 10:14 PM GMT

PDB ID : 4X4V
Title : Crystal structure of the *A.fulgidus* CCA-adding enzyme in complex with a human MenBeta minihelix ending in CCACC and AMPcPP
Authors : Kuhn, C.-D.; Joshua-Tor, L.
Deposited on : 2014-12-03
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)
RNA backbone	2183	1022 (3.00-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	457	% 93%
1	C	457	4% 89% 5% 6%
2	B	37	27% 73% 16% 11%
2	D	37	22% 51% 19% 30%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MG	A	502	-	-	-	X
4	MG	C	502	-	-	-	X
5	TAR	A	503	-	-	-	X
5	TAR	A	504	-	-	-	X
5	TAR	B	501	-	-	-	X
7	5GP	B	503	-	-	-	X
7	5GP	D	101	-	-	-	X

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 16922 atoms, of which 8000 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CCA-adding enzyme.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	443	7343	2354	3675	639	662	13	0	0	0
1	C	431	7161	2300	3583	623	643	12	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	438	SER	-	expression tag	UNP O28126
A	439	ASN	-	expression tag	UNP O28126
A	440	SER	-	expression tag	UNP O28126
A	441	SER	-	expression tag	UNP O28126
A	442	SER	-	expression tag	UNP O28126
A	443	VAL	-	expression tag	UNP O28126
A	444	ASP	-	expression tag	UNP O28126
A	445	LYS	-	expression tag	UNP O28126
A	446	LEU	-	expression tag	UNP O28126
A	447	ALA	-	expression tag	UNP O28126
A	448	ALA	-	expression tag	UNP O28126
A	449	ALA	-	expression tag	UNP O28126
A	450	LEU	-	expression tag	UNP O28126
A	451	GLU	-	expression tag	UNP O28126
A	452	HIS	-	expression tag	UNP O28126
A	453	HIS	-	expression tag	UNP O28126
A	454	HIS	-	expression tag	UNP O28126
A	455	HIS	-	expression tag	UNP O28126
A	456	HIS	-	expression tag	UNP O28126
A	457	HIS	-	expression tag	UNP O28126
C	438	SER	-	expression tag	UNP O28126
C	439	ASN	-	expression tag	UNP O28126
C	440	SER	-	expression tag	UNP O28126
C	441	SER	-	expression tag	UNP O28126
C	442	SER	-	expression tag	UNP O28126

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	443	VAL	-	expression tag	UNP O28126
C	444	ASP	-	expression tag	UNP O28126
C	445	LYS	-	expression tag	UNP O28126
C	446	LEU	-	expression tag	UNP O28126
C	447	ALA	-	expression tag	UNP O28126
C	448	ALA	-	expression tag	UNP O28126
C	449	ALA	-	expression tag	UNP O28126
C	450	LEU	-	expression tag	UNP O28126
C	451	GLU	-	expression tag	UNP O28126
C	452	HIS	-	expression tag	UNP O28126
C	453	HIS	-	expression tag	UNP O28126
C	454	HIS	-	expression tag	UNP O28126
C	455	HIS	-	expression tag	UNP O28126
C	456	HIS	-	expression tag	UNP O28126
C	457	HIS	-	expression tag	UNP O28126

- Molecule 2 is a RNA chain called Human MenBeta minihelix ending in CCACC.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P			
2	B	33	1100	323	371	132	240	34	0	1	0
2	D	26	836	245	284	97	184	26	0	0	0

- Molecule 3 is DIPHOSPHOMETHYLPHOSPHONIC ACID ADENOSYL ESTER (three-letter code: APC) (formula: C₁₁H₁₈N₅O₁₂P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
3	A	1	Total	C	H	N	O	P	0	0
			45	11	14	5	12	3		
3	C	1	Total	C	H	N	O	P	0	0
			45	11	14	5	12	3		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	C	1	Total	Mg	0	0
			1	1		

- Molecule 5 is D(-)-TARTARIC ACID (three-letter code: TAR) (formula: C₄H₆O₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			14	4	4	6		
5	A	1	Total	C	H	O	0	0
			14	4	4	6		
5	B	1	Total	C	H	O	0	0
			14	4	4	6		
5	C	1	Total	C	H	O	0	0
			14	4	4	6		

- Molecule 6 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			S
6	A	1	25	6	13	1	4	1	0	0

- Molecule 7 is GUANOSINE-5'-MONOPHOSPHATE (three-letter code: 5GP) (formula: $C_{10}H_{14}N_5O_8P$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	H	N	O	P		
7	B	1	35	10	11	5	8	1	0	0
7	B	1	29	10	9	5	5		0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
7	D	1	30	10	10	5	5	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	131	Total 131	O 131	0	0
8	B	21	Total 21	O 21	0	0
8	C	59	Total 59	O 59	0	0
8	D	4	Total 4	O 4	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CCA-adding enzyme




- Molecule 1: CCA-adding enzyme



- Molecule 2: Human MenBeta minihelix ending in CCACC



- Molecule 2: Human MenBeta minihelix ending in CCACC



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	111.06Å 215.60Å 58.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.68 – 2.60 29.68 – 2.59	Depositor EDS
% Data completeness (in resolution range)	98.5 (29.68-2.60) 98.5 (29.68-2.59)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.66 (at 2.61Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.184 , 0.237 0.199 , 0.245	Depositor DCC
R_{free} test set	2212 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	36.6	Xtrriage
Anisotropy	0.154	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 46.0	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Outliers	0 of 43901 reflections	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	16922	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: APC, MG, 5GP, MES, TAR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/3751	0.55	0/5040
1	C	0.38	0/3658	0.47	0/4913
2	B	0.57	1/812 (0.1%)	0.88	0/1261
2	D	0.58	1/611 (0.2%)	0.78	0/941
All	All	0.44	2/8832 (0.0%)	0.59	0/12155

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	G	OP3-P	-10.89	1.48	1.61
2	D	1	G	OP3-P	-10.03	1.49	1.61

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3668	3675	3668	10	0
1	C	3578	3583	3576	13	0
2	B	729	371	372	4	0
2	D	552	284	287	3	0
3	A	31	14	14	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	31	14	14	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	20	8	8	1	0
5	B	10	4	4	0	0
5	C	10	4	4	1	0
6	A	12	13	12	0	0
7	B	44	20	24	1	0
7	D	20	10	12	2	0
8	A	131	0	0	3	0
8	B	21	0	0	0	0
8	C	59	0	0	0	0
8	D	4	0	0	0	0
All	All	8922	8000	7995	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 28 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:ARG:NH2	8:A:623:HOH:O	2.15	0.78
1:C:30:GLU:OE2	1:C:34:ARG:NE	2.21	0.74
1:C:358:PHE:O	1:C:363:ARG:NH2	2.29	0.65
2:B:4[A]:G:H2'	2:B:5:C:C6	2.33	0.64
2:D:17:A:O4'	7:D:101:5GP:N2	2.31	0.64

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/457 (96%)	429 (97%)	10 (2%)	2 (0%)	34	60
1	C	427/457 (93%)	411 (96%)	15 (4%)	1 (0%)	52	77
All	All	868/914 (95%)	840 (97%)	25 (3%)	3 (0%)	46	72

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	120	PRO
1	C	119	GLU
1	A	119	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	393/404 (97%)	390 (99%)	3 (1%)	86	95
1	C	381/404 (94%)	379 (100%)	2 (0%)	92	98
All	All	774/808 (96%)	769 (99%)	5 (1%)	90	97

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	LEU
1	A	125	SER
1	A	137	LEU
1	C	118	LYS
1	C	145	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	30/37 (81%)	0	0
2	D	23/37 (62%)	2 (8%)	1 (4%)
All	All	53/74 (71%)	2 (3%)	1 (1%)

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	D	6	U
2	D	21	C

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	D	20	C

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	APC	A	501	4	25,33,33	2.22	5 (20%)	30,52,52	2.60	3 (10%)
5	TAR	A	503	-	3,9,9	0.44	0	6,12,12	0.92	0
5	TAR	A	504	-	3,9,9	0.61	0	6,12,12	0.92	0
6	MES	A	505	-	12,12,12	1.96	1 (8%)	16,16,16	1.80	6 (37%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	TAR	B	501	-	3,9,9	0.45	0	6,12,12	0.71	0
7	5GP	B	502	-	21,26,26	0.91	1 (4%)	23,40,40	1.49	2 (8%)
7	5GP	B	503	-	17,22,26	1.03	1 (5%)	17,33,40	1.14	2 (11%)
3	APC	C	501	4	25,33,33	2.20	5 (20%)	30,52,52	2.39	3 (10%)
5	TAR	C	503	-	3,9,9	0.60	0	6,12,12	0.42	0
7	5GP	D	101	-	17,22,26	0.79	0	17,33,40	1.07	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	APC	A	501	4	-	0/15/38/38	0/3/3/3
5	TAR	A	503	-	-	0/4/12/12	0/0/0/0
5	TAR	A	504	-	-	0/4/12/12	0/0/0/0
6	MES	A	505	-	-	0/6/14/14	1/1/1/1
5	TAR	B	501	-	-	0/4/12/12	0/0/0/0
7	5GP	B	502	-	-	0/6/26/26	0/3/3/3
7	5GP	B	503	-	-	0/2/22/26	0/3/3/3
3	APC	C	501	4	-	0/15/38/38	0/3/3/3
5	TAR	C	503	-	-	0/4/12/12	0/0/0/0
7	5GP	D	101	-	-	0/2/22/26	0/3/3/3

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	505	MES	C8-S	-5.91	1.67	1.78
3	C	501	APC	C2'-C3'	-2.95	1.45	1.53
3	A	501	APC	C2'-C3'	-2.77	1.45	1.53
3	A	501	APC	PB-O2B	-2.40	1.50	1.56
3	C	501	APC	PB-O2B	-2.13	1.51	1.56

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	APC	N3-C2-N1	-12.39	119.41	128.89
3	C	501	APC	N3-C2-N1	-11.10	120.40	128.89
3	C	501	APC	C4'-O4'-C1'	-4.63	104.63	109.72
3	A	501	APC	C4'-O4'-C1'	-4.59	104.67	109.72
7	B	502	5GP	C4-C5-N7	-3.29	106.45	109.48

There are no chirality outliers.

There are no torsion outliers.

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	505	MES	C2-C3-C5-C6-N4-O1

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	503	TAR	1	0
7	B	503	5GP	1	0
5	C	503	TAR	1	0
7	D	101	5GP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	443/457 (96%)	-0.27	6 (1%) 78 74	18, 30, 60, 90	0
1	C	431/457 (94%)	0.13	19 (4%) 38 30	23, 53, 83, 107	0
2	B	33/37 (89%)	1.38	10 (30%) 1 0	23, 62, 117, 120	0
2	D	26/37 (70%)	1.05	8 (30%) 1 0	41, 78, 130, 135	0
All	All	933/988 (94%)	0.01	43 (4%) 36 29	18, 39, 82, 135	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	27	G	5.9
2	B	5	C	5.6
2	B	18	G	5.2
1	C	22	ARG	5.2
2	B	6	U	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	MG	A	502	1/1	0.92	0.44	13.41	42,42,42,42	0
5	TAR	B	501	10/10	0.70	0.55	9.65	66,73,95,114	0
4	MG	C	502	1/1	0.81	0.50	9.45	71,71,71,71	0
5	TAR	A	503	10/10	0.90	0.47	5.83	48,56,68,77	0
5	TAR	A	504	10/10	0.82	0.34	3.59	63,73,86,87	0
7	5GP	D	101	20/24	0.44	0.48	1.92	54,61,73,79	30
7	5GP	B	503	20/24	0.69	0.41	1.79	48,57,70,70	29
5	TAR	C	503	10/10	0.85	0.20	1.70	35,58,76,78	0
6	MES	A	505	12/12	0.95	0.22	1.14	32,39,48,50	0
7	5GP	B	502	24/24	0.74	0.37	0.88	54,63,80,83	35
3	APC	C	501	31/31	0.95	0.23	0.54	47,63,77,89	0
3	APC	A	501	31/31	0.98	0.17	0.23	23,31,37,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.