



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:19 PM GMT

PDB ID : 4XEY
Title : Crystal structure of an SH2-kinase domain construct of c-Abl tyrosine kinase
Authors : Lorenz, S.; Deng, P.; Kuriyan, J.
Deposited on : 2014-12-25
Resolution : 2.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	408	<div> <div>5%</div> <div>60%</div> <div>6%</div> <div>34%</div> </div>
1	B	408	<div> <div>9%</div> <div>76%</div> <div>12%</div> <div>12%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5180 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein kinase ABL1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	359	Total	C	N	O	S	0	0	0
			2919	1872	489	541	17			
1	A	269	Total	C	N	O	S	0	0	0
			2195	1413	357	408	17			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	135	MET	-	expression tag	UNP P00519
B	136	ALA	-	expression tag	UNP P00519
B	137	SER	-	expression tag	UNP P00519
B	535	LEU	-	expression tag	UNP P00519
B	536	GLU	-	expression tag	UNP P00519
B	537	HIS	-	expression tag	UNP P00519
B	538	HIS	-	expression tag	UNP P00519
B	539	HIS	-	expression tag	UNP P00519
B	540	HIS	-	expression tag	UNP P00519
B	541	HIS	-	expression tag	UNP P00519
B	542	HIS	-	expression tag	UNP P00519
A	135	MET	-	expression tag	UNP P00519
A	136	ALA	-	expression tag	UNP P00519
A	137	SER	-	expression tag	UNP P00519
A	535	LEU	-	expression tag	UNP P00519
A	536	GLU	-	expression tag	UNP P00519
A	537	HIS	-	expression tag	UNP P00519
A	538	HIS	-	expression tag	UNP P00519
A	539	HIS	-	expression tag	UNP P00519
A	540	HIS	-	expression tag	UNP P00519
A	541	HIS	-	expression tag	UNP P00519
A	542	HIS	-	expression tag	UNP P00519

- Molecule 2 is N-(2-CHLORO-6-METHYLPHENYL)-2-({6-[4-(2-HYDROXYETHYL)PIPE RAZIN-1-YL]-2-METHYLPYRIMIDIN-4-YL}AMINO)-1,3-THIAZOLE-5-CARBOXAMID

E (three-letter code: 1N1) (formula: C₂₂H₂₆ClN₇O₂S).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	B	1	Total	C	Cl	N	O	S	0	0
			33	22	1	7	2	1		
2	A	1	Total	C	Cl	N	O	S	0	0
			33	22	1	7	2	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tyrosine-protein kinase ABL1



• Molecule 1: Tyrosine-protein kinase ABL1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	114.12Å 125.45Å 56.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.80 – 2.89 46.80 – 2.89	Depositor EDS
% Data completeness (in resolution range)	99.5 (46.80-2.89) 99.6 (46.80-2.89)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 2.91Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.231 , 0.259 0.237 , 0.267	Depositor DCC
R_{free} test set	958 reflections (5.41%)	DCC
Wilson B-factor (Å ²)	49.8	Xtriage
Anisotropy	0.796	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 38.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 18657 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	5180	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1N1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/2250	0.44	0/3044
1	B	0.31	0/2993	0.46	0/4051
All	All	0.28	0/5243	0.45	0/7095

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2195	0	2150	13	0
1	B	2919	0	2849	45	0
2	A	33	0	26	1	0
2	B	33	0	26	0	0
All	All	5180	0	5051	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:296:THR:HG22	1:A:298:GLU:H	1.13	1.09
1:B:325:THR:HG22	1:B:325:THR:O	1.69	0.90
1:B:258:ARG:NH2	1:B:324:CYS:HB3	1.86	0.89
1:B:303:LEU:HD21	1:B:325:THR:HG23	1.59	0.84
1:B:189:ARG:NH2	1:B:191:TYR:OH	2.12	0.83
1:B:303:LEU:HD22	1:B:325:THR:HG21	1.65	0.79
1:B:303:LEU:HD21	1:B:325:THR:CG2	2.16	0.76
1:B:303:LEU:CD2	1:B:325:THR:HG21	2.16	0.76
1:B:303:LEU:CD2	1:B:325:THR:CG2	2.69	0.69
1:B:325:THR:CG2	1:B:325:THR:O	2.42	0.67
1:A:296:THR:HG22	1:A:298:GLU:N	1.99	0.62
1:B:168:PHE:CD2	1:B:232:LEU:HD23	2.36	0.61
1:A:292:LEU:HD23	1:A:302:PHE:CD2	2.37	0.59
1:B:213:THR:HG22	1:B:214:LEU:N	2.17	0.58
1:B:170:VAL:HG13	1:B:182:ILE:HG22	1.86	0.58
1:B:258:ARG:HH21	1:B:324:CYS:HB3	1.67	0.57
1:B:150:PRO:HA	1:B:172:GLU:HG3	1.85	0.57
1:B:175:SER:O	1:B:194:ARG:NH1	2.40	0.55
1:A:487:VAL:O	1:A:491:MET:HG3	2.07	0.55
1:B:149:GLY:C	1:B:172:GLU:HG2	2.30	0.52
1:B:366:ILE:HD13	1:B:447:LEU:HD21	1.92	0.51
1:B:174:GLU:O	1:B:174:GLU:HG3	2.11	0.51
1:B:291:THR:OG1	1:B:292:LEU:N	2.44	0.50
1:B:176:SER:HB2	1:B:194:ARG:HH12	1.76	0.50
1:B:176:SER:HB3	1:B:194:ARG:NH2	2.26	0.50
1:B:258:ARG:NH1	1:B:327:GLU:OE2	2.45	0.50
1:B:221:HIS:HA	1:B:224:VAL:O	2.12	0.50
1:B:176:SER:CB	1:B:194:ARG:NH2	2.76	0.49
1:A:366:ILE:HD13	1:A:447:LEU:HD21	1.94	0.48
1:A:293:LYS:HG2	1:A:295:ASP:H	1.77	0.48
1:B:258:ARG:NH2	1:B:324:CYS:CB	2.70	0.48
1:B:213:THR:CG2	1:B:214:LEU:N	2.77	0.47
1:A:358:VAL:O	1:A:362:MET:HG3	2.14	0.47
1:B:258:ARG:HH21	1:B:324:CYS:CB	2.27	0.46
1:B:176:SER:HB2	1:B:194:ARG:NH1	2.31	0.46
1:B:258:ARG:HG3	1:B:331:TYR:CE2	2.51	0.46
1:B:181:SER:HB2	1:B:193:TYR:O	2.15	0.45
1:B:195:ILE:HD13	1:B:214:LEU:HD11	1.98	0.45
1:B:175:SER:HB3	1:B:194:ARG:HD2	1.99	0.45
1:A:291:THR:OG1	1:A:292:LEU:N	2.50	0.45
1:A:324:CYS:HB2	1:A:331:TYR:HB2	1.98	0.44
2:A:601:1N1:H161	2:A:601:1N1:H12	1.90	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:169:LEU:HD12	1:B:169:LEU:C	2.39	0.43
1:A:510:GLN:O	1:A:514:THR:HG23	2.18	0.43
1:B:220:HIS:O	1:B:223:THR:HG22	2.18	0.42
1:B:479:ARG:NE	1:B:483:CYS:O	2.50	0.42
1:A:296:THR:HG22	1:A:297:MET:N	2.35	0.42
1:B:204:TYR:HB3	1:B:210:ARG:HA	2.01	0.42
1:B:324:CYS:HB2	1:B:331:TYR:HB2	2.01	0.41
1:B:258:ARG:CZ	1:B:324:CYS:HB3	2.47	0.41
1:B:203:LEU:O	1:B:217:LEU:HD23	2.19	0.41
1:B:469:GLU:O	1:B:472:GLU:HB2	2.21	0.41
1:B:232:LEU:HD12	1:B:232:LEU:N	2.35	0.41
1:A:345:TYR:O	1:A:349:CYS:HB3	2.21	0.41
1:B:345:TYR:O	1:B:349:CYS:HB3	2.19	0.41
1:A:457:SER:HA	1:A:458:PRO:HD3	1.92	0.40
1:B:175:SER:CB	1:B:194:ARG:HD2	2.51	0.40
1:B:167:SER:HA	1:B:234:TYR:O	2.21	0.40
1:B:155:ALA:HA	1:B:158:TYR:CD2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/408 (65%)	257 (97%)	8 (3%)	0	100	100
1	B	349/408 (86%)	329 (94%)	20 (6%)	0	100	100
All	All	614/816 (75%)	586 (95%)	28 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/357 (66%)	236 (100%)	1 (0%)	93	98
1	B	316/357 (88%)	315 (100%)	1 (0%)	94	99
All	All	553/714 (78%)	551 (100%)	2 (0%)	93	98

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	400	ASP
1	A	400	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	509	HIS
1	A	509	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	1N1	A	601	-	32,36,36	0.86	2 (6%)	39,50,50	2.47	15 (38%)
2	1N1	B	601	-	32,36,36	0.87	2 (6%)	39,50,50	2.30	11 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1N1	A	601	-	-	0/13/29/29	0/4/4/4
2	1N1	B	601	-	-	0/13/29/29	0/4/4/4

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	1N1	C4-N2	-2.70	1.38	1.43
2	A	601	1N1	C4-N2	-2.61	1.38	1.43
2	B	601	1N1	C5-CL	2.27	1.79	1.73
2	A	601	1N1	C5-CL	2.27	1.79	1.73

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	1N1	C17-C16-N5	-3.63	103.43	110.63
2	A	601	1N1	N4-C14-N3	-3.30	119.10	125.58
2	B	601	1N1	N4-C14-N3	-3.20	119.29	125.58
2	B	601	1N1	C17-C16-N5	-2.84	104.99	110.63
2	A	601	1N1	N3-C13-N5	-2.71	113.56	116.63
2	B	601	1N1	C12-C11-N4	-2.66	117.29	123.43
2	A	601	1N1	C12-C11-N4	-2.52	117.63	123.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	1N1	O-C3-C2	-2.46	116.00	121.23
2	B	601	1N1	O-C3-C2	-2.39	116.14	121.23
2	A	601	1N1	C21-C20-N6	-2.31	104.82	113.41
2	A	601	1N1	C19-N5-C13	-2.18	115.03	120.22
2	A	601	1N1	C9-C4-N2	2.32	122.44	118.99
2	B	601	1N1	C8-C9-C4	2.44	121.58	117.78
2	A	601	1N1	C8-C9-C4	2.47	121.63	117.78
2	B	601	1N1	C15-C14-N4	2.54	121.53	117.20
2	A	601	1N1	C15-C14-N4	2.57	121.58	117.20
2	B	601	1N1	C14-N3-C13	3.31	118.29	115.64
2	A	601	1N1	C20-N6-C18	3.40	119.99	111.27
2	A	601	1N1	C14-N3-C13	3.62	118.53	115.64
2	B	601	1N1	C20-N6-C18	3.86	121.18	111.27
2	B	601	1N1	C2-C3-N2	4.60	120.44	113.75
2	A	601	1N1	C2-C3-N2	4.99	121.01	113.75
2	B	601	1N1	C18-N6-C17	5.96	121.81	108.90
2	B	601	1N1	C19-N5-C16	6.33	124.89	111.59
2	A	601	1N1	C18-N6-C17	6.72	123.45	108.90
2	A	601	1N1	C19-N5-C16	6.81	125.89	111.59

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	1N1	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	269/408 (65%)	0.52	21 (7%) 16 10	32, 44, 82, 98	0
1	B	359/408 (87%)	0.54	35 (9%) 10 6	24, 43, 101, 124	0
All	All	628/816 (76%)	0.53	56 (8%) 12 7	24, 43, 93, 124	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	198	ALA	8.3
1	B	197	THR	7.7
1	B	196	ASN	7.3
1	B	212	ASN	7.3
1	B	199	SER	5.6
1	B	201	GLY	5.3
1	B	251	TYR	4.7
1	B	226	ASP	4.7
1	B	177	PRO	4.6
1	A	259	THR	4.3
1	B	210	ARG	4.3
1	B	200	ASP	4.2
1	B	297	MET	3.9
1	A	258	ARG	3.8
1	B	176	SER	3.7
1	B	203	LEU	3.6
1	A	251	TYR	3.6
1	A	280	TRP	3.5
1	B	202	LYS	3.5
1	B	193	TYR	3.5
1	B	325	THR	3.5
1	B	296	THR	3.4
1	A	261	ILE	3.4
1	B	214	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	146	TRP	3.2
1	B	173	SER	3.2
1	B	195	ILE	3.2
1	B	174	GLU	3.2
1	A	522	SER	3.2
1	B	217	LEU	3.1
1	B	141	LEU	3.1
1	B	179	GLN	3.0
1	A	257	GLU	2.9
1	B	175	SER	2.9
1	B	204	TYR	2.9
1	A	294	GLU	2.9
1	A	256	MET	2.8
1	B	523	ASP	2.8
1	A	250	ASN	2.7
1	A	283	TYR	2.6
1	A	295	ASP	2.6
1	A	329	PRO	2.6
1	B	221	HIS	2.5
1	A	260	ASP	2.5
1	A	262	THR	2.4
1	A	328	PRO	2.3
1	A	521	ILE	2.3
1	B	181	SER	2.3
1	A	520	SER	2.3
1	B	148	HIS	2.2
1	A	276	TYR	2.2
1	B	211	PHE	2.2
1	A	325	THR	2.2
1	A	352	GLN	2.1
1	B	220	HIS	2.1
1	B	150	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	1N1	B	601	33/33	0.94	0.20	-0.00	29,34,45,54	0
2	1N1	A	601	33/33	0.91	0.22	-0.21	42,51,61,65	0

6.5 Other polymers [i](#)

There are no such residues in this entry.