



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 10:22 PM GMT

PDB ID : 4XKY  
Title : Structure of dihydrodipicolinate synthase from the commensal bacterium *Bacteroides thetaiotaomicron* at 2.1 Å resolution  
Authors : Mank, N.J.; Arnette, A.K.; Klapper, V.G.; Chruszcz, M.  
Deposited on : 2015-01-12  
Resolution : 2.10 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	 93% 7%
2	B	300	 93% 6%
2	C	300	 93% 6%
2	D	300	 93% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	A	401	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9698 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dihydrodipicolinate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	299	2288	1468	377	427	16	0	3	0

- Molecule 2 is a protein called Dihydrodipicolinate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	298	2298	1471	383	428	16	0	4	0
2	C	299	2284	1465	378	425	16	0	3	0
2	D	300	2308	1482	378	432	16	0	4	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	D	1	Total Na 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	127	Total O 127 127	0	0
5	B	167	Total O 167 167	0	0
5	C	77	Total O 77 77	0	0
5	D	144	Total O 144 144	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dihydrodipicolinate synthase




- Molecule 2: Dihydrodipicolinate synthase



- Molecule 2: Dihydrodipicolinate synthase



- Molecule 2: Dihydrodipicolinate synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.90Å 91.99Å 208.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.62 – 2.10 48.58 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.1 (48.62-2.10) 99.2 (48.58-2.10)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.71 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.185 , 0.221 0.192 , 0.226	Depositor DCC
$R_{free}$ test set	3845 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	33.3	Xtrriage
Anisotropy	0.317	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 39.4	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Outliers	1 of 76494 reflections (0.001%)	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9698	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MHO, ACT, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/2327	0.71	0/3165
2	B	0.57	0/2350	0.75	0/3196
2	C	0.46	0/2333	0.70	0/3174
2	D	0.57	0/2357	0.79	1/3205 (0.0%)
All	All	0.53	0/9367	0.74	1/12740 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	235	ARG	NE-CZ-NH2	-5.53	117.54	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2288	0	2292	11	0
2	B	2298	0	2304	10	0
2	C	2284	0	2276	10	0
2	D	2308	0	2305	11	0
3	A	4	0	3	0	0
4	D	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	127	0	0	1	0
5	B	167	0	0	3	0
5	C	77	0	0	1	0
5	D	144	0	0	1	0
All	All	9698	0	9180	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:138[B]:MET:SD	2:C:139:PRO:HD2	2.24	0.77
2:D:113:THR:HB	2:D:117:GLU:OE1	1.83	0.77
1:A:105:SER:HB3	1:A:133:LEU:HD11	1.73	0.71
2:D:256:SER:HA	5:D:566:HOH:O	1.94	0.67
2:D:105:SER:HB3	2:D:133:LEU:HD11	1.78	0.64
1:A:113:THR:CB	1:A:117:GLU:OE1	2.49	0.60
2:B:138:MET:HG3	2:B:139:PRO:HD2	1.83	0.60
1:A:167[A]:SER:O	1:A:168[A]:ALA:HB3	2.01	0.60
2:D:59:ARG:NH2	2:D:82:ASP:OD1	2.24	0.60
2:D:15:PRO:HB2	2:D:23:ASP:HB3	1.84	0.58
1:A:104:VAL:HG12	1:A:136:TYR:CD2	2.40	0.56
2:C:167[B]:SER:O	2:C:168[B]:ALA:HB3	2.06	0.56
2:D:217[B]:GLU:CD	2:D:217[B]:GLU:H	2.09	0.56
2:B:138:MET:HE1	5:B:530:HOH:O	2.07	0.53
2:B:15:PRO:HB2	2:B:23:ASP:HB3	1.90	0.53
2:C:235:ARG:HD3	5:C:441:HOH:O	2.09	0.51
2:C:60:MET:HG2	2:C:94:LEU:HD22	1.93	0.51
2:D:51:SER:OG	2:D:59:ARG:HD3	2.11	0.51
2:B:105:SER:HB3	2:B:133:LEU:HD11	1.94	0.50
2:D:10:PRO:HD2	2:D:41:ALA:O	2.11	0.50
2:C:105:SER:HB3	2:C:133:LEU:HD11	1.95	0.49
2:B:20:ASP:HA	2:B:58:LEU:HD22	1.95	0.49
1:A:138:MET:HB3	1:A:142:THR:HG23	1.95	0.49
1:A:15:PRO:HB2	1:A:23:ASP:HB3	1.95	0.48
1:A:28:GLN:HE21	1:A:68:ARG:HH22	1.62	0.47
2:D:252:GLY:O	2:D:253:GLN:CB	2.63	0.47
2:B:10:PRO:HD2	2:B:41:ALA:O	2.15	0.46
2:C:15:PRO:HB2	2:C:23:ASP:HB3	1.96	0.46
2:C:34:LEU:HD21	2:C:213:ASN:HA	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:153:ARG:O	2:C:156:GLU:HG2	2.14	0.46
2:B:104:VAL:HG12	2:B:136:TYR:CD2	2.52	0.45
2:B:9:ILE:O	2:B:208:VAL:HA	2.17	0.44
2:D:109:TYR:HB2	2:D:141:HIS:CB	2.48	0.43
1:A:190[A]:VAL:HG23	5:A:556:HOH:O	2.18	0.42
2:B:260:LYS:NZ	5:B:475:HOH:O	2.52	0.42
1:A:9:ILE:O	1:A:208:VAL:HA	2.19	0.42
1:A:167[A]:SER:O	1:A:168[A]:ALA:CB	2.67	0.42
2:C:16:LEU:CD2	2:C:58:LEU:HD21	2.50	0.42
2:B:181:LYS:HE3	5:B:555:HOH:O	2.20	0.41
2:D:75:PRO:HA	2:D:101:ASP:OD2	2.20	0.41
1:A:10:PRO:HD2	1:A:41:ALA:O	2.22	0.40
2:C:139:PRO:O	2:C:140:THR:C	2.59	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	300/300 (100%)	296 (99%)	4 (1%)	0	100	100
2	B	300/300 (100%)	294 (98%)	6 (2%)	0	100	100
2	C	300/300 (100%)	296 (99%)	4 (1%)	0	100	100
2	D	302/300 (101%)	295 (98%)	7 (2%)	0	100	100
All	All	1202/1200 (100%)	1181 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/250 (98%)	244 (100%)	1 (0%)	93	96
2	B	250/251 (100%)	246 (98%)	4 (2%)	70	76
2	C	244/251 (97%)	243 (100%)	1 (0%)	93	96
2	D	248/251 (99%)	246 (99%)	2 (1%)	86	91
All	All	987/1003 (98%)	979 (99%)	8 (1%)	86	91

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	170	THR
2	B	105	SER
2	B	145	ASN
2	B	167	SER
2	B	253	GLN
2	C	136	TYR
2	D	136	TYR
2	D	283	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	40	HIS
2	B	98	HIS
2	B	253	GLN
2	C	238	GLN
2	D	28	GLN
2	D	183	ASN
2	D	224	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	MHO	A	1	1	7,8,9	1.12	0	6,9,11	1.99	3 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MHO	A	1	1	-	0/5/7/9	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	MHO	O-C-CA	-2.34	119.40	125.49
1	A	1	MHO	CE-SD-CG	2.39	103.39	97.59
1	A	1	MHO	OD1-SD-CE	2.92	111.24	106.41

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ACT	A	401	-	1,3,3	1.89	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACT	A	401	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	298/300 (99%)	-0.30	3 (1%) 84 87	23, 37, 67, 101	0
2	B	298/300 (99%)	-0.35	4 (1%) 79 84	23, 30, 55, 101	0
2	C	299/300 (99%)	-0.19	8 (2%) 58 65	30, 45, 65, 111	0
2	D	300/300 (100%)	-0.20	10 (3%) 50 59	20, 31, 63, 104	0
All	All	1195/1200 (99%)	-0.26	25 (2%) 67 72	20, 36, 65, 111	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	255	GLY	6.0
2	C	142	THR	3.7
2	C	140	THR	3.4
1	A	254	HIS	3.3
2	C	139	PRO	3.3
1	A	109	TYR	3.2
2	C	141	HIS	3.1
2	D	252	GLY	3.0
2	D	254	HIS	3.0
2	B	113	THR	2.9
2	C	138[A]	MET	2.9
2	D	113	THR	2.8
2	C	156	GLU	2.6
2	D	142	THR	2.6
2	D	144	VAL	2.5
2	B	2	ASN	2.5
2	D	253	GLN	2.4
2	C	144	VAL	2.4
2	D	139	PRO	2.2
2	D	111	PHE	2.2
2	B	1	MET	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	253	GLN	2.2
1	A	141	HIS	2.1
2	C	299	GLY	2.1
2	D	112	ALA	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	MHO	A	1	9/10	0.47	0.24	-	64,72,88,94	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	ACT	A	401	4/4	0.92	0.22	5.09	43,47,48,52	0
4	NA	D	401	1/1	0.91	0.13	0.30	43,43,43,43	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.