



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 08:14 PM GMT

PDB ID : 4XZR
Title : Structure of yeast importin a bound to the membrane protein Nuclear Localization Signal sequence of INM protein Heh1
Authors : Lokareddy, R.K.; Cingolani, G.
Deposited on : 2015-02-04
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtrriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1640 (2.28-2.24)
Clashscore	102246	1095 (2.26-2.26)
Ramachandran outliers	100387	1063 (2.26-2.26)
Sidechain outliers	100360	1063 (2.26-2.26)
RSRZ outliers	91569	1647 (2.28-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	54	
2	B	423	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3546 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inner nuclear membrane protein SRC1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	A	18	148	87	31	30	0	0	0

- Molecule 2 is a protein called Importin subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	423	3285	2078	555	636	16	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	87	ALA	-	expression tag	UNP Q02821

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	5	5	5	0	0
3	B	108	108	108	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Inner nuclear membrane protein SRC1



- Molecule 2: Importin subunit alpha



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	129.68Å 58.31Å 95.37Å 90.00° 129.32° 90.00°	Depositor
Resolution (Å)	29.01 – 2.25 29.01 – 2.24	Depositor EDS
% Data completeness (in resolution range)	97.1 (29.01-2.25) 87.9 (29.01-2.24)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.11 (at 2.24Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: dev_1696)	Depositor
R, R_{free}	0.196 , 0.215 0.203 , 0.223	Depositor DCC
R_{free} test set	2423 reflections (4.97%)	DCC
Wilson B-factor (Å ²)	42.6	Xtrriage
Anisotropy	0.092	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 51.2	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Outliers	0 of 25273 reflections	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3546	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/147	1.03	2/190 (1.1%)
2	B	0.27	0/3337	0.43	2/4540 (0.0%)
All	All	0.28	0/3484	0.47	4/4730 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	4
All	All	0	5

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	176	LYS	N-CA-C	-8.63	87.69	111.00
2	B	120	ARG	C-N-CD	6.61	142.29	128.40
2	B	89	LEU	C-N-CD	6.60	142.25	128.40
1	A	176	LYS	N-CA-CB	6.58	122.44	110.60

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	175	LYS	Peptide
2	B	118	GLU	Peptide
2	B	119	HIS	Peptide
2	B	120	ARG	Peptide
2	B	89	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	148	0	149	17	0
2	B	3285	0	3340	45	1
3	A	5	0	0	0	0
3	B	108	0	0	1	0
All	All	3546	0	3489	54	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:175:LYS:HB3	1:A:176:LYS:HA	1.36	1.04
2:B:117:ARG:HB3	2:B:121:PRO:HG3	1.37	1.03
2:B:117:ARG:HD2	2:B:121:PRO:HG3	1.51	0.89
2:B:117:ARG:HB3	2:B:121:PRO:CG	2.12	0.79
1:A:175:LYS:HB3	1:A:176:LYS:CA	2.13	0.76
1:A:175:LYS:HE3	1:A:177:ARG:HE	1.53	0.74
2:B:393:GLU:HG3	2:B:394:VAL:HG13	1.77	0.67
2:B:117:ARG:HD2	2:B:121:PRO:CG	2.28	0.63
2:B:414:GLY:HA3	2:B:421:ILE:HG12	1.81	0.61
2:B:392:LEU:HD22	2:B:407:ILE:HD12	1.82	0.61
1:A:175:LYS:NZ	2:B:366:SER:OG	2.33	0.61
1:A:175:LYS:CB	1:A:176:LYS:HA	2.23	0.59
2:B:90:PRO:O	2:B:92:MET:N	2.39	0.56
1:A:195:LYS:HZ2	2:B:191:GLU:HG2	1.74	0.53
1:A:175:LYS:HE3	1:A:177:ARG:NE	2.23	0.53
2:B:279:TRP:CD2	2:B:318:PRO:HB3	2.43	0.53
2:B:117:ARG:CB	2:B:121:PRO:HG3	2.26	0.52
2:B:87:ALA:HB1	2:B:88:GLU:CD	2.30	0.51
1:A:175:LYS:CB	1:A:176:LYS:CA	2.83	0.51
2:B:119:HIS:O	2:B:120:ARG:HG3	2.11	0.50
2:B:90:PRO:C	2:B:92:MET:H	2.15	0.50
2:B:490:GLN:HB2	2:B:498:TYR:CE1	2.47	0.49
1:A:175:LYS:NZ	2:B:366:SER:CB	2.75	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:ASN:HD22	1:A:196:ILE:HG23	1.78	0.49
2:B:483:MET:HE2	2:B:509:PHE:HB3	1.95	0.48
2:B:92:MET:HA	2:B:95:GLN:HG2	1.95	0.48
2:B:117:ARG:HB3	2:B:121:PRO:CB	2.45	0.47
2:B:118:GLU:O	2:B:119:HIS:ND1	2.47	0.47
2:B:365:ILE:HG21	2:B:388:LEU:HD11	1.97	0.46
2:B:298:VAL:HG23	2:B:300:ILE:HG13	1.98	0.46
2:B:91:GLN:H	2:B:94:GLN:HB2	1.79	0.46
2:B:496:LYS:HD2	2:B:496:LYS:HA	1.54	0.46
2:B:496:LYS:HE2	2:B:500:LYS:H	1.80	0.46
1:A:176:LYS:HE3	2:B:329:GLY:O	2.16	0.46
1:A:175:LYS:CE	1:A:177:ARG:HE	2.27	0.45
1:A:196:ILE:HG13	1:A:196:ILE:H	1.62	0.45
2:B:490:GLN:HB2	2:B:498:TYR:CD1	2.52	0.44
1:A:178:LYS:NZ	2:B:287:GLY:O	2.43	0.44
2:B:218:GLU:HB3	2:B:219:PRO:HD3	1.99	0.44
2:B:390:LYS:O	2:B:394:VAL:HG22	2.18	0.43
2:B:397:TYR:OH	2:B:441:ASP:OD2	2.33	0.43
2:B:127:ILE:HD13	2:B:168:VAL:HG21	2.01	0.42
1:A:175:LYS:NZ	2:B:366:SER:HB2	2.34	0.42
2:B:117:ARG:HB3	2:B:121:PRO:HB3	2.00	0.42
1:A:175:LYS:HZ3	2:B:366:SER:HB2	1.83	0.42
2:B:279:TRP:CE2	2:B:318:PRO:HB3	2.54	0.42
1:A:194:ASN:OD1	2:B:117:ARG:HD3	2.18	0.42
2:B:463:LYS:HB2	2:B:468:LEU:HB2	2.02	0.42
2:B:120:ARG:O	2:B:120:ARG:NE	2.53	0.42
2:B:210:TYR:OH	3:B:693:HOH:O	2.21	0.41
2:B:91:GLN:HA	2:B:94:GLN:HB2	2.01	0.41
2:B:247:LYS:HA	2:B:248:PRO:HA	1.86	0.41
2:B:398:LYS:O	2:B:401:LYS:NZ	2.49	0.40
2:B:437:LEU:HD22	2:B:445:ILE:HG23	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:120:ARG:NH2	2:B:498:TYR:CE2[2_666]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	14/54 (26%)	12 (86%)	2 (14%)	0	100	100
2	B	421/423 (100%)	405 (96%)	16 (4%)	0	100	100
All	All	435/477 (91%)	417 (96%)	18 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	17/53 (32%)	15 (88%)	2 (12%)	6	4
2	B	364/364 (100%)	357 (98%)	7 (2%)	65	75
All	All	381/417 (91%)	372 (98%)	9 (2%)	57	67

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	174	ARG
1	A	189	SER
2	B	88	GLU
2	B	117	ARG
2	B	120	ARG
2	B	373	THR
2	B	412	SER
2	B	496	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	502	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	194	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	18/54 (33%)	1.29	4 (22%) 1 1	56, 90, 109, 111	0
2	B	423/423 (100%)	0.41	36 (8%) 13 14	34, 54, 112, 133	0
All	All	441/477 (92%)	0.44	40 (9%) 11 13	34, 54, 111, 133	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	87	ALA	9.2
2	B	469	ASN	7.1
2	B	470	ILE	7.0
2	B	91	GLN	6.9
2	B	498	TYR	5.1
2	B	468	LEU	5.0
2	B	118	GLU	5.0
2	B	502	TYR	4.9
2	B	467	GLY	4.6
2	B	487	PHE	4.5
1	A	173	THR	4.4
2	B	493	GLU	4.3
2	B	94	GLN	3.9
2	B	465	ALA	3.7
2	B	119	HIS	3.7
1	A	197	ASP	3.7
2	B	101	MET	3.7
2	B	466	ARG	3.5
2	B	398	LYS	3.4
2	B	100	ASP	3.3
2	B	464	GLU	3.3
2	B	494	ASN	3.0
2	B	120	ARG	3.0
2	B	489	CYS	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	89	LEU	2.8
2	B	90	PRO	2.8
1	A	174	ARG	2.8
2	B	281	ILE	2.8
2	B	495	ASP	2.7
2	B	102	GLN	2.6
1	A	189	SER	2.6
2	B	504	ILE	2.6
2	B	497	ILE	2.5
2	B	323	VAL	2.4
2	B	388	LEU	2.4
2	B	506	GLU	2.4
2	B	490	GLN	2.4
2	B	463	LYS	2.4
2	B	88	GLU	2.1
2	B	481	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.