



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 08:18 PM GMT

PDB ID : 4XZS
Title : Crystal Structure of TRIAP1-MBP fusion
Authors : Miliara, X.; Garnett, J.A.; Abid-Ali, F.; Perez-Dorado, I.; Matthews, S.J.
Deposited on : 2015-02-04
Resolution : 2.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4587 (2.14-2.10)
Clashscore	102246	5132 (2.14-2.10)
Ramachandran outliers	100387	5080 (2.14-2.10)
Sidechain outliers	100360	5081 (2.14-2.10)
RSRZ outliers	91569	4597 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	446	 86% 8% 6%
1	B	446	 85% 9% 6%

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 6794 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein, TP53-regulated inhibitor of apoptosis 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	421	3249	2083	531	624	11	0	4	0
1	B	420	3233	2071	531	620	11	0	2	0

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP P0AEX9
A	82	ALA	ASP	conflict	UNP P0AEX9
A	83	ALA	LYS	conflict	UNP P0AEX9
A	172	ALA	GLU	conflict	UNP P0AEX9
A	173	ALA	ASN	conflict	UNP P0AEX9
A	239	ALA	LYS	conflict	UNP P0AEX9
A	359	ALA	GLU	conflict	UNP P0AEX9
A	362	ALA	LYS	conflict	UNP P0AEX9
A	363	ALA	ASP	conflict	UNP P0AEX9
A	367	ASN	-	linker	UNP P0AEX9
A	368	ALA	-	linker	UNP P0AEX9
A	369	ALA	-	linker	UNP P0AEX9
A	370	ALA	-	linker	UNP P0AEX9
B	0	MET	-	initiating methionine	UNP P0AEX9
B	82	ALA	ASP	conflict	UNP P0AEX9
B	83	ALA	LYS	conflict	UNP P0AEX9
B	172	ALA	GLU	conflict	UNP P0AEX9
B	173	ALA	ASN	conflict	UNP P0AEX9
B	239	ALA	LYS	conflict	UNP P0AEX9
B	359	ALA	GLU	conflict	UNP P0AEX9
B	362	ALA	LYS	conflict	UNP P0AEX9
B	363	ALA	ASP	conflict	UNP P0AEX9
B	367	ASN	-	linker	UNP P0AEX9
B	368	ALA	-	linker	UNP P0AEX9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	369	ALA	-	linker	UNP P0AEX9
B	370	ALA	-	linker	UNP P0AEX9

- Molecule 2 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 12 6 6	0	0
2	A	1	Total C O 11 6 5	0	0
2	B	1	Total C O 12 6 6	0	0
2	B	1	Total C O 11 6 5	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	141	Total O 141 141	0	0
3	B	125	Total O 125 125	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltose-binding periplasmic protein,TP53-regulated inhibitor of apoptosis 1



- Molecule 1: Maltose-binding periplasmic protein,TP53-regulated inhibitor of apoptosis 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	75.39Å 56.01Å 100.28Å 90.00° 106.35° 90.00°	Depositor
Resolution (Å)	48.41 – 2.12 48.41 – 2.12	Depositor EDS
% Data completeness (in resolution range)	98.6 (48.41-2.12) 98.6 (48.41-2.12)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 2.12Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.196 , 0.242 0.204 , 0.244	Depositor DCC
R_{free} test set	2246 reflections (5.22%)	DCC
Wilson B-factor (Å ²)	28.7	Xtrriage
Anisotropy	0.547	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 43.8	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Outliers	0 of 45295 reflections	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6794	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 73.80 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.7217e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.65	0/3339	0.70	1/4539 (0.0%)
1	B	0.61	0/3314	0.70	0/4505
All	All	0.63	0/6653	0.70	1/9044 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	209	ASP	CB-CG-OD1	5.51	123.26	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3249	0	3162	25	0
1	B	3233	0	3140	28	0
2	A	23	0	21	0	0
2	B	23	0	21	2	0
3	A	141	0	0	3	0
3	B	125	0	0	3	0
All	All	6794	0	6344	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:170:LYS:HE2	1:B:180:ASP:OD2	1.69	0.91
1:B:170:LYS:CE	1:B:180:ASP:OD2	2.38	0.72
1:B:247:LEU:H	1:B:323:ASN:HD21	1.40	0.70
1:B:89:LEU:HD23	1:B:107:PRO:HG2	1.75	0.68
1:B:79:ILE:HD12	1:B:81:PRO:HD3	1.75	0.68
1:A:157:THR:HG23	3:A:676:HOH:O	1.93	0.67
1:A:247:LEU:H	1:A:323:ASN:HD21	1.41	0.66
1:B:5:GLY:H	1:B:272:ASN:HD21	1.44	0.66
1:A:181:VAL:O	1:A:365:GLN:NE2	2.29	0.65
1:B:153:GLU:CD	1:B:344:ARG:HH12	2.00	0.65
1:A:5:GLY:H	1:A:272:ASN:HD21	1.46	0.64
1:B:65:ASP:OD2	2:B:502:GLC:O3	2.15	0.61
1:A:196:VAL:HG12	1:A:200:LYS:HD2	1.82	0.60
1:B:171:TYR:HB2	1:B:176:TYR:CE1	2.39	0.58
1:B:373:VAL:HG11	1:B:420:ALA:HB2	1.86	0.57
1:B:337:SER:HB2	3:B:707:HOH:O	2.05	0.56
1:B:229:PRO:HA	1:B:232:TRP:CE2	2.42	0.54
1:B:5:GLY:H	1:B:272:ASN:ND2	2.07	0.53
1:A:5:GLY:H	1:A:272:ASN:ND2	2.08	0.51
1:B:153:GLU:OE2	1:B:344:ARG:NH1	2.44	0.50
1:A:183:VAL:H	1:A:365:GLN:HE22	1.59	0.49
1:B:68:GLY:HA3	1:B:332:ASN:O	2.14	0.48
1:A:68:GLY:HA3	1:A:332:ASN:O	2.14	0.47
1:B:381:LYS:HB2	1:B:413:TYR:CE1	2.50	0.47
1:A:411:LYS:O	1:A:415:GLN:HG3	2.14	0.46
1:B:340:TRP:CD1	2:B:502:GLC:H4	2.50	0.46
1:A:144:LYS:NZ	3:A:603:HOH:O	2.50	0.45
1:B:198:LEU:HD13	1:B:204:MET:CE	2.46	0.44
1:A:181:VAL:O	1:A:365:GLN:OE1	2.35	0.44
1:B:132:ILE:HD13	1:B:147:LEU:HD22	1.99	0.44
1:A:184:ASP:HB2	1:A:365:GLN:HB2	1.98	0.44
1:B:337:SER:HB3	3:B:612:HOH:O	2.17	0.44
1:A:205[B]:ASN:ND2	1:A:207:ASP:OD1	2.51	0.44
1:A:47:PHE:HB2	1:A:60:ILE:HD12	2.00	0.43
1:A:62:TRP:CD1	1:A:63:ALA:N	2.86	0.43
1:A:321:MET:HA	1:A:321:MET:CE	2.49	0.43
1:B:158:TRP:N	1:B:159:PRO:CD	2.81	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:391:TRP:O	1:A:395:LYS:HB2	2.18	0.43
1:A:183:VAL:HG12	1:A:365:GLN:NE2	2.33	0.42
1:A:229:PRO:HA	1:A:232:TRP:CE2	2.54	0.42
1:A:79:ILE:HD12	1:A:81:PRO:HD3	2.02	0.42
1:B:205:ASN:HB2	1:B:208:THR:HG23	2.02	0.42
1:A:136:ASP:HA	1:A:146:ALA:HB2	2.01	0.42
1:B:373:VAL:CG1	1:B:420:ALA:HB2	2.50	0.41
1:A:66:ARG:NE	3:A:605:HOH:O	2.53	0.41
1:B:171:TYR:CD1	1:B:176:TYR:CZ	3.07	0.41
1:B:361:LEU:HD23	1:B:361:LEU:HA	1.91	0.41
1:B:172:ALA:O	1:B:173:ALA:HB3	2.21	0.41
1:A:288:GLU:CD	1:A:288:GLU:H	2.23	0.41
1:B:62:TRP:CD1	1:B:63:ALA:N	2.89	0.41
1:A:158:TRP:HH2	1:A:167:TYR:O	2.04	0.40
1:A:181:VAL:O	1:A:365:GLN:CD	2.59	0.40
1:B:26:LYS:HE3	3:B:608:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	423/446 (95%)	417 (99%)	6 (1%)	0	100	100
1	B	420/446 (94%)	416 (99%)	4 (1%)	0	100	100
All	All	843/892 (94%)	833 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	329/353 (93%)	328 (100%)	1 (0%)	94	97
1	B	326/353 (92%)	325 (100%)	1 (0%)	94	97
All	All	655/706 (93%)	653 (100%)	2 (0%)	94	97

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	258	PHE
1	B	258	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	272	ASN
1	A	323	ASN
1	A	365	GLN
1	A	389	ASN
1	A	418	GLN
1	B	39	HIS
1	B	253	GLN
1	B	272	ASN
1	B	323	ASN
1	B	389	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GLC	A	501	2	12,12,12	0.69	0	17,17,17	1.76	4 (23%)
2	GLC	A	502	2	11,11,12	0.77	0	15,15,17	1.16	2 (13%)
2	GLC	B	501	2	12,12,12	0.43	0	17,17,17	0.50	0
2	GLC	B	502	2	11,11,12	0.25	0	15,15,17	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	501	2	-	0/2/22/22	0/1/1/1
2	GLC	A	502	2	-	0/2/19/22	0/1/1/1
2	GLC	B	501	2	-	0/2/22/22	0/1/1/1
2	GLC	B	502	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	502	GLC	O3-C3-C2	-2.85	104.78	110.01
2	A	501	GLC	O3-C3-C4	2.32	115.59	110.36
2	A	502	GLC	C1-O5-C5	2.56	115.91	112.14
2	A	501	GLC	C1-C2-C3	2.62	114.97	110.68
2	A	501	GLC	C1-O5-C5	3.57	120.37	113.54
2	A	501	GLC	O5-C1-C2	4.07	117.14	110.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	502	GLC	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	421/446 (94%)	-0.17	5 (1%) 81 85	23, 38, 68, 92	0
1	B	420/446 (94%)	-0.09	4 (0%) 84 87	28, 42, 70, 93	0
All	All	841/892 (94%)	-0.13	9 (1%) 82 86	23, 40, 69, 93	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	421	ILE	2.7
1	B	173	ALA	2.5
1	A	394	GLU	2.2
1	A	365	GLN	2.2
1	B	52	ALA	2.1
1	A	419	LYS	2.1
1	B	394	GLU	2.0
1	B	403	GLY	2.0
1	A	406	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q < 0.9
2	GLC	A	501	12/12	0.94	0.12	0.91	21,25,27,27	0
2	GLC	B	501	12/12	0.93	0.11	0.55	22,26,30,33	0
2	GLC	B	502	11/12	0.90	0.12	0.53	18,20,21,21	0
2	GLC	A	502	11/12	0.98	0.06	-1.51	15,18,20,20	0

6.5 Other polymers

There are no such residues in this entry.