



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 06:43 AM GMT

PDB ID : 2Y3X
Title : Catalytic domain of mouse 2',3'-cyclic nucleotide 3'- phosphodiesterase, complexed with sulfate
Authors : Myllykoski, M.; Kursula, P.
Deposited on : 2011-01-04
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)

There are 4 unique types of molecules in this entry. The entry contains 5198 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 2', 3'-CYCLIC-NUCLEOTIDE 3'-PHOSPHODIESTERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	215	Total	C 1671	N 1078	O 281	S 307	5	0	0
1	B	215	Total	C 1672	N 1079	O 281	S 307	5	0	0
1	E	210	Total	C 1634	N 1056	O 275	S 298	5	0	0

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	158	GLY	-	EXPRESSION TAG	UNP P16330
B	158	GLY	-	EXPRESSION TAG	UNP P16330
E	158	GLY	-	EXPRESSION TAG	UNP P16330

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	58	Total O 58 58	0	0
4	B	96	Total O 96 96	0	0
4	E	5	Total O 5 5	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 2', 3'-CYCLIC-NUCLEOTIDE 3'-PHOSPHODIESTERASE



- Molecule 1: 2', 3'-CYCLIC-NUCLEOTIDE 3'-PHOSPHODIESTERASE



- Molecule 1: 2', 3'-CYCLIC-NUCLEOTIDE 3'-PHOSPHODIESTERASE



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	82.43Å 82.43Å 86.31Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	71.40 – 2.10 32.98 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (71.40-2.10) 99.8 (32.98-2.10)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.57 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R , R_{free}	0.190 , 0.228 0.223 , 0.259	Depositor DCC
R_{free} test set	1981 reflections (5.46%)	DCC
Wilson B-factor (Å ²)	39.8	Xtriage
Anisotropy	0.308	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 28.0	EDS
Estimated twinning fraction	0.788 for H, K, L 0.212 for -H, H+K, -L 0.068 for -h,-k,l 0.085 for h,-h-k,-l 0.159 for -k,-h,-l	Xtriage
Reported twinning fraction	0.788 for H, K, L 0.212 for -H, H+K, -L	Depositor
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtriage
Outliers	0 of 38236 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5198	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.74	0 / 1710	0.92	3 / 2303 (0.1%)
1	B	0.99	3 / 1709 (0.2%)	0.85	0 / 2299
1	E	0.39	0 / 1671	0.80	3 / 2249 (0.1%)
All	All	0.75	3 / 5090 (0.1%)	0.86	6 / 6851 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	E	0	2
All	All	0	4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	314	CYS	CB-SG	-9.27	1.66	1.82
1	B	179	GLU	CG-CD	5.41	1.60	1.51
1	B	236	CYS	CB-SG	-5.33	1.73	1.81

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	348	ARG	NE-CZ-NH2	-20.06	110.27	120.30
1	E	348	ARG	NE-CZ-NH2	-19.00	110.80	120.30
1	A	348	ARG	NE-CZ-NH1	15.96	128.28	120.30
1	E	348	ARG	NE-CZ-NH1	15.20	127.90	120.30
1	A	348	ARG	CD-NE-CZ	9.40	136.76	123.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	E	348	ARG	CD-NE-CZ	8.67	135.74	123.60

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	340	GLY	Peptide
1	A	348	ARG	Sidechain
1	E	340	GLY	Peptide
1	E	348	ARG	Sidechain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1671	0	1694	17	1
1	B	1672	0	1697	16	0
1	E	1634	0	1660	21	1
2	A	25	0	0	0	0
2	B	25	0	0	0	0
3	B	12	0	16	4	0
4	A	58	0	0	0	0
4	B	96	0	0	4	0
4	E	5	0	0	1	0
All	All	5198	0	5067	49	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (\AA)	Clash overlap (\AA)
1:A:329:ASP:OD2	1:A:358:ARG:NH1	2.11	0.82
1:A:334:VAL:HG23	1:A:339:GLN:HB3	1.62	0.81
1:E:334:VAL:HG12	1:E:339:GLN:HB3	1.61	0.81

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:329:ASP:OD2	1:E:358:ARG:NH1	2.18	0.76
3:B:1380:GOL:H11	4:B:2069:HOH:O	1.87	0.74
1:A:182:ARG:HH12	1:A:229:LEU:HD21	1.52	0.73
1:A:333:GLN:OE1	1:A:358:ARG:NH2	2.29	0.66
1:B:333:GLN:HG3	1:B:358:ARG:HH12	1.59	0.66
1:E:182:ARG:HH12	1:E:229:LEU:HD21	1.61	0.65
3:B:1380:GOL:C1	4:B:2069:HOH:O	2.46	0.64
1:A:211:PRO:O	1:A:213:GLU:HG2	2.00	0.61
1:E:333:GLN:OE1	1:E:358:ARG:NH2	2.33	0.60
1:B:224:ARG:HB2	4:B:2066:HOH:O	2.03	0.58
1:A:278:GLN:HG3	1:A:327:LEU:HD21	1.86	0.58
1:E:213:GLU:OE1	4:E:2002:HOH:O	2.17	0.57
1:E:331:LEU:HA	1:E:334:VAL:HG22	1.85	0.57
1:B:183:LYS:HE2	1:B:187:VAL:CG2	2.33	0.57
1:B:333:GLN:HG3	1:B:358:ARG:NH1	2.20	0.55
1:B:290:PRO:HG2	1:B:293:LEU:HD21	1.89	0.55
1:A:255:ARG:HD3	1:E:296:PRO:CG	2.37	0.54
1:E:333:GLN:HE21	1:E:338:SER:HB3	1.74	0.53
1:A:255:ARG:HG3	1:E:296:PRO:HB2	1.91	0.52
1:A:333:GLN:HE21	1:A:338:SER:HB3	1.75	0.52
1:B:330:ILE:O	1:B:334:VAL:HG23	2.10	0.52
1:B:183:LYS:HD3	4:B:2007:HOH:O	2.10	0.51
1:E:278:GLN:HG3	1:E:327:LEU:HD21	1.92	0.51
1:A:262:LYS:NZ	1:E:246:GLU:HB3	2.27	0.50
1:B:183:LYS:HE2	1:B:187:VAL:HG21	1.93	0.50
1:B:183:LYS:CE	1:B:187:VAL:HG21	2.42	0.49
1:B:370:LYS:HD2	3:B:1379:GOL:H11	1.96	0.48
1:A:182:ARG:NH1	1:A:229:LEU:HD21	2.25	0.47
1:A:255:ARG:HD3	1:E:296:PRO:HG3	1.97	0.47
1:E:330:ILE:HG23	1:E:354:LEU:HD13	1.96	0.46
1:E:302:LEU:HD22	1:E:332:GLN:NE2	2.31	0.46
1:E:183:LYS:O	1:E:187:VAL:HG23	2.16	0.46
1:A:182:ARG:NH1	1:A:224:ARG:HH11	2.14	0.45
1:B:213:GLU:OE2	1:E:317:ASP:HA	2.15	0.45
1:A:343:VAL:HG11	1:A:353:SER:HB2	1.98	0.45
1:B:183:LYS:HE2	1:B:187:VAL:HG23	1.99	0.44
1:A:183:LYS:O	1:A:187:VAL:HG23	2.17	0.44
1:B:179:GLU:HA	1:B:179:GLU:OE1	2.18	0.44
1:A:262:LYS:HZ1	1:E:246:GLU:HB3	1.83	0.43
1:A:281:LEU:HD22	1:A:285:GLU:HB3	2.01	0.43
1:E:339:GLN:HA	1:E:354:LEU:HD23	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:214:LYS:HE2	1:E:317:ASP:OD2	2.19	0.43
1:E:346:LEU:HD11	1:E:351:LEU:HD22	2.02	0.42
1:B:174:THR:OG1	3:B:1379:GOL:H12	2.21	0.41
1:B:339:GLN:NE2	1:B:353:SER:HB3	2.35	0.41
1:E:202:LEU:HD13	1:E:217:LEU:HD21	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:LYS:NZ	1:E:205:PHE:O[3_664]	2.00	0.20

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	213/221 (96%)	210 (99%)	3 (1%)	0	100 100
1	B	211/221 (96%)	203 (96%)	8 (4%)	0	100 100
1	E	206/221 (93%)	202 (98%)	4 (2%)	0	100 100
All	All	630/663 (95%)	615 (98%)	15 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	176/180 (98%)	172 (98%)	4 (2%)	58	62
1	B	176/180 (98%)	174 (99%)	2 (1%)	80	85
1	E	172/180 (96%)	163 (95%)	9 (5%)	29	25
All	All	524/540 (97%)	509 (97%)	15 (3%)	50	53

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	182	ARG
1	A	249	GLN
1	A	334	VAL
1	A	338	SER
1	B	224	ARG
1	B	295	LYS
1	E	165	LEU
1	E	182	ARG
1	E	202	LEU
1	E	249	GLN
1	E	293	LEU
1	E	296	PRO
1	E	302	LEU
1	E	338	SER
1	E	348	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	1379	-	4,4,4	0.35	0	6,6,6	0.43	0
2	SO4	A	1380	-	4,4,4	1.13	0	6,6,6	1.10	1 (16%)
2	SO4	A	1381	-	4,4,4	0.32	0	6,6,6	0.17	0
2	SO4	A	1382	-	4,4,4	0.24	0	6,6,6	0.12	0
2	SO4	A	1383	-	4,4,4	0.25	0	6,6,6	0.25	0
3	GOL	B	1379	-	5,5,5	0.41	0	5,5,5	0.42	0
3	GOL	B	1380	-	5,5,5	0.34	0	5,5,5	0.90	0
2	SO4	B	1381	-	4,4,4	0.22	0	6,6,6	0.15	0
2	SO4	B	1382	-	4,4,4	0.83	0	6,6,6	0.68	0
2	SO4	B	1383	-	4,4,4	0.34	0	6,6,6	0.37	0
2	SO4	B	1384	-	4,4,4	0.28	0	6,6,6	0.26	0
2	SO4	B	1385	-	4,4,4	0.27	0	6,6,6	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	1379	-	-	0/0/0/0	0/0/0/0
2	SO4	A	1380	-	-	0/0/0/0	0/0/0/0
2	SO4	A	1381	-	-	0/0/0/0	0/0/0/0
2	SO4	A	1382	-	-	0/0/0/0	0/0/0/0
2	SO4	A	1383	-	-	0/0/0/0	0/0/0/0
3	GOL	B	1379	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1380	-	-	0/4/4/4	0/0/0/0
2	SO4	B	1381	-	-	0/0/0/0	0/0/0/0
2	SO4	B	1382	-	-	0/0/0/0	0/0/0/0
2	SO4	B	1383	-	-	0/0/0/0	0/0/0/0
2	SO4	B	1384	-	-	0/0/0/0	0/0/0/0
2	SO4	B	1385	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	1380	SO4	O4-S-O3	2.20	117.94	108.98

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1379	GOL	2	0
3	B	1380	GOL	2	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	215/221 (97%)	0.69	9 (4%) 40 49	31, 50, 75, 87	0
1	B	215/221 (97%)	0.59	6 (2%) 56 64	24, 39, 67, 88	0
1	E	210/221 (95%)	4.64	174 (82%) 0 0	92, 124, 164, 186	0
All	All	640/663 (96%)	1.95	189 (29%) 1 1	24, 54, 149, 186	0

All (189) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	207	SER	18.0
1	E	291	SER	14.8
1	E	203	ARG	14.3
1	E	204	HIS	12.4
1	E	206	ILE	11.9
1	E	318	VAL	11.2
1	E	214	LYS	10.9
1	E	224	ARG	10.4
1	E	315	ALA	10.4
1	E	234	LYS	10.3
1	E	202	LEU	10.3
1	E	248	ALA	10.3
1	E	200	LYS	10.2
1	E	331	LEU	9.9
1	E	317	ASP	9.6
1	E	330	ILE	9.6
1	E	198	PHE	9.5
1	E	319	GLN	9.5
1	E	289	TRP	9.2
1	E	195	HIS	9.2
1	E	360	MET	8.9
1	E	164	PHE	8.7
1	E	288	LEU	8.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	295	LYS	8.3
1	E	247	TYR	8.2
1	E	377	TYR	8.2
1	B	160	LEU	8.2
1	E	205	PHE	8.0
1	E	223	LYS	7.8
1	E	225	PRO	7.8
1	E	336	GLY	7.8
1	E	298	ALA	7.8
1	E	263	LEU	7.7
1	E	253	VAL	7.7
1	E	217	LEU	7.6
1	E	286	LEU	7.6
1	E	371	ALA	7.4
1	E	251	GLU	7.4
1	E	293	LEU	7.4
1	E	294	ASP	7.4
1	E	270	VAL	7.3
1	E	303	PRO	7.3
1	E	213	GLU	7.1
1	E	348	ARG	7.0
1	E	235	PHE	6.9
1	E	297	SER	6.8
1	E	340	GLY	6.8
1	E	238	TYR	6.8
1	E	323	THR	6.7
1	E	165	LEU	6.6
1	E	281	LEU	6.5
1	E	378	GLY	6.5
1	E	199	LYS	6.5
1	E	359	TRP	6.5
1	E	357	GLY	6.3
1	E	313	GLY	6.3
1	E	171	TRP	6.3
1	E	287	GLN	6.3
1	E	375	GLY	6.1
1	E	322	GLN	6.1
1	E	276	GLY	6.1
1	E	306	SER	5.8
1	E	259	LYS	5.8
1	E	172	PHE	5.8
1	E	316	ALA	5.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	337	GLY	5.6
1	E	354	LEU	5.6
1	E	376	TYR	5.6
1	A	338	SER	5.6
1	E	342	ALA	5.6
1	E	220	TYR	5.5
1	E	245	GLU	5.5
1	E	279	VAL	5.5
1	E	261	PHE	5.3
1	B	225	PRO	5.3
1	E	192	LEU	5.3
1	B	164	PHE	5.3
1	E	310	VAL	5.2
1	E	175	LYS	5.2
1	E	233	THR	5.1
1	E	304	PRO	5.1
1	E	283	ASP	5.1
1	E	215	LEU	5.1
1	E	290	PRO	5.1
1	E	346	LEU	5.0
1	E	364	THR	5.0
1	E	280	VAL	5.0
1	E	355	GLY	5.0
1	E	320	PRO	4.6
1	E	169	PHE	4.5
1	E	244	ALA	4.5
1	E	335	LYS	4.5
1	E	356	LYS	4.5
1	E	373	PHE	4.5
1	E	334	VAL	4.4
1	E	243	GLY	4.4
1	E	242	ALA	4.3
1	E	312	LEU	4.3
1	E	307	ARG	4.2
1	E	240	LYS	4.2
1	E	328	LEU	4.1
1	E	321	VAL	4.1
1	E	201	GLU	4.1
1	E	370	LYS	4.0
1	E	363	LEU	4.0
1	E	221	PHE	3.9
1	E	324	GLY	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	299	SER	3.9
1	E	369	VAL	3.9
1	E	258	GLY	3.8
1	E	329	ASP	3.8
1	E	262	LYS	3.8
1	E	231	CYS	3.7
1	E	257	TYR	3.7
1	E	341	GLU	3.7
1	E	351	LEU	3.7
1	E	228	VAL	3.7
1	E	269	PHE	3.7
1	E	343	VAL	3.7
1	E	285	GLU	3.7
1	E	196	LYS	3.6
1	E	275	ALA	3.6
1	E	167	LEU	3.6
1	E	182	ARG	3.5
1	E	197	ALA	3.5
1	E	358	ARG	3.4
1	E	292	ASP	3.4
1	E	296	PRO	3.4
1	B	161	GLU	3.3
1	E	255	ARG	3.3
1	E	216	GLU	3.3
1	E	252	VAL	3.3
1	E	176	LYS	3.3
1	E	219	SER	3.1
1	E	227	GLY	3.1
1	E	361	LEU	3.1
1	E	268	LEU	3.1
1	E	277	ALA	3.0
1	E	352	TYR	3.0
1	E	362	SER	3.0
1	A	214	LYS	3.0
1	E	265	ILE	2.9
1	E	273	LYS	2.9
1	E	282	THR	2.9
1	E	174	THR	2.8
1	E	333	GLN	2.8
1	E	246	GLU	2.8
1	E	327	LEU	2.8
1	E	249	GLN	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	284	GLN	2.8
1	E	353	SER	2.8
1	E	267	ALA	2.8
1	E	272	PRO	2.8
1	E	367	MET	2.7
1	E	226	PRO	2.7
1	E	372	ILE	2.7
1	A	209	ASP	2.7
1	B	163	ASP	2.7
1	E	266	SER	2.7
1	A	215	LEU	2.6
1	E	314	CYS	2.6
1	E	236	CYS	2.6
1	E	302	LEU	2.6
1	B	207	SER	2.5
1	E	300	GLU	2.5
1	E	187	VAL	2.5
1	E	278	GLN	2.5
1	E	190	GLU	2.5
1	E	229	LEU	2.5
1	A	224	ARG	2.4
1	E	347	PRO	2.4
1	A	245	GLU	2.3
1	E	264	SER	2.3
1	E	325	LEU	2.3
1	E	232	THR	2.2
1	E	189	LEU	2.2
1	E	194	ASN	2.2
1	E	193	GLY	2.2
1	A	295	LYS	2.2
1	A	340	GLY	2.1
1	E	338	SER	2.1
1	E	186	GLN	2.1
1	A	378	GLY	2.1
1	E	326	ASP	2.1
1	E	301	GLY	2.0
1	E	168	TYR	2.0
1	E	365	LYS	2.0
1	E	183	LYS	2.0
1	E	332	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	SO4	A	1381	5/5	0.72	0.21	0.71	100,101,101,103	0
2	SO4	A	1380	5/5	0.98	0.18	0.70	35,42,52,55	0
2	SO4	B	1383	5/5	0.89	0.15	0.12	78,80,83,86	0
3	GOL	B	1380	6/6	0.87	0.13	-0.21	43,50,54,59	0
2	SO4	A	1379	5/5	0.94	0.15	-0.31	67,67,71,71	0
2	SO4	B	1382	5/5	0.98	0.13	-0.45	44,48,53,55	0
3	GOL	B	1379	6/6	0.81	0.18	-0.56	66,70,73,74	0
2	SO4	B	1381	5/5	0.95	0.10	-0.81	71,74,77,78	0
2	SO4	A	1383	5/5	0.86	0.18	-	97,97,100,101	0
2	SO4	B	1384	5/5	0.92	0.10	-	92,94,96,96	0
2	SO4	A	1382	5/5	0.69	0.21	-	131,132,132,133	0
2	SO4	B	1385	5/5	0.90	0.21	-	84,87,89,90	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.