



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 06:45 AM GMT

PDB ID : 2Y9Y
Title : Chromatin Remodeling Factor ISW1a(del_ATPase)
Authors : Yamada, K.; Frouws, T.D.; Angst, B.; Fitzgerald, D.J.; Deluca, C.; Schimmele, K.; Sargent, D.F.; Richmond, T.J.
Deposited on : 2011-02-17
Resolution : 3.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1624 (3.32-3.20)
Clashscore	102246	1806 (3.32-3.20)
Ramachandran outliers	100387	1773 (3.32-3.20)
Sidechain outliers	100360	1771 (3.32-3.20)
RSRZ outliers	91569	1632 (3.32-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	374	<div> <div>7%</div> <div>48%</div> <div>24%</div> <div>•</div> <div>26%</div> </div>
2	B	624	<div> <div>10%</div> <div>55%</div> <div>36%</div> <div>•</div> <div>5%</div> </div>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 7170 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IMITATION SWITCH PROTEIN 1 (DEL_ATPASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	277	Total	C	N	O	S	0	0	0
			2317	1473	399	437	8			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	762	MET	-	EXPRESSION TAG	UNP P38144
A	815	GLN	LYS	ENGINEERED MUTATION	UNP P38144
A	844	LYS	ASP	ENGINEERED MUTATION	UNP P38144
A	848	GLN	GLU	ENGINEERED MUTATION	UNP P38144
A	851	GLN	LYS	ENGINEERED MUTATION	UNP P38144
A	853	GLU	GLN	ENGINEERED MUTATION	UNP P38144
A	1130	HIS	-	EXPRESSION TAG	UNP P38144
A	1131	HIS	-	EXPRESSION TAG	UNP P38144
A	1132	HIS	-	EXPRESSION TAG	UNP P38144
A	1133	HIS	-	EXPRESSION TAG	UNP P38144
A	1134	HIS	-	EXPRESSION TAG	UNP P38144
A	1135	HIS	-	EXPRESSION TAG	UNP P38144

- Molecule 2 is a protein called ISWI ONE COMPLEX PROTEIN 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	592	Total	C	N	O	S	0	0	0
			4853	3123	818	897	15			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	126	MET	-	EXPRESSION TAG	UNP P43596
B	682	LYS	ASN	ENGINEERED MUTATION	UNP P43596


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: IMITATION SWITCH PROTEIN 1 (DEL_ATPASE)



• Molecule 2: ISWI ONE COMPLEX PROTEIN 3





4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	206.97Å 206.97Å 215.31Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.87 – 3.25 29.87 – 3.25	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.87-3.25) 100.0 (29.87-3.25)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.93 (at 3.24Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.283 , 0.296 0.275 , 0.278	Depositor DCC
R_{free} test set	1380 reflections (4.93%)	DCC
Wilson B-factor (Å ²)	120.0	Xtriage
Anisotropy	0.190	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 97.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 28010 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7170	wwPDB-VP
Average B, all atoms (Å ²)	153.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/2368	0.35	0/3186
2	B	0.21	0/4974	0.35	0/6723
All	All	0.21	0/7342	0.35	0/9909

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2317	0	2300	87	0
2	B	4853	0	4857	204	0
All	All	7170	0	7157	287	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 287 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:327:LYS:HB2	2:B:328:PRO:HD3	1.50	0.91
1:A:918:LYS:HB2	1:A:922:GLU:HG3	1.53	0.89

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:293:LYS:HA	2:B:296:SER:HB3	1.54	0.87
2:B:339:THR:HB	2:B:340:PRO:HD3	1.61	0.83
2:B:413:LEU:HD22	2:B:533:LEU:HB2	1.58	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/374 (74%)	214 (78%)	52 (19%)	9 (3%)	5	32
2	B	588/624 (94%)	481 (82%)	92 (16%)	15 (3%)	7	39
All	All	863/998 (86%)	695 (80%)	144 (17%)	24 (3%)	6	37

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	799	PRO
1	A	916	PRO
1	A	904	ARG
1	A	793	PRO
2	B	162	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/343 (74%)	250 (98%)	4 (2%)	70	89
2	B	543/571 (95%)	514 (95%)	29 (5%)	28	68
All	All	797/914 (87%)	764 (96%)	33 (4%)	37	75

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	319	LEU
2	B	373	ARG
2	B	716	ILE
2	B	330	THR
2	B	353	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	335	GLN
2	B	407	HIS
2	B	726	GLN
2	B	353	ASN
2	B	415	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	277/374 (74%)	0.50	28 (10%) 9 6	91, 154, 262, 349	0
2	B	592/624 (94%)	0.54	60 (10%) 9 6	79, 132, 259, 319	0
All	All	869/998 (87%)	0.53	88 (10%) 9 6	79, 138, 261, 349	0

The worst 5 of 88 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	274	SER	11.6
2	B	139	GLU	10.0
2	B	469	LYS	9.1
2	B	275	ALA	8.5
2	B	470	GLU	8.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.