



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 11:42 PM GMT

PDB ID : 1YEE
Title : STRUCTURE OF A CATALYTIC ANTIBODY, IGG2A FAB FRAGMENT (D2.5)
Authors : Golinelli-Pimpaneau, B.; Knossow, M.
Deposited on : 1997-04-15
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	L	219	
2	H	222	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3495 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IGG2A FAB FRAGMENT (D2.5).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	219	Total	C	N	O	S	0	0	0
			1690	1061	283	339	7			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	2	ILE	VAL	CONFLICT	PIR S16112
L	7	SER	THR	CONFLICT	PIR S16112
L	10	THR	SER	CONFLICT	PIR S16112
L	12	SER	PRO	CONFLICT	PIR S16112
L	14	THR	SER	CONFLICT	PIR S16112
L	15	ILE	LEU	CONFLICT	PIR S16112
L	17	GLN	ASP	CONFLICT	PIR S16112
L	18	PRO	GLN	CONFLICT	PIR S16112
L	24	LYS	ARG	CONFLICT	PIR S16112
L	27B	LEU	VAL	CONFLICT	PIR S16112
L	27D	TYR	HIS	CONFLICT	PIR S16112
L	30	LYS	ASN	CONFLICT	PIR S16112
L	34	SER	TYR	CONFLICT	PIR S16112
L	36	LEU	TYR	CONFLICT	PIR S16112
L	39	ARG	LYS	CONFLICT	PIR S16112
L	46	ARG	PRO	CONFLICT	PIR S16112
L	50	LEU	ARG	CONFLICT	PIR S16112
L	53	LYS	ASN	CONFLICT	PIR S16112
L	54	LEU	ARG	CONFLICT	PIR S16112
L	55	ASP	PHE	CONFLICT	PIR S16112
L	63	THR	SER	CONFLICT	PIR S16112
L	81	ALA	GLU	CONFLICT	PIR S16112
L	85	LEU	VAL	CONFLICT	PIR S16112
L	87	TYR	PHE	CONFLICT	PIR S16112
L	89	VAL	PHE	CONFLICT	PIR S16112
L	94	PHE	VAL	CONFLICT	PIR S16112
L	103	LYS	ARG	CONFLICT	PIR S16112

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
L	107	LEU	LYS	CONFLICT	PIR S16112

- Molecule 2 is a protein called IGG2A FAB FRAGMENT (D2.5).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	222	Total	C	N	O	S	0	0	0
			1688	1081	273	328	6			

- Molecule 3 is 4-NITRO-BENZYLPHOSPHONOBUTANOYL-GLYCINE (three-letter code: PNB) (formula: $C_{13}H_{17}N_2O_8P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	H	1	Total	C	N	O	P	0	0
			24	13	2	8	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	49	Total	O	0	0
			49	49		
4	L	44	Total	O	0	0
			44	44		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


• Molecule 1: IGG2A FAB FRAGMENT (D2.5)

Chain L: 



• Molecule 2: IGG2A FAB FRAGMENT (D2.5)

Chain H: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.93 Å 76.98 Å 45.80 Å 90.00° 94.68° 90.00°	Depositor
Resolution (Å)	7.00 – 2.20	Depositor
% Data completeness (in resolution range)	95.0 (7.00-2.20)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.204 , 0.260	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3495	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PNB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.64	0/1727	0.86	3/2345 (0.1%)
2	H	0.61	0/1734	0.88	1/2370 (0.0%)
All	All	0.63	0/3461	0.87	4/4715 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	11	LEU	CA-CB-CG	8.82	135.58	115.30
1	L	83	LEU	CA-CB-CG	5.65	128.30	115.30
1	L	36	LEU	CA-CB-CG	5.52	128.00	115.30
2	H	108	LEU	CA-CB-CG	5.03	126.86	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1690	0	1654	44	0
2	H	1688	0	1665	44	0
3	H	24	0	15	0	0
4	H	49	0	0	3	0
4	L	44	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3495	0	3334	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:195:GLU:HG2	1:L:206:VAL:HG22	1.53	0.88
1:L:117:ILE:H	2:H:129:GLY:HA3	1.38	0.86
2:H:18:VAL:HG22	2:H:82(C):LEU:HD11	1.59	0.85
2:H:169:VAL:HG12	2:H:189:VAL:HG23	1.60	0.83
2:H:119:PRO:HB3	2:H:145:TYR:HB3	1.67	0.76
1:L:160:LEU:HD23	2:H:177:GLN:NE2	2.02	0.74
2:H:126:PRO:CB	2:H:130:ASP:HB3	2.20	0.70
1:L:160:LEU:HD21	2:H:175:VAL:HG11	1.73	0.70
2:H:126:PRO:HB2	2:H:130:ASP:HB3	1.76	0.67
1:L:160:LEU:HD21	2:H:175:VAL:CG1	2.25	0.66
2:H:108:LEU:HD13	4:H:556:HOH:O	1.96	0.65
1:L:50:LEU:HB2	1:L:53:LYS:HD3	1.78	0.64
1:L:147:LYS:HE2	1:L:154:GLU:HG3	1.79	0.62
1:L:39:ARG:HG2	4:L:245:HOH:O	1.98	0.62
1:L:167:ASP:OD1	1:L:169:LYS:HG2	2.02	0.60
2:H:170:HIS:HD2	4:H:571:HOH:O	1.84	0.60
1:L:11:LEU:HD13	1:L:13:VAL:HG23	1.84	0.59
1:L:147:LYS:HE3	1:L:148:TRP:H	1.67	0.59
1:L:146:VAL:HA	1:L:195:GLU:O	2.02	0.59
2:H:201:GLN:HE22	2:H:203:ILE:HG13	1.69	0.57
2:H:13:ARG:HD3	2:H:112:SER:O	2.06	0.55
1:L:119:PRO:HG2	2:H:127:VAL:HG21	1.87	0.55
2:H:38:LYS:HB2	2:H:48:ILE:HD11	1.88	0.55
2:H:201:GLN:HE22	2:H:203:ILE:CG1	2.21	0.53
1:L:108:ARG:NH1	1:L:111:ALA:HB2	2.24	0.53
2:H:61:VAL:HG13	2:H:64:LYS:CE	2.39	0.52
1:L:117:ILE:N	2:H:129:GLY:HA3	2.18	0.52
1:L:160:LEU:CD2	2:H:175:VAL:HG11	2.39	0.52
1:L:183:LYS:O	1:L:187:GLU:HG3	2.10	0.52
2:H:12:VAL:HG21	2:H:18:VAL:CG1	2.41	0.50
2:H:64:LYS:HD3	4:H:570:HOH:O	2.12	0.50
1:L:108:ARG:HH12	1:L:111:ALA:HB2	1.77	0.50
2:H:40:ARG:O	2:H:42:ALA:N	2.44	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:24:LYS:HE2	1:L:70:ASP:OD1	2.11	0.49
2:H:131:THR:HG21	2:H:135:SER:OG	2.12	0.49
2:H:196:THR:O	2:H:201:GLN:HG3	2.12	0.49
2:H:61:VAL:HG13	2:H:64:LYS:HE2	1.95	0.49
1:L:53:LYS:N	1:L:53:LYS:HD2	2.28	0.49
1:L:2:ILE:HG13	1:L:93:HIS:CD2	2.47	0.49
2:H:12:VAL:HG21	2:H:18:VAL:HG13	1.94	0.48
2:H:12:VAL:HG11	2:H:82(C):LEU:HD12	1.95	0.48
2:H:126:PRO:HB3	2:H:130:ASP:HB3	1.93	0.48
1:L:119:PRO:HG2	2:H:127:VAL:CG2	2.43	0.48
1:L:136:LEU:HD13	1:L:175:MET:HG2	1.96	0.48
1:L:147:LYS:HE3	1:L:148:TRP:N	2.29	0.47
1:L:142:LYS:HB3	1:L:142:LYS:HE2	1.59	0.47
1:L:39:ARG:HB2	1:L:42:GLN:HG3	1.96	0.47
1:L:160:LEU:HD23	2:H:177:GLN:CD	2.34	0.47
2:H:97:PHE:O	2:H:99:PRO:HD3	2.15	0.46
1:L:147:LYS:HE2	1:L:154:GLU:CG	2.45	0.46
1:L:143:ASP:O	1:L:198:HIS:HD2	1.99	0.45
2:H:172:PHE:HA	2:H:173:PRO:HD3	1.84	0.45
1:L:212:ASN:O	1:L:213:GLU:HG2	2.18	0.44
1:L:40:PRO:HG2	1:L:83:LEU:CD2	2.47	0.44
1:L:74:LYS:HG3	4:L:224:HOH:O	2.17	0.44
1:L:117:ILE:HG22	2:H:129:GLY:N	2.32	0.43
1:L:120:PRO:HG2	1:L:130:ALA:HB1	2.01	0.43
1:L:11:LEU:HD13	1:L:13:VAL:CG2	2.49	0.43
2:H:131:THR:HB	2:H:135:SER:O	2.18	0.42
2:H:196:THR:HG22	2:H:201:GLN:CD	2.40	0.42
1:L:169:LYS:HD3	1:L:169:LYS:N	2.35	0.42
2:H:133:GLY:O	2:H:134:SER:HB3	2.20	0.42
1:L:189:HIS:O	1:L:211:ARG:HD3	2.19	0.42
1:L:193:THR:HG23	1:L:206:VAL:HG13	2.02	0.42
1:L:2:ILE:CD1	1:L:27:GLN:NE2	2.83	0.41
2:H:201:GLN:O	2:H:201:GLN:OE1	2.39	0.41
2:H:19:LYS:HB3	2:H:19:LYS:HE2	1.70	0.41
2:H:130:ASP:O	2:H:131:THR:CB	2.69	0.41
1:L:27(D):TYR:HB3	1:L:28:ASN:OD1	2.21	0.41
1:L:110:ASP:HB3	1:L:200:THR:HG22	2.03	0.41
1:L:117:ILE:HG22	2:H:129:GLY:CA	2.51	0.40
2:H:196:THR:O	2:H:201:GLN:OE1	2.40	0.40
2:H:62:LYS:HE2	2:H:62:LYS:HB3	1.94	0.40
1:L:117:ILE:HG22	2:H:129:GLY:HA3	2.04	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:119:PRO:CG	2:H:127:VAL:HB	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	217/219 (99%)	211 (97%)	5 (2%)	1 (0%)	34	35
2	H	220/222 (99%)	208 (94%)	7 (3%)	5 (2%)	8	4
All	All	437/441 (99%)	419 (96%)	12 (3%)	6 (1%)	14	10

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	41	ALA
2	H	42	ALA
2	H	131	THR
2	H	132	THR
2	H	130	ASP
1	L	51	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	195/195 (100%)	177 (91%)	18 (9%)	11	11
2	H	190/190 (100%)	171 (90%)	19 (10%)	9	8
All	All	385/385 (100%)	348 (90%)	37 (10%)	10	9

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	11	LEU
1	L	14	THR
1	L	27(C)	LEU
1	L	36	LEU
1	L	37	LEU
1	L	39	ARG
1	L	42	GLN
1	L	43	SER
1	L	47	LEU
1	L	85	LEU
1	L	89	VAL
1	L	103	LYS
1	L	105	GLU
1	L	108	ARG
1	L	147	LYS
1	L	155	ARG
1	L	175	MET
1	L	207	LYS
2	H	1	GLU
2	H	5	GLN
2	H	11	LEU
2	H	17	SER
2	H	18	VAL
2	H	61	VAL
2	H	64	LYS
2	H	100	VAL
2	H	100(E)	VAL
2	H	100(F)	LEU
2	H	108	LEU
2	H	115	LYS
2	H	132	THR
2	H	137	THR
2	H	138	LEU
2	H	164	LEU
2	H	169	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	H	194	SER
2	H	201	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	27	GLN
1	L	156	GLN
2	H	170	HIS
2	H	177	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PNB	H	551	-	19,24,24	3.23	5 (26%)	25,32,32	2.26	10 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PNB	H	551	-	-	0/20/22/22	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	551	PNB	P1-C8	-9.05	1.73	1.80
3	H	551	PNB	P1-O3	-6.86	1.39	1.56
3	H	551	PNB	C12-N2	-5.89	1.34	1.46
3	H	551	PNB	P1-O2	-4.42	1.40	1.51
3	H	551	PNB	P1-O1	2.10	1.59	1.57

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	551	PNB	C5-C4-N1	-5.22	115.27	119.48
3	H	551	PNB	C8-C9-C10	-3.30	103.45	112.11
3	H	551	PNB	O1-P1-O2	-3.00	105.99	113.98
3	H	551	PNB	C2-C3-C4	-2.25	117.00	120.15
3	H	551	PNB	C5-C6-C1	-2.21	118.01	121.04
3	H	551	PNB	O3-P1-O2	2.25	117.20	110.12
3	H	551	PNB	C10-C11-N2	2.83	121.37	116.46
3	H	551	PNB	P1-C8-C9	2.92	119.67	114.67
3	H	551	PNB	C9-C10-C11	3.20	121.72	113.24
3	H	551	PNB	C3-C4-N1	4.06	122.76	119.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.