



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 19, 2017 – 01:04 PM EST

PDB ID : 2YFD
Title : STRUCTURAL AND FUNCTIONAL INSIGHTS OF DR2231 PROTEIN, THE MAZG-LIKE NUCLEOSIDE TRIPHOSPHATE PYROPHOSPHOHYDROLASE FROM DEINOCOCCUS RADIODURANS, COMPLEXED WITH Mg and dUMP
Authors : Goncalves, A.M.D.; De Sanctis, D.; Mcsweeney, S.M.
Deposited on : 2011-04-05
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028442
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1609 (1.76-1.76)
Clashscore	102246	1730 (1.76-1.76)
Ramachandran outliers	100387	1711 (1.76-1.76)
Sidechain outliers	100360	1711 (1.76-1.76)
RSRZ outliers	91569	1610 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	154	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> % </div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 77% 12% 10% </div> </div>
1	B	154	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> 4% </div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 79% 5% 16% </div> </div>
1	C	154	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> 2% </div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 81% 8% 11% </div> </div>
1	D	154	<div> <div style="width: 100%; height: 10px; background-color: red; position: relative;"> 7% </div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> 78% 6% 16% </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACT	B	1146	-	-	-	X
5	ACT	D	1148	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8777 atoms, of which 4163 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MAZG-LIKE NUCLEOSIDE TRIPHOSPHATE PYROPHOSPHOHYDROLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	138	Total	C	H	N	O	0	2	0	
			2135	666	1064	204	201				
1	B	130	Total	C	H	N	O	0	0	0	
			1980	623	984	184	189				
1	C	137	Total	C	H	N	O	S	0	4	0
			2136	668	1063	201	203	1			
1	D	130	Total	C	H	N	O	S	0	8	0
			2048	640	1017	194	196	1			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	EXPRESSION TAG	UNP Q9RS96
A	-4	ILE	-	EXPRESSION TAG	UNP Q9RS96
A	-3	ASP	-	EXPRESSION TAG	UNP Q9RS96
A	-2	PRO	-	EXPRESSION TAG	UNP Q9RS96
A	-1	PHE	-	EXPRESSION TAG	UNP Q9RS96
A	0	THR	-	EXPRESSION TAG	UNP Q9RS96
B	-5	GLY	-	EXPRESSION TAG	UNP Q9RS96
B	-4	ILE	-	EXPRESSION TAG	UNP Q9RS96
B	-3	ASP	-	EXPRESSION TAG	UNP Q9RS96
B	-2	PRO	-	EXPRESSION TAG	UNP Q9RS96
B	-1	PHE	-	EXPRESSION TAG	UNP Q9RS96
B	0	THR	-	EXPRESSION TAG	UNP Q9RS96
C	-5	GLY	-	EXPRESSION TAG	UNP Q9RS96
C	-4	ILE	-	EXPRESSION TAG	UNP Q9RS96
C	-3	ASP	-	EXPRESSION TAG	UNP Q9RS96
C	-2	PRO	-	EXPRESSION TAG	UNP Q9RS96
C	-1	PHE	-	EXPRESSION TAG	UNP Q9RS96
C	0	THR	-	EXPRESSION TAG	UNP Q9RS96
D	-5	GLY	-	EXPRESSION TAG	UNP Q9RS96
D	-4	ILE	-	EXPRESSION TAG	UNP Q9RS96

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-3	ASP	-	EXPRESSION TAG	UNP Q9RS96
D	-2	PRO	-	EXPRESSION TAG	UNP Q9RS96
D	-1	PHE	-	EXPRESSION TAG	UNP Q9RS96
D	0	THR	-	EXPRESSION TAG	UNP Q9RS96

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0

- Molecule 3 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (three-letter code: UMP) (formula: C₉H₁₃N₂O₈P).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	P	0	0
			31	9	11	2	8	1		
3	C	1	Total	C	H	N	O	P	0	0
			30	9	10	2	8	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		
4	D	1	Total	Cl	0	0
			1	1		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	H	O	0	0
			7	2	3	2		
5	D	1	Total	C	H	O	0	0
			7	2	3	2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	D	1	Total	C	H	O	0	0
			14	3	8	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	102	Total	O	0	0
			102	102		
7	B	98	Total	O	0	0
			98	98		
7	C	97	Total	O	0	0
			97	97		
7	D	86	Total	O	0	0
			86	86		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: MAZG-LIKE NUCLEOSIDE TRIPHOSPHATE PYROPHOSPHOHYDROLASE




• Molecule 1: MAZG-LIKE NUCLEOSIDE TRIPHOSPHATE PYROPHOSPHOHYDROLASE



• Molecule 1: MAZG-LIKE NUCLEOSIDE TRIPHOSPHATE PYROPHOSPHOHYDROLASE



• Molecule 1: MAZG-LIKE NUCLEOSIDE TRIPHOSPHATE PYROPHOSPHOHYDROLASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	78.09 Å 149.53 Å 52.38 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.93 – 1.77 43.50 – 1.77	Depositor EDS
% Data completeness (in resolution range)	93.9 (19.93-1.77) 93.9 (43.50-1.77)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 1.77 Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.184 , 0.211 0.184 , 0.184	Depositor DCC
R_{free} test set	2901 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	18.6	Xtriage
Anisotropy	0.269	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 47.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8777	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.19 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.3643e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, ACT, UMP, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	1/1104 (0.1%)	0.55	0/1499
1	B	0.46	0/1017	0.52	0/1385
1	C	0.70	1/1109 (0.1%)	0.63	1/1508 (0.1%)
1	D	0.36	0/1077	0.48	0/1466
All	All	0.56	2/4307 (0.0%)	0.55	1/5858 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	117	ARG	CB-CG	-5.26	1.38	1.52
1	A	145	PRO	CA-CB	-5.20	1.43	1.53

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	13	ARG	NE-CZ-NH1	5.78	123.19	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1071	1064	1045	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	996	984	983	6	0
1	C	1073	1063	1043	10	0
1	D	1031	1017	982	8	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	20	11	10	0	0
3	C	20	10	10	0	0
4	A	1	0	0	1	0
4	D	1	0	0	0	0
5	B	4	3	3	0	0
5	D	4	3	3	0	0
6	D	6	8	8	0	0
7	A	102	0	0	2	0
7	B	98	0	0	0	0
7	C	97	0	0	0	0
7	D	86	0	0	1	0
All	All	4614	4163	4087	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:28:ARG:HH12	1:C:143:HIS:CD2	1.93	0.86
4:A:1148:CL:CL	7:A:2024:HOH:O	2.48	0.69
1:A:16:GLU:HA	1:A:19:ARG:HE	1.59	0.66
1:A:16:GLU:CB	1:A:19:ARG:HH21	2.13	0.62
1:A:28:ARG:NH2	1:A:144:ALA:O	2.33	0.60
1:A:130:ARG:HD3	7:A:2089:HOH:O	2.01	0.59
1:A:104:GLU:HG3	1:A:132:ALA:HB1	1.86	0.58
1:A:74:ALA:HA	1:B:138:ILE:CD1	2.35	0.56
1:C:74:ALA:CB	1:D:135:ARG:HG2	2.37	0.55
1:C:104:GLU:HG3	1:C:132:ALA:HB1	1.88	0.54
1:C:35:GLU:HA	1:C:38:ARG:NH1	2.24	0.53
1:A:28:ARG:NH1	1:A:144:ALA:O	2.42	0.53
1:C:54[A]:GLU:HG3	1:C:76:LEU:HB2	1.92	0.52
1:C:28:ARG:HH12	1:C:143:HIS:CG	2.27	0.51
1:A:75:PRO:O	1:A:79:GLU:HG2	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:36:LEU:C	1:B:36:LEU:HD23	2.33	0.49
1:A:16:GLU:HB2	1:A:19:ARG:HH21	1.77	0.49
1:A:35:GLU:H	1:A:35:GLU:CD	2.15	0.49
1:D:138:ILE:O	1:D:142:GLN:HG3	2.13	0.48
1:A:138:ILE:O	1:A:142:GLN:HG2	2.14	0.48
1:A:16:GLU:HA	1:A:19:ARG:HH21	1.78	0.48
1:C:35:GLU:H	1:C:35:GLU:CD	2.19	0.45
1:C:54[A]:GLU:HG2	1:C:76:LEU:N	2.31	0.45
1:C:74:ALA:HB3	1:D:135:ARG:HG2	1.99	0.44
1:C:54[A]:GLU:HG2	1:C:76:LEU:CA	2.47	0.44
1:B:138:ILE:O	1:B:142:GLN:HG3	2.19	0.43
1:A:16:GLU:CA	1:A:19:ARG:HH21	2.31	0.43
1:B:25:THR:HG23	1:B:25:THR:O	2.19	0.42
1:B:135:ARG:O	1:B:139:GLU:HG3	2.20	0.42
1:D:25:THR:O	1:D:25:THR:HG23	2.21	0.41
1:D:6:CYS:HA	1:D:7:PRO:HD3	1.86	0.41
1:A:126:PRO:HD2	1:A:129:TRP:HB2	2.03	0.41
1:A:25:THR:HA	1:A:26:PRO:HD3	1.87	0.40
1:B:130:ARG:HA	1:B:131:PRO:HD3	1.81	0.40
1:D:36:LEU:C	1:D:36:LEU:HD23	2.42	0.40
1:D:144:ALA:HB3	7:D:2084:HOH:O	2.21	0.40
1:D:74:ALA:HB3	1:D:75:PRO:HD3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	136/154 (88%)	135 (99%)	1 (1%)	0	100	100
1	B	126/154 (82%)	124 (98%)	2 (2%)	0	100	100
1	C	137/154 (89%)	135 (98%)	2 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	134/154 (87%)	132 (98%)	2 (2%)	0	100	100
All	All	533/616 (86%)	526 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/120 (90%)	107 (99%)	1 (1%)	84	72
1	B	100/120 (83%)	100 (100%)	0	100	100
1	C	111/120 (92%)	108 (97%)	3 (3%)	52	27
1	D	106/120 (88%)	106 (100%)	0	100	100
All	All	425/480 (88%)	421 (99%)	4 (1%)	88	72

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	140	ARG
1	C	27	GLU
1	C	110[A]	LEU
1	C	110[B]	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 6 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	UMP	A	1147	-	16,21,21	2.59	4 (25%)	21,31,31	1.68	4 (19%)
5	ACT	B	1146	-	0,3,3	0.00	-	0,3,3	0.00	-
3	UMP	C	1145	-	16,21,21	2.55	5 (31%)	21,31,31	1.72	4 (19%)
6	GOL	D	1147	-	5,5,5	0.28	0	5,5,5	0.42	0
5	ACT	D	1148	-	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	UMP	A	1147	-	-	0/6/22/22	0/2/2/2
5	ACT	B	1146	-	-	0/0/0/0	0/0/0/0
3	UMP	C	1145	-	-	0/6/22/22	0/2/2/2
6	GOL	D	1147	-	-	0/4/4/4	0/0/0/0
5	ACT	D	1148	-	-	0/0/0/0	0/0/0/0

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1145	UMP	O3'-C3'	-2.81	1.37	1.43
3	A	1147	UMP	O3'-C3'	-2.68	1.37	1.43
3	C	1145	UMP	C4-N3	2.05	1.39	1.36
3	A	1147	UMP	O4'-C1'	3.24	1.49	1.42
3	C	1145	UMP	O4'-C1'	3.53	1.50	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1147	UMP	O4-C4	4.91	1.36	1.24
3	C	1145	UMP	O4-C4	5.00	1.37	1.24
3	C	1145	UMP	C6-N1	6.54	1.44	1.35
3	A	1147	UMP	C6-N1	7.06	1.44	1.35

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1147	UMP	O3'-C3'-C4'	2.01	118.39	110.10
3	C	1145	UMP	C5-C4-N3	2.25	122.10	117.07
3	A	1147	UMP	O4'-C1'-N1	2.40	111.91	107.71
3	A	1147	UMP	O4'-C4'-C5'	2.86	119.53	109.29
3	C	1145	UMP	O4'-C1'-N1	3.00	112.96	107.71
3	C	1145	UMP	O4'-C4'-C5'	3.37	121.34	109.29
3	C	1145	UMP	C2'-C1'-N1	4.25	124.64	114.14
3	A	1147	UMP	C2'-C1'-N1	4.56	125.38	114.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	138/154 (89%)	0.13	2 (1%) 78 84	9, 18, 40, 49	0
1	B	130/154 (84%)	0.24	6 (4%) 36 42	8, 17, 42, 65	0
1	C	137/154 (88%)	0.19	3 (2%) 65 72	8, 20, 43, 57	0
1	D	130/154 (84%)	0.36	11 (8%) 13 16	8, 18, 50, 75	0
All	All	535/616 (86%)	0.23	22 (4%) 41 47	8, 18, 45, 75	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	22	GLY	5.5
1	D	6	CYS	3.6
1	B	130	ARG	3.6
1	D	127	GLU	3.5
1	C	6	CYS	3.5
1	B	124	LEU	3.4
1	B	143	HIS	3.2
1	D	129	TRP	3.1
1	D	130	ARG	2.9
1	D	113	ALA	2.8
1	A	145	PRO	2.7
1	D	126	PRO	2.6
1	D	140	ARG	2.6
1	C	67	ALA	2.6
1	D	135	ARG	2.5
1	A	19	ARG	2.3
1	D	131	PRO	2.3
1	B	129	TRP	2.2
1	B	127	GLU	2.1
1	D	134	VAL	2.0
1	B	144	ALA	2.0
1	D	144	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	ACT	D	1148	4/4	0.74	0.20	13.88	30,36,40,44	0
5	ACT	B	1146	4/4	0.68	0.22	5.53	33,41,42,45	0
6	GOL	D	1147	6/6	0.91	0.12	1.31	17,29,34,34	0
2	MG	D	1145	1/1	0.98	0.11	0.80	28,28,28,28	0
4	CL	D	1146	1/1	0.99	0.10	-0.32	30,30,30,30	0
3	UMP	C	1145	20/20	0.98	0.10	-0.44	10,11,16,18	0
3	UMP	A	1147	20/20	0.98	0.10	-0.58	9,12,19,19	0
4	CL	A	1148	1/1	0.89	0.07	-1.78	48,48,48,48	0
2	MG	C	1144	1/1	0.98	0.11	-	21,21,21,21	0
2	MG	B	1145	1/1	0.95	0.09	-	29,29,29,29	0
2	MG	A	1146	1/1	0.95	0.10	-	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.