wwPDB NMR Structure Validation Summary Report (i)

Apr 27, 2016 – 05:42 AM BST

PDB ID	:	2YUE
Title	:	Solution structure of the NEUZ (NHR) domain in Neuralized from Drosophila melanogaster
Authors	:	He, F.; Muto, Y.; Inoue, M.; Kigawa, T.; Shirouzu, M.; Tarada, T.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on	:	2007-04-06

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 90%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	168	55% 30% • • 13%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 11 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:9-A:30, A:37-A:59, A:66-A:166 (146)	0.32	11

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 6, 7, 9, 10, 11, 12, 13, 15, 16, 19, 20
2	8, 14, 17
Single-model clusters	5; 18

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2620 atoms, of which 1286 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Protein neuralized.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	168	2620	845	1286	233	250	6	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	EXPRESSION TAG	UNP P29503
A	2	SER	-	EXPRESSION TAG	UNP P29503
A	3	SER	-	EXPRESSION TAG	UNP P29503
A	4	GLY	-	EXPRESSION TAG	UNP P29503
A	5	SER	-	EXPRESSION TAG	UNP P29503
A	6	SER	-	EXPRESSION TAG	UNP P29503
A	7	GLY	-	EXPRESSION TAG	UNP P29503

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Protein neuralized

Chain A: 55% 30% •• 13%

G1 S2 S3 G4 S5 S6 G7 P8 S13 V14 H15 R20 I21 S22 R23 D24 G25 T26 L27 A28 R29 R30 F31 E32 S33 F34 C35 R36 A37 I38 T39 F40 S41 V45 R46 I47 R50 I51 K54 F55 A56 E57 I58 S59 M60 M61 W62 M63 G64 G65 I66 R67 F68 D74

P75 Y85 A86 C87 P88 D89 L90 T91 A98 K99 A100 L101 E107 V116 V122 I123 I126 E130 L135 T136 D139 T140 L144 V147 I150 C154 T155 L160 R163 Y167 Q168

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 11. Colouring as in section 4.1 above.

- Molecule 1: Protein neuralized

Chain A: 53% 32% •• 13%

G1 S2 S3 G4 S5 S6 G7 P8 S13 V14 H15 R20 I21 S22 R23 D24 G25 T26 L27 A28 R29 R30 F31 E32 S33 F34 C35 R36 A37 I38 T39 F40 S41 V45 R46 I47 R50 I51 K54 E57 I58 S59 M60 M61 W62 M63 G64 G65 I66 R67 F68 G69 F70 T71

D74 P75 L78 Y85 A86 C87 P88 D89 L90 T91 W97 A98 K99 A100 L101 V116 V122 I123 E130 K131 G132 L135 T136 D139 T140 L144 W145 T146 D149 I150 C154 T155 L160 R163 Y167 Q168

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations, target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.1
CYANA	refinement	2.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 11351
Number of chemical shift lists	1
Total number of shifts	2050
Number of shifts mapped to atoms	2050
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	90%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1170	1152	1152	22±4
All	All	23400	23040	23040	449

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

5 of 115 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:45:VAL:HG21	1:A:51:ILE:HD11	0.78	1.53	4	14
1:A:66:ILE:HD11	1:A:101:LEU:HD13	0.74	1.58	9	5
1:A:51:ILE:HD13	1:A:146:THR:HG21	0.71	1.63	16	7
1:A:74:ASP:OD1	1:A:75:PRO:HD2	0.68	1.89	19	17
1:A:90:LEU:HD22	1:A:96:PHE:CE1	0.67	2.25	3	4

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	146/168 (87%)	124±3 (85±2%)	17±3 (12±2%)	5±2 (4±1%)	8 37
All	All	2920/3360 (87%)	2477 (85%)	340 (12%)	103 (4%)	8 37

5 of 12 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	91	THR	20
1	A	89	ASP	19
1	A	26	THR	15
1	A	25	GLY	15
1	A	151	TYR	7

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	126/143 (88%)	103±3 (82±2%)	23±3 (18±2%)	5 39
All	All	2520/2860 (88%)	2056 (82%)	464 (18%)	5 39

5 of 49 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	85	TYR	20
1	A	99	LYS	20
1	A	23	ARG	20
1	A	136	THR	20
1	A	91	THR	20

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 90% for the well-defined parts and 89% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 11351

Chemical shift list name: *assigned_ chem_ shift_ list_ 1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	2050
Number of shifts mapped to atoms	2050
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	7

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	161	0.15 ± 0.17	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	146	-0.04 ± 0.16	None needed (< 0.5 ppm)
$^{13}C'$	154	0.33 ± 0.14	None needed (< 0.5 ppm)
^{15}N	153	0.66 ± 0.38	None needed (imprecise)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 90%, i.e. 1643 atoms were assigned a chemical shift out of a possible 1823. 0 out of 19 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	703/720 (98%)	282/287 (98%)	283/292 (97%)	138/141 (98%)
Sidechain	777/931 (83%)	476/543 (88%)	288/339 (85%)	13/49 (27%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	163/172 (95%)	85/91 (93%)	76/76 (100%)	2/5 (40%)
Overall	1643/1823 (90%)	843/921 (92%)	647/707 (92%)	153/195 (78%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	74	ASP	HA	1.69	6.15 – 3.05	-9.4
1	A	75	PRO	HB3	-0.46	3.81 – 0.21	-6.9
1	A	28	ALA	HB1	-0.12	2.61 – 0.11	-5.9
1	A	28	ALA	HB3	-0.12	2.61 – 0.11	-5.9
1	A	28	ALA	HB2	-0.12	2.61 – 0.11	-5.9
1	A	131	LYS	HB3	0.21	3.10 – 0.40	-5.7
1	A	139	ASP	H	5.17	11.17 – 5.47	-5.5

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: