



Full wwPDB X-ray Structure Validation Report i

Jun 20, 2016 – 11:35 AM EDT

PDB ID : 4YXV
Title : PksG, a HMG-CoA Synthase from Bacillus subtilus
Authors : Till, M.; Nair, A.V.; Robson, A.; Race, P.R.
Deposited on : 2015-03-23
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027790
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027790


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3340 (2.80-2.72)
Clashscore	102246	3829 (2.80-2.72)
Ramachandran outliers	100387	3767 (2.80-2.72)
Sidechain outliers	100360	3770 (2.80-2.72)
RSRZ outliers	91569	3352 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 6264 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyketide biosynthesis 3-hydroxy-3-methylglutaryl-ACP synthase PksG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	398	3112	1982	525	582	23	0	0	0
1	B	401	3111	1979	521	587	24	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	114	ALA	CYS	engineered mutation	UNP P40830
B	114	ALA	CYS	engineered mutation	UNP P40830


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	22	Total O 22 22	0	0
2	B	19	Total O 19 19	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Polyketide biosynthesis 3-hydroxy-3-methylglutaryl-ACP synthase PksG



- Molecule 1: Polyketide biosynthesis 3-hydroxy-3-methylglutaryl-ACP synthase PksG



4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	74.67 Å 123.63 Å 199.35 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.00 – 2.75 63.92 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.9 (64.00-2.75) 99.9 (63.92-2.75)	Depositor EDS
R_{merge}	0.47	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	1.97 (at 2.77 Å)	Xtriage
Refinement program	REFMAC	Depositor
R , R_{free}	0.191 , 0.241 0.194 , 0.245	Depositor DCC
R_{free} test set	1243 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	48.3	Xtriage
Anisotropy	0.054	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 37.2	EDS
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.028 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.042 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6264	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	0/3177	0.78	3/4283 (0.1%)
1	B	0.63	0/3175	0.79	2/4283 (0.0%)
All	All	0.63	0/6352	0.78	5/8566 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	222	ASP	CB-CG-OD2	5.83	123.55	118.30
1	A	222	ASP	CB-CG-OD2	5.76	123.48	118.30
1	B	106	ARG	NE-CZ-NH1	5.49	123.05	120.30
1	A	378	ARG	N-CA-C	-5.29	96.71	111.00
1	A	106	ARG	NE-CZ-NH1	5.02	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3112	0	3050	38	0
1	B	3111	0	3036	34	0
2	A	22	0	0	1	0
2	B	19	0	0	0	0
All	All	6264	0	6086	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:183:VAL:O	1:B:339:THR:HB	1.86	0.75
1:A:183:VAL:O	1:A:339:THR:HB	1.87	0.75
1:A:278:THR:O	1:A:282:THR:HG23	1.87	0.74
1:B:278:THR:O	1:B:282:THR:HG23	1.86	0.74
1:B:13:GLY:H	1:B:355:HIS:HE1	1.35	0.73
1:A:13:GLY:H	1:A:355:HIS:HE1	1.37	0.71
1:A:251:THR:HB	1:A:257:VAL:HG21	1.72	0.70
1:B:251:THR:HB	1:B:257:VAL:HG21	1.74	0.67
1:B:123:MET:HE3	1:B:123:MET:HA	1.77	0.67
1:A:123:MET:HA	1:A:123:MET:HE3	1.78	0.64
1:B:79:THR:HG23	1:B:92:SER:OG	2.01	0.61
1:B:36:ASN:OD1	1:B:36:ASN:N	2.33	0.60
1:A:79:THR:HG23	1:A:92:SER:OG	2.01	0.60
1:B:123:MET:HA	1:B:123:MET:CE	2.32	0.59
1:A:234:ARG:HH12	1:B:104:ASN:HD21	1.52	0.58
1:A:249:TYR:O	1:A:251:THR:HG22	2.04	0.57
1:A:233:LYS:NZ	2:A:502:HOH:O	2.31	0.57
1:B:249:TYR:O	1:B:251:THR:HG22	2.04	0.56
1:A:378:ARG:HG2	1:A:379:PHE:N	2.21	0.55
1:A:206:PRO:HA	1:B:87:PHE:CE2	2.42	0.55
1:A:123:MET:HA	1:A:123:MET:CE	2.37	0.55
1:A:50:ASP:H	1:A:53:THR:HB	1.76	0.50
1:A:340:THR:HB	1:A:341:PRO:HD2	1.93	0.50
1:B:249:TYR:O	1:B:251:THR:CG2	2.60	0.50
1:A:20:VAL:HG13	1:A:166:PRO:HB2	1.93	0.50
1:A:90:SER:HB2	1:A:109:GLU:HB2	1.93	0.50
1:A:50:ASP:HB2	1:A:51:PRO:CD	2.42	0.50
1:A:87:PHE:CE2	1:B:206:PRO:HA	2.47	0.49
1:B:165:GLU:N	1:B:166:PRO:CD	2.76	0.49
1:A:165:GLU:N	1:A:166:PRO:CD	2.76	0.49
1:A:104:ASN:HD21	1:B:234:ARG:HH12	1.59	0.49
1:B:50:ASP:H	1:B:53:THR:HB	1.78	0.48
1:B:90:SER:HB2	1:B:109:GLU:HB2	1.94	0.48
1:B:50:ASP:HB2	1:B:51:PRO:CD	2.44	0.48
1:A:249:TYR:O	1:A:251:THR:CG2	2.61	0.47
1:B:340:THR:HB	1:B:341:PRO:HD2	1.95	0.47
1:B:79:THR:HG21	1:B:95:ILE:HD12	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:26:TYR:CG	1:B:364:MET:HG3	2.51	0.45
1:B:383:ASN:HA	1:B:409:ILE:O	2.16	0.45
1:A:378:ARG:HG2	1:A:379:PHE:H	1.81	0.45
1:A:383:ASN:HA	1:A:409:ILE:O	2.16	0.45
1:A:123:MET:HE2	1:B:123:MET:HB3	1.99	0.45
1:A:79:THR:HG21	1:A:95:ILE:HD12	1.98	0.45
1:B:190:ALA:HB1	1:B:231:TYR:CE1	2.52	0.45
1:A:373:GLY:O	1:A:376:MET:HB2	2.16	0.44
1:A:401:LYS:HB3	1:A:402:PRO:HD2	1.98	0.44
1:B:20:VAL:HG13	1:B:166:PRO:HB2	1.98	0.44
1:B:50:ASP:OD2	1:B:414:ARG:NH1	2.48	0.43
1:A:107:LEU:O	1:B:112:GLN:HG2	2.18	0.43
1:A:314:ARG:HD3	1:A:316:ASP:OD1	2.19	0.43
1:A:190:ALA:HB1	1:A:231:TYR:CE1	2.53	0.43
1:B:285:LYS:N	1:B:286:PRO:CD	2.81	0.43
1:B:373:GLY:O	1:B:376:MET:HB2	2.19	0.43
1:A:27:ARG:HB3	1:A:29:LEU:HD13	2.01	0.43
1:B:137:LYS:HD3	1:B:176:LEU:HD11	2.00	0.42
1:B:263:THR:O	1:B:267:LYS:HB2	2.19	0.42
1:A:314:ARG:O	1:A:319:LYS:NZ	2.51	0.42
1:A:50:ASP:OD2	1:A:414:ARG:NH1	2.50	0.42
1:B:401:LYS:HB3	1:B:402:PRO:HD2	2.00	0.42
1:A:73:ARG:NH1	1:A:75:GLU:OE2	2.46	0.42
1:A:358:ARG:HH11	1:A:358:ARG:HG2	1.86	0.41
1:A:358:ARG:CG	1:A:358:ARG:HH11	2.32	0.41
1:B:32:ALA:O	1:B:34:PHE:N	2.54	0.41
1:A:112:GLN:HG2	1:B:107:LEU:O	2.21	0.41
1:A:378:ARG:CG	1:A:379:PHE:N	2.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	390/420 (93%)	367 (94%)	20 (5%)	3 (1%)	24 55
1	B	395/420 (94%)	367 (93%)	24 (6%)	4 (1%)	19 48
All	All	785/840 (94%)	734 (94%)	44 (6%)	7 (1%)	21 52

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	112	GLN
1	B	34	PHE
1	B	112	GLN
1	B	397	SER
1	A	204	PRO
1	B	204	PRO
1	A	200	ASP

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	327/347 (94%)	302 (92%)	25 (8%)	16 39
1	B	326/347 (94%)	297 (91%)	29 (9%)	12 31
All	All	653/694 (94%)	599 (92%)	54 (8%)	14 35

All (54) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	LEU
1	A	53	THR
1	A	79	THR
1	A	107	LEU
1	A	112	GLN
1	A	183	VAL
1	A	193	TYR
1	A	198	VAL
1	A	205	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	215	LEU
1	A	216	SER
1	A	217	LEU
1	A	233	LYS
1	A	241	GLN
1	A	251	THR
1	A	266	ARG
1	A	267	LYS
1	A	282	THR
1	A	339	THR
1	A	342	GLN
1	A	346	ARG
1	A	354	LYS
1	A	369	LEU
1	A	385	LYS
1	A	401	LYS
1	B	30	ASP
1	B	36	ASN
1	B	46	LEU
1	B	53	THR
1	B	79	THR
1	B	107	LEU
1	B	112	GLN
1	B	162	SER
1	B	163	TYR
1	B	183	VAL
1	B	198	VAL
1	B	205	ILE
1	B	215	LEU
1	B	216	SER
1	B	217	LEU
1	B	233	LYS
1	B	241	GLN
1	B	251	THR
1	B	266	ARG
1	B	267	LYS
1	B	282	THR
1	B	317	THR
1	B	339	THR
1	B	342	GLN
1	B	346	ARG
1	B	354	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	364	MET
1	B	369	LEU
1	B	396	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	104	ASN
1	A	112	GLN
1	A	126	ASN
1	A	131	GLN
1	A	185	GLN
1	A	191	ASN
1	A	241	GLN
1	A	342	GLN
1	A	344	GLN
1	A	355	HIS
1	B	104	ASN
1	B	112	GLN
1	B	126	ASN
1	B	131	GLN
1	B	185	GLN
1	B	191	ASN
1	B	241	GLN
1	B	342	GLN
1	B	344	GLN
1	B	355	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	398/420 (94%)	-0.20	3 (0%)	87	83	26, 42, 90, 127
1	B	401/420 (95%)	-0.16	3 (0%)	89	86	27, 45, 95, 118
All	All	799/840 (95%)	-0.18	6 (0%)	87	83	26, 43, 93, 127

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	206	PRO	3.5
1	B	28	HIS	3.5
1	A	420	SER	3.1
1	B	202	CYS	2.1
1	A	205	ILE	2.1
1	A	202	CYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates i

There are no carbohydrates in this entry.

6.4 Ligands i

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.