



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2016 – 08:47 PM GMT

PDB ID : 4Z2T
Title : Crystal Structure of 2-hydroxybiphenyl 3-monooxygenase W225Y from *Pseudomonas azelaica*
Authors : Kanteev, M.; Bregman-Cohen, A.; Fishman, A.
Deposited on : 2015-03-30
Resolution : 2.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4776 (2.50-2.42)
Clashscore	102246	1030 (2.48-2.44)
Ramachandran outliers	100387	1024 (2.48-2.44)
Sidechain outliers	100360	1024 (2.48-2.44)
RSRZ outliers	91569	4787 (2.50-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	592	
1	B	592	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9201 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-hydroxybiphenyl-3-monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	559	Total	C	N	O	S	0	0	0
			4301	2717	756	808	20			
1	B	564	Total	C	N	O	S	0	0	0
			4339	2741	761	817	20			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	MET	-	initiating methionine	UNP O06647
A	-4	HIS	-	expression tag	UNP O06647
A	-3	HIS	-	expression tag	UNP O06647
A	-2	HIS	-	expression tag	UNP O06647
A	-1	HIS	-	expression tag	UNP O06647
A	0	HIS	-	expression tag	UNP O06647
A	1	HIS	-	expression tag	UNP O06647
A	225	TYR	TRP	engineered mutation	UNP O06647
B	-5	MET	-	initiating methionine	UNP O06647
B	-4	HIS	-	expression tag	UNP O06647
B	-3	HIS	-	expression tag	UNP O06647
B	-2	HIS	-	expression tag	UNP O06647
B	-1	HIS	-	expression tag	UNP O06647
B	0	HIS	-	expression tag	UNP O06647
B	1	HIS	-	expression tag	UNP O06647
B	225	TYR	TRP	engineered mutation	UNP O06647


- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	242	Total	O	0	0
			242	242		
3	B	213	Total	O	0	0
			213	213		



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	154.38Å 130.58Å 78.75Å 90.00° 98.79° 90.00°	Depositor
Resolution (Å)	43.85 – 2.45 43.85 – 2.45	Depositor EDS
% Data completeness (in resolution range)	97.0 (43.85-2.45) 97.1 (43.85-2.45)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.60 (at 2.45Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.218 , 0.236 0.225 , 0.246	Depositor DCC
R_{free} test set	2792 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	27.5	Xtriage
Anisotropy	0.570	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	2 of 54992 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9201	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.97	7/4393 (0.2%)	0.81	9/5951 (0.2%)
1	B	0.88	4/4433 (0.1%)	0.81	13/6007 (0.2%)
All	All	0.93	11/8826 (0.1%)	0.81	22/11958 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	76	TYR	C-N	-17.31	0.94	1.34
1	A	416	ALA	C-N	-9.87	1.11	1.34
1	A	520	ILE	C-N	-9.69	1.11	1.34
1	B	360	ARG	C-N	-9.00	1.13	1.34
1	A	358	ALA	C-N	-7.71	1.16	1.34

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	77	MET	O-C-N	-11.92	102.94	123.20
1	B	359	GLU	C-N-CA	-10.12	96.40	121.70
1	A	77	MET	CA-C-N	9.72	135.64	116.20
1	B	360	ARG	O-C-N	-8.80	108.62	122.70
1	B	198	ILE	N-CA-C	-8.47	88.12	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	360	ARG	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4301	0	4251	149	0
1	B	4339	0	4286	149	0
2	A	53	0	31	5	0
2	B	53	0	31	9	0
3	A	242	0	0	2	1
3	B	213	0	0	4	2
All	All	9201	0	8599	299	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 299 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:70:LEU:O	1:A:245:ARG:NH2	1.76	1.16
1:A:223:MET:HE2	1:A:225:TYR:HE1	1.10	1.15
1:A:191:PRO:HG2	1:A:301:VAL:CG2	1.79	1.12
1:B:46:ARG:HH11	1:B:46:ARG:CG	1.68	1.06
1:A:270:ALA:HA	1:A:273:ILE:HD12	1.38	1.06

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:816:HOH:O	3:B:816:HOH:O[2_655]	1.33	0.87
3:A:807:HOH:O	3:B:881:HOH:O[4_555]	2.03	0.17

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	551/592 (93%)	527 (96%)	23 (4%)	1 (0%)	52	64
1	B	558/592 (94%)	533 (96%)	22 (4%)	3 (0%)	34	41
All	All	1109/1184 (94%)	1060 (96%)	45 (4%)	4 (0%)	39	49

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	198	ILE
1	B	391	PRO
1	B	270	ALA
1	A	442	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	448/473 (95%)	444 (99%)	4 (1%)	84	91
1	B	452/473 (96%)	445 (98%)	7 (2%)	72	84
All	All	900/946 (95%)	889 (99%)	11 (1%)	78	87

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	46	ARG
1	B	243	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	319	THR
1	A	370	ARG
1	B	277	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	147	HIS
1	B	48	HIS
1	B	297	GLN
1	B	513	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	601	-	52,58,58	1.29	6 (11%)	52,89,89	2.16	9 (17%)
2	FAD	B	601	-	52,58,58	1.30	6 (11%)	52,89,89	2.16	9 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	601	-	-	0/30/50/50	0/6/6/6
2	FAD	B	601	-	-	0/30/50/50	0/6/6/6

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	FAD	C9A-N10	2.47	1.42	1.38
2	B	601	FAD	C9A-N10	2.54	1.42	1.38
2	B	601	FAD	C5A-C4A	2.70	1.46	1.40
2	A	601	FAD	C5A-C4A	2.73	1.46	1.40
2	A	601	FAD	C8-C7	2.83	1.48	1.41

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	FAD	N3A-C2A-N1A	-7.97	122.61	128.87
2	B	601	FAD	N3A-C2A-N1A	-7.79	122.75	128.87
2	B	601	FAD	C4-C4X-C10	-4.80	116.87	119.94
2	A	601	FAD	C4-C4X-C10	-4.79	116.88	119.94
2	A	601	FAD	C4X-C4-N3	-3.76	118.61	123.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	FAD	5	0
2	B	601	FAD	9	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	4
1	B	2

The worst 5 of 6 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	70:LEU	C	71:ALA	N	1.18
1	A	358:ALA	C	359:GLU	N	1.16
1	B	360:ARG	C	361:SER	N	1.13
1	A	416:ALA	C	417:LEU	N	1.11
1	A	520:ILE	C	521:GLU	N	1.11

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	559/592 (94%)	0.27	39 (6%) 19 21	10, 24, 63, 131	0
1	B	564/592 (95%)	0.39	49 (8%) 13 13	11, 25, 73, 140	0
All	All	1123/1184 (94%)	0.33	88 (7%) 16 16	10, 25, 67, 140	0

The worst 5 of 88 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	198	ILE	8.1
1	B	197	GLY	7.9
1	B	196	MET	7.8
1	B	390	ALA	7.2
1	B	200	ASP	7.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FAD	B	601	53/53	0.93	0.17	0.19	15,31,43,46	0
2	FAD	A	601	53/53	0.95	0.15	-0.41	16,30,42,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.