



Full wwPDB X-ray Structure Validation Report ⓘ

May 4, 2016 – 05:09 PM EDT

PDB ID : 4ZBC
Title : A dehydrated form of glucose isomerase collected at 100K.
Authors : Sandy, J.
Deposited on : 2015-04-14
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027457
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> % 92% 6% • </div> </div>
1	B	388	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="display: flex; justify-content: space-between; margin-top: 5px;"> % 90% 8% • </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MN	A	403	-	-	-	X
2	MN	B	405	-	-	-	X
3	GLC	A	407	-	-	-	X
3	GLC	A	408	-	-	-	X
3	GLC	B	406	-	-	-	X
3	GLC	B	407	-	-	-	X
4	FRU	A	409	-	-	-	X
4	FRU	A	410	-	-	-	X
4	FRU	B	408	-	-	-	X
4	FRU	B	409	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6755 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Xylose isomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	387	Total	C	N	O	S	0	0	0
			3045	1913	550	574	8			
1	B	387	Total	C	N	O	S	0	2	0
			3061	1923	556	574	8			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	388	GLY	-	expression tag	UNP P24300
B	388	GLY	-	expression tag	UNP P24300

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	5	Total	Mn	0	0
			5	5		
2	A	6	Total	Mn	0	0
			6	6		

- Molecule 3 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			12	6	6		
3	A	1	Total	C	O	0	0
			12	6	6		
3	B	1	Total	C	O	0	0
			12	6	6		
3	B	1	Total	C	O	0	0
			12	6	6		

- Molecule 4 is FRUCTOSE (three-letter code: FRU) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			12	6	6		
4	A	1	Total	C	O	0	0
			12	6	6		
4	B	1	Total	C	O	0	0
			12	6	6		
4	B	1	Total	C	O	0	0
			12	6	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	282	Total	O	0	0
			282	282		
5	B	260	Total	O	0	0
			260	260		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Xylose isomerase



- Molecule 1: Xylose isomerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	81.63Å 93.64Å 97.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	67.58 – 2.00 67.58 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.3 (67.58-2.00) 99.3 (67.58-2.00)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.40 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.136 , 0.182 0.150 , 0.193	Depositor DCC
R_{free} test set	2431 reflections (5.02%)	DCC
Wilson B-factor (Å ²)	13.6	Xtriage
Anisotropy	0.612	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 50.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.015 for -h,l,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6755	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, FRU, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.95	3/3117 (0.1%)	1.10	18/4219 (0.4%)
1	B	0.92	1/3139 (0.0%)	1.07	21/4248 (0.5%)
All	All	0.93	4/6256 (0.1%)	1.08	39/8467 (0.5%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	328	GLU	CG-CD	7.13	1.62	1.51
1	A	355	SER	CB-OG	-6.23	1.34	1.42
1	B	160	GLU	CD-OE1	5.44	1.31	1.25
1	A	331	ARG	CZ-NH1	5.13	1.39	1.33

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	331	ARG	NE-CZ-NH2	-14.71	112.94	120.30
1	B	266	ARG	NE-CZ-NH2	-13.71	113.44	120.30
1	A	266	ARG	NE-CZ-NH2	-11.94	114.33	120.30
1	B	74	ARG	NE-CZ-NH2	-11.05	114.77	120.30
1	A	331	ARG	CG-CD-NE	-9.56	91.73	111.80
1	A	381	ASP	CB-CG-OD1	8.75	126.17	118.30
1	A	266	ARG	NE-CZ-NH1	8.54	124.57	120.30
1	A	387	ARG	NE-CZ-NH2	-8.35	116.12	120.30
1	A	387	ARG	NE-CZ-NH1	8.16	124.38	120.30
1	A	55	ASP	CB-CG-OD1	8.16	125.64	118.30
1	B	316	ARG	NE-CZ-NH1	-7.96	116.32	120.30
1	B	259	ARG	NE-CZ-NH2	-7.45	116.57	120.30
1	B	76	ARG	NE-CZ-NH2	-7.23	116.68	120.30
1	B	35	ASP	CB-CG-OD1	7.23	124.80	118.30
1	A	74	ARG	NE-CZ-NH2	-7.09	116.75	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	55	ASP	CB-CG-OD2	-7.01	111.99	118.30
1	B	109	ARG	NE-CZ-NH1	6.70	123.65	120.30
1	A	368	ARG	NE-CZ-NH2	-6.42	117.09	120.30
1	A	188	ARG	NE-CZ-NH1	6.37	123.48	120.30
1	B	266	ARG	NE-CZ-NH1	6.16	123.38	120.30
1	B	368	ARG	NE-CZ-NH1	6.10	123.35	120.30
1	A	368	ARG	NE-CZ-NH1	5.92	123.26	120.30
1	A	381	ASP	CB-CG-OD2	-5.89	113.00	118.30
1	B	336	ASP	CB-CG-OD1	5.83	123.55	118.30
1	B	74	ARG	NE-CZ-NH1	5.78	123.19	120.30
1	B	42	ARG	NE-CZ-NH1	5.74	123.17	120.30
1	B	24	ASP	CB-CG-OD1	5.74	123.47	118.30
1	B	35	ASP	CB-CG-OD2	-5.71	113.16	118.30
1	B	323	ASP	CB-CG-OD1	5.57	123.31	118.30
1	B	109	ARG	NE-CZ-NH2	-5.46	117.57	120.30
1	B	316	ARG	NE-CZ-NH2	5.45	123.03	120.30
1	A	337	GLU	OE1-CD-OE2	-5.41	116.81	123.30
1	B	76	ARG	NE-CZ-NH1	5.27	122.94	120.30
1	B	160	GLU	OE1-CD-OE2	5.24	129.58	123.30
1	A	42	ARG	NE-CZ-NH1	5.23	122.91	120.30
1	A	336	ASP	CB-CG-OD1	5.18	122.96	118.30
1	B	80	ASP	CB-CG-OD1	5.12	122.91	118.30
1	A	245	ASP	CB-CG-OD1	5.09	122.89	118.30
1	B	361	ASP	CB-CG-OD2	-5.01	113.79	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3045	0	2914	16	0
1	B	3061	0	2940	21	3
2	A	6	0	0	0	0
2	B	5	0	0	0	0
3	A	24	0	21	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	24	0	18	5	0
4	A	24	0	24	0	0
4	B	24	0	24	3	2
5	A	282	0	0	7	2
5	B	260	0	0	10	0
All	All	6755	0	5941	41	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:406:GLC:C6	3:B:407:GLC:O6	1.66	1.39
1:B:80:ASP:CG	5:B:502:HOH:O	1.63	1.35
1:A:331:ARG:HD2	5:A:503:HOH:O	1.25	1.31
4:B:408:FRU:H5	5:B:504:HOH:O	1.01	1.18
1:A:80:ASP:HB3	5:A:631:HOH:O	1.42	1.17
1:A:327:GLN:HG2	5:A:762:HOH:O	1.68	0.92
1:A:340:ARG:NH2	5:A:502:HOH:O	2.04	0.89
4:B:408:FRU:C5	5:B:504:HOH:O	1.79	0.75
1:B:80:ASP:CB	5:B:502:HOH:O	2.25	0.71
1:B:205[B]:ARG:NH1	5:B:505:HOH:O	2.24	0.70
1:B:354:ARG:NE	1:B:359:GLU:OE2	2.23	0.68
1:B:80:ASP:OD2	5:B:502:HOH:O	1.93	0.65
3:B:406:GLC:C6	3:B:407:GLC:C6	2.74	0.65
1:B:85:LYS:HE2	1:B:85:LYS:HA	1.80	0.64
1:B:76:ARG:NH1	5:B:506:HOH:O	2.31	0.63
3:B:406:GLC:O6	3:B:407:GLC:O5	2.17	0.61
1:B:382:HIS:CE1	5:B:526:HOH:O	2.54	0.60
1:B:382:HIS:HE1	5:B:526:HOH:O	1.86	0.58
1:B:85:LYS:CA	1:B:85:LYS:HE2	2.37	0.54
1:B:201:ALA:HB1	1:B:205[B]:ARG:HH22	1.73	0.52
1:A:152:ARG:CZ	3:A:408:GLC:C6	2.92	0.47
1:B:245:ASP:OD1	1:B:285:HIS:HD2	1.96	0.46
1:A:340:ARG:HD2	1:B:109:ARG:HD2	1.96	0.46
1:B:181:GLU:OE1	3:B:406:GLC:O2	2.33	0.46
1:A:52:THR:OG1	1:A:285:HIS:HE1	1.98	0.46
1:A:24:ASP:HB2	1:A:25:PRO:CD	2.46	0.46
1:A:97:PRO:HB2	1:B:366:ALA:HB1	1.98	0.45
1:A:245:ASP:OD1	1:A:285:HIS:HD2	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:20:TRP:CZ3	4:B:408:FRU:H61	2.52	0.44
1:A:76:ARG:NH2	1:A:128:GLU:HG2	2.31	0.44
5:A:674:HOH:O	1:B:205[A]:ARG:HD2	2.17	0.44
1:A:80:ASP:CB	5:A:631:HOH:O	2.24	0.43
1:A:374:ARG:HG3	5:A:647:HOH:O	2.18	0.43
1:B:85:LYS:CE	1:B:85:LYS:HA	2.47	0.42
1:A:152:ARG:CZ	3:A:408:GLC:H62	2.49	0.42
1:B:328:GLU:OE2	5:B:503:HOH:O	2.22	0.42
1:A:52:THR:OG1	1:A:285:HIS:CE1	2.73	0.41
1:B:52:THR:OG1	1:B:285:HIS:HE1	2.03	0.41
1:B:24:ASP:HB2	1:B:25:PRO:CD	2.51	0.41
1:A:387:ARG:HD2	1:A:387:ARG:HA	1.96	0.40
3:B:406:GLC:O6	3:B:407:GLC:C6	2.70	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:651:HOH:O	5:A:733:HOH:O[2_455]	1.51	0.69
5:A:663:HOH:O	5:A:663:HOH:O[2_455]	1.73	0.47
1:B:358:GLU:OE1	4:B:409:FRU:O1[1_655]	1.96	0.24
1:B:351:LEU:O	4:B:409:FRU:O2[1_655]	1.98	0.22
1:B:316:ARG:NH1	1:B:384:LEU:O[2_455]	2.01	0.19

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/388 (99%)	371 (96%)	13 (3%)	1 (0%)	46	41
1	B	387/388 (100%)	374 (97%)	11 (3%)	2 (0%)	34	26
All	All	772/776 (100%)	745 (96%)	24 (3%)	3 (0%)	39	33

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	GLU
1	B	186	GLU
1	B	357	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/304 (100%)	299 (99%)	4 (1%)	76	79
1	B	305/304 (100%)	302 (99%)	3 (1%)	82	85
All	All	608/608 (100%)	601 (99%)	7 (1%)	78	81

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	ARG
1	A	76	ARG
1	A	176	ILE
1	A	325	GLU
1	B	2	ASN
1	B	8	GLU
1	B	42	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	285	HIS
1	B	2	ASN
1	B	285	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 19 ligands modelled in this entry, 11 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	GLC	A	407	2	12,12,12	1.11	1 (8%)	17,17,17	2.77	6 (35%)
3	GLC	A	408	-	12,12,12	0.86	0	17,17,17	1.98	7 (41%)
4	FRU	A	409	-	11,12,12	1.19	0	10,18,18	0.71	0
4	FRU	A	410	-	11,12,12	1.56	2 (18%)	10,18,18	2.02	3 (30%)
3	GLC	B	406	2	12,12,12	1.97	3 (25%)	17,17,17	4.56	11 (64%)
3	GLC	B	407	-	12,12,12	1.24	1 (8%)	17,17,17	2.66	9 (52%)
4	FRU	B	408	-	11,12,12	1.03	1 (9%)	10,18,18	2.05	2 (20%)
4	FRU	B	409	-	11,12,12	0.74	0	10,18,18	1.24	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	A	407	2	-	0/2/22/22	0/1/1/1
3	GLC	A	408	-	-	0/2/22/22	0/1/1/1
4	FRU	A	409	-	-	0/5/24/24	0/1/1/1
4	FRU	A	410	-	-	0/5/24/24	0/1/1/1
3	GLC	B	406	2	-	0/2/22/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	B	407	-	-	0/2/22/22	0/1/1/1
4	FRU	B	408	-	-	0/5/24/24	0/1/1/1
4	FRU	B	409	-	-	0/5/24/24	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	408	FRU	C1-C2	2.13	1.55	1.52
3	B	407	GLC	O5-C1	2.16	1.47	1.43
3	B	406	GLC	C1-C2	2.47	1.58	1.52
4	A	410	FRU	O3-C3	2.48	1.47	1.42
3	A	407	GLC	O5-C1	2.74	1.48	1.43
4	A	410	FRU	O5-C2	3.34	1.48	1.43
3	B	406	GLC	O5-C5	3.62	1.53	1.44
3	B	406	GLC	O1-C1	4.13	1.54	1.39

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	406	GLC	C6-C5-C4	-8.24	92.33	112.99
4	B	408	FRU	O2-C2-O5	-5.67	97.60	109.42
3	B	406	GLC	O6-C6-C5	-5.34	93.48	111.30
3	B	407	GLC	C4-C3-C2	-4.21	103.03	110.79
4	A	410	FRU	O3-C3-C4	-3.73	100.18	113.43
3	B	406	GLC	O5-C1-C2	-3.53	103.83	110.00
4	A	410	FRU	O2-C2-O5	-3.48	102.17	109.42
3	B	406	GLC	C4-C3-C2	-3.42	104.49	110.79
3	A	407	GLC	O4-C4-C3	-2.51	104.70	110.36
3	B	407	GLC	O1-C1-C2	-2.45	102.28	109.05
3	A	408	GLC	O5-C5-C6	-2.28	100.47	106.38
3	A	408	GLC	O4-C4-C3	-2.25	105.28	110.36
3	A	408	GLC	C1-O5-C5	-2.10	109.53	113.54
3	B	407	GLC	O5-C5-C4	-2.08	105.69	109.67
3	A	408	GLC	O2-C2-C1	2.08	114.29	109.74
4	B	408	FRU	O1-C1-C2	2.21	115.90	111.54
3	B	407	GLC	O5-C5-C6	2.25	112.21	106.38
3	B	407	GLC	O4-C4-C5	2.27	115.20	109.23
3	A	408	GLC	O6-C6-C5	2.40	119.31	111.30
3	B	407	GLC	O3-C3-C4	2.63	116.29	110.36
4	B	409	FRU	O4-C4-C3	2.68	120.71	112.09
4	A	410	FRU	O4-C4-C5	2.70	119.06	111.01
3	B	406	GLC	O2-C2-C1	3.30	116.96	109.74

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	406	GLC	C1-O5-C5	3.40	120.05	113.54
3	A	408	GLC	O1-C1-C2	3.46	118.62	109.05
3	B	406	GLC	O5-C5-C4	3.57	116.49	109.67
3	B	407	GLC	O6-C6-C5	3.70	123.67	111.30
3	B	407	GLC	O5-C1-C2	3.89	116.81	110.00
3	A	407	GLC	C4-C3-C2	3.94	118.04	110.79
3	B	406	GLC	O1-C1-C2	3.95	119.97	109.05
3	A	407	GLC	O2-C2-C1	4.33	119.22	109.74
3	A	408	GLC	O4-C4-C5	4.47	121.00	109.23
3	A	407	GLC	O4-C4-C5	4.66	121.49	109.23
3	A	407	GLC	O2-C2-C3	4.98	121.60	110.36
3	A	407	GLC	O3-C3-C2	5.57	122.92	110.36
3	B	407	GLC	C1-O5-C5	6.31	125.61	113.54
3	B	406	GLC	C1-C2-C3	6.34	121.07	110.68
3	B	406	GLC	O5-C5-C6	7.56	125.96	106.38
3	B	406	GLC	O4-C4-C3	8.79	130.18	110.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	408	GLC	2	0
3	B	406	GLC	5	0
3	B	407	GLC	4	0
4	B	408	FRU	3	0
4	B	409	FRU	0	2

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	387/388 (99%)	-0.64	2 (0%) 91 92	8, 12, 26, 55	0
1	B	387/388 (99%)	-0.63	2 (0%) 91 92	8, 12, 26, 55	0
All	All	774/776 (99%)	-0.63	4 (0%) 91 92	8, 12, 26, 55	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	ASN	4.7
1	B	2	ASN	2.8
1	B	388	GLY	2.5
1	A	175	ASP	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	MN	B	405	1/1	0.96	0.23	19.30	58,58,58,58	0
2	MN	A	403	1/1	1.00	0.19	11.26	47,47,47,47	0
4	FRU	A	410	12/12	0.70	0.30	7.92	35,40,43,43	0
3	GLC	B	406	12/12	0.90	0.17	7.53	14,17,21,30	0
3	GLC	A	408	12/12	0.88	0.18	6.42	18,30,33,38	0
3	GLC	B	407	12/12	0.74	0.27	5.36	25,35,42,49	0
4	FRU	A	409	12/12	0.92	0.16	4.65	21,24,25,27	0
4	FRU	B	408	12/12	0.92	0.22	4.10	20,25,30,44	0
4	FRU	B	409	12/12	0.81	0.23	3.91	33,50,57,62	0
3	GLC	A	407	12/12	0.94	0.12	3.57	14,16,17,19	0
2	MN	B	402	1/1	1.00	0.07	0.36	12,12,12,12	0
2	MN	B	403	1/1	1.00	0.05	-1.56	25,25,25,25	0
2	MN	A	402	1/1	1.00	0.05	-3.21	12,12,12,12	0
2	MN	B	401	1/1	1.00	0.06	-	11,11,11,11	0
2	MN	A	405	1/1	0.98	0.20	-	46,46,46,46	0
2	MN	A	406	1/1	0.98	0.07	-	32,32,32,32	0
2	MN	A	401	1/1	0.99	0.05	-	12,12,12,12	0
2	MN	B	404	1/1	0.97	0.16	-	44,44,44,44	0
2	MN	A	404	1/1	1.00	0.06	-	15,15,15,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.