Full wwPDB X-ray Structure Validation Report (i)

Feb 19, 2016 – 09:01 PM GMT

PDB ID	:	4ZFG
Title	:	Dual-specificity Fab 5A12 in complex with Angiopoietin 2
Authors	:	Harris, S.F.; Wu, P.
Deposited on	:	2015-04-21
Resolution	:	2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026982
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20026982

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	5193 (2.30-2.26)
Clashscore	102246	5929 (2.30-2.26)
Ramachandran outliers	100387	5851 (2.30-2.26)
Sidechain outliers	100360	5850 (2.30-2.26)
RSRZ outliers	91569	5204 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	232	2%, 89%, 6%, 5%
2	H	228	8%, 86%, 8%, 6%
3	L	215	5%, 85%, 12%, • •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	SO4	H	301	-	-	-	X

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 5445 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Angiopoietin-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	220	1781	1129	302	339	11	0	1	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	274	ALA	-	expression tag	UNP O15123
A	275	GLY	-	expression tag	UNP O15123
A	276	SER	-	expression tag	UNP O15123
A	497	GLY	-	expression tag	UNP O15123
A	498	ASN	-	expression tag	UNP O15123
A	499	SER	-	expression tag	UNP O15123
A	500	HIS	-	expression tag	UNP O15123
A	501	HIS	-	expression tag	UNP O15123
A	502	HIS	-	expression tag	UNP O15123
A	503	HIS	-	expression tag	UNP O15123
A	504	HIS	-	expression tag	UNP O15123
A	505	HIS	-	expression tag	UNP O15123

- Molecule 2 is a protein called Fragment antigen binding 5A12 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	215	1618	1034	265	313	6	0	1	0

- Molecule 3 is a protein called Fragment antigen binding 5A12 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	213	1628	1023	271	329	5	0	1	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
4	A	1	1	1	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
5	A	1	5	4	1	0	0
			Total	O	S		
5	A	1	5	4	1	0	0
			Total	O	S		
5	A	1	5	4	1	0	0
			Total	O	S		
5	H	1	5	4	1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	122	122	122	0	0
			Total	O		
6	H	145	145	145	0	0
			Total	O		
6	L	130	130	130	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Angiopoietin-2

Chain A:

- Molecule 2: Fragment antigen binding 5A12 heavy chain

Chain H:





- Molecule 3: Fragment antigen binding 5A12 light chain

Chain L:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	181.67Å 109.29Å 43.95Å 90.00° 100.47° 90.00°	Depositor
Resolution (Å)	43.21 – 2.27 43.21 – 2.27	Depositor EDS
% Data completeness (in resolution range)	99.5 (43.21-2.27) 99.3 (43.21-2.27)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.22 (at 2.27Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R, R_{free}	0.177 , 0.223 0.177 , 0.220	Depositor DCC
R_{free} test set	1947 reflections (5.28%)	DCC
Wilson B-factor (Å^2)	46.9	Xtriage
Anisotropy	0.722	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.35 , 62.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.34$	Xtriage
Outliers	0 of 38887 reflections	Xtriage
F_o,F_c correlation	0.97	EDS
Total number of atoms	5445	wwPDB-VP
Average B, all atoms (Å^2)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.50	0/1832	0.66	0/2474
2	H	0.51	0/1661	0.72	0/2269
3	L	0.51	1/1663 (0.1%)	0.73	0/2256
All	All	0.51	1/5156 (0.0%)	0.71	0/6999

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	L	4	MET	SD-CE	-5.72	1.45	1.77

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1781	0	1659	4	0
2	H	1618	0	1575	9	0
3	L	1628	0	1594	19	0
4	A	1	0	0	0	0
5	A	15	0	0	0	0
5	H	5	0	0	0	0
6	A	122	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	H	145	0	0	0	0
6	L	130	0	0	7	0
All	All	5445	0	4828	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:58:VAL:HG23	6:L:309:HOH:O	1.70	0.91
3:L:37:GLN:HB2	3:L:47:LEU:HD11	1.69	0.74
1:A:297:ILE:HG12	1:A:349:PRO:HB2	1.69	0.74
3:L:55:TYR:HD2	6:L:309:HOH:O	1.76	0.67
3:L:120:PRO:HD3	3:L:132:VAL:HG22	1.79	0.65
2:H:140:CYS:HG	2:H:196:CYS:HG	0.72	0.64
3:L:163:VAL:HG22	3:L:175:LEU:HD22	1.80	0.63
3:L:4:MET:HE3	3:L:28:LEU:HD11	1.81	0.61
3:L:163:VAL:HG22	3:L:175:LEU:CD2	2.33	0.59
2:H:96:VAL:O	2:H:100(A):TYR:HA	2.04	0.57
2:H:33:TRP:HB2	2:H:95:PHE:HB3	1.88	0.55
2:H:6:GLU:HA	2:H:21:SER:O	2.10	0.52
3:L:136:LEU:HB2	3:L:175:LEU:HB3	1.95	0.49
2:H:94:ARG:O	2:H:100(C):MET:HA	2.13	0.47
3:L:164:THR:CG2	6:L:362:HOH:O	2.61	0.47
3:L:23:CYS:CB	3:L:88:CYS:HG	2.21	0.46
3:L:20:THR:HG23	6:L:416:HOH:O	2.16	0.45
1:A:415:SER:HB2	1:A:474:TRP:CE2	2.52	0.45
3:L:15:VAL:HA	3:L:78:LEU:O	2.17	0.45
2:H:12:VAL:HG21	2:H:82(C):LEU:HD13	1.99	0.45
2:H:140:CYS:CB	2:H:196:CYS:HG	2.26	0.44
2:H:61:ASP:HA	2:H:64:LYS:HD2	1.99	0.44
3:L:164:THR:HG22	3:L:174:SER:H	1.83	0.44
3:L:164:THR:HG23	6:L:362:HOH:O	2.16	0.44
3:L:210:ASN:O	3:L:211:ARG:HB2	2.18	0.43
3:L:47:LEU:HA	3:L:58:VAL:HG21	2.00	0.42
2:H:126:PRO:HG3	2:H:138:LEU:HB3	2.01	0.42
1:A:424:SER:O	1:A:444:GLY:HA2	2.19	0.42
1:A:283:ASP:HA	1:A:300:LEU:HD13	2.01	0.42
3:L:21:ILE:HG12	3:L:102:THR:HG21	2.02	0.41
3:L:20:THR:HG21	6:L:354:HOH:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:100:GLN:HG2	6:L:337:HOH:O	2.19	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/232 (94%)	207 (94%)	11 (5%)	1 (0%)	34	39
2	H	212/228 (93%)	208 (98%)	4 (2%)	0	100	100
3	L	212/215 (99%)	203 (96%)	8 (4%)	1 (0%)	34	39
All	All	643/675 (95%)	618 (96%)	23 (4%)	2 (0%)	46	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	L	211	ARG
1	A	462	GLN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/197 (95%)	183 (97%)	5 (3%)	52	68
2	H	176/188 (94%)	171 (97%)	5 (3%)	51	66

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	L	186/187 (100%)	174 (94%)	12 (6%)	21	26
All	All	550/572 (96%)	528 (96%)	22 (4%)	38	50

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	366	GLN
1	A	368	ARG
1	A	371	LEU
1	A	397	LEU
1	A	412	LYS
2	H	12	VAL
2	H	30	SER
2	H	150	VAL
2	H	193	THR
2	H	197	ASN
3	L	1	ASP
3	L	27	GLN
3	L	73	LEU
3	L	97	THR
3	L	100	GLN
3	L	127	SER
3	L	129	THR
3	L	142	ARG
3	L	164	THR
3	L	194	CYS
3	L	210	ASN
3	L	211	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	418	GLN
3	L	100	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
5	SO4	A	602	-	4,4,4	1.06	0	6,6,6	0.50	0
5	SO4	A	603	-	4,4,4	0.14	0	6,6,6	0.24	0
5	SO4	A	604	-	4,4,4	0.23	0	6,6,6	0.15	0
5	SO4	H	301	-	4,4,4	0.33	0	6,6,6	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SO4	A	602	-	-	0/0/0/0	0/0/0/0
5	SO4	A	603	-	-	0/0/0/0	0/0/0/0
5	SO4	A	604	-	-	0/0/0/0	0/0/0/0
5	SO4	H	301	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	220/232 (94%)	0.25	4 (1%)	71	77	41, 55, 78, 124	0
2	H	215/228 (94%)	0.50	18 (8%)	14	18	35, 56, 101, 132	0
3	L	213/215 (99%)	0.33	10 (4%)	35	43	39, 57, 95, 110	0
All	All	648/675 (96%)	0.36	32 (4%)	33	41	35, 56, 92, 132	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	127	SER	8.7
2	H	128	SER	7.7
1	A	278	GLN	6.9
2	H	211	VAL	5.8
2	H	189	LEU	5.1
2	H	188	SER	5.0
1	A	279	ILE	4.9
2	H	191	THR	4.7
3	L	191	VAL	4.4
1	A	281	PHE	3.9
3	L	192	TYR	3.8
2	H	125	ALA	3.5
2	H	212	GLU	3.5
3	L	125	LEU	3.3
3	L	190	LYS	3.0
3	L	154	LEU	2.9
3	L	209	PHE	2.9
2	H	210	LYS	2.8
2	H	126	PRO	2.6
3	L	148	TRP	2.6
2	H	138	LEU	2.6
3	L	188	LYS	2.6
3	L	184	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	154	TRP	2.5
2	H	194	TYR	2.4
2	H	158	ALA	2.4
2	H	56	TYR	2.2
1	A	292	HIS	2.1
2	H	124	LEU	2.0
2	H	204	ASN	2.0
3	L	208	SER	2.0
2	H	190	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å^2)	Q<0.9
5	SO4	H	301	5/5	0.87	0.24	6.04	91,99,101,102	0
5	SO4	A	602	5/5	0.99	0.16	0.46	49,50,52,58	0
5	SO4	A	604	5/5	0.82	0.16	-0.34	108,109,112,112	0
4	CA	A	601	1/1	1.00	0.09	-1.68	54,54,54,54	0
5	SO4	A	603	5/5	0.91	0.20	-	89,91,94,96	0

6.5 Other polymers (i)

There are no such residues in this entry.