



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2016 – 09:23 PM EDT

PDB ID : 4ZI3
Title : BART-like domain of BARTL1/CCDC104 aa1-133 in complex with Arl3FL bound to GppNHp in P1 21 1
Authors : Lokaj, M.; Koerner, C.; Koesling, S.; Wittinghofer, A.
Deposited on : 2015-04-27
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027107
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0122
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027107


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	190	<div> <div>3%</div> <div>82%</div> <div>13%</div> <div>• 5%</div> </div>
1	B	190	<div> <div>19%</div> <div>70%</div> <div>17%</div> <div>• 11%</div> </div>
2	C	135	<div> <div>2%</div> <div>82%</div> <div>9%</div> <div>9%</div> </div>
3	D	133	<div> <div>4%</div> <div>86%</div> <div>11%</div> <div>•</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5335 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-ribosylation factor-like protein 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	180	Total	C	N	O	S	0	7	0
			1497	947	265	281	4			
1	B	170	Total	C	N	O	S	0	2	0
			1370	870	236	260	4			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	183	LEU	-	expression tag	UNP Q9WUL7
A	184	GLU	-	expression tag	UNP Q9WUL7
A	185	HIS	-	expression tag	UNP Q9WUL7
A	186	HIS	-	expression tag	UNP Q9WUL7
A	187	HIS	-	expression tag	UNP Q9WUL7
A	188	HIS	-	expression tag	UNP Q9WUL7
A	189	HIS	-	expression tag	UNP Q9WUL7
A	190	HIS	-	expression tag	UNP Q9WUL7
B	183	LEU	-	expression tag	UNP Q9WUL7
B	184	GLU	-	expression tag	UNP Q9WUL7
B	185	HIS	-	expression tag	UNP Q9WUL7
B	186	HIS	-	expression tag	UNP Q9WUL7
B	187	HIS	-	expression tag	UNP Q9WUL7
B	188	HIS	-	expression tag	UNP Q9WUL7
B	189	HIS	-	expression tag	UNP Q9WUL7
B	190	HIS	-	expression tag	UNP Q9WUL7

- Molecule 2 is a protein called Cilia- and flagella-associated protein 36.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	123	Total	C	N	O	S	0	4	0
			1040	662	164	209	5			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	GLY	-	expression tag	UNP Q8C6E0
C	0	PRO	-	expression tag	UNP Q8C6E0

- Molecule 3 is a protein called Cilia- and flagella-associated protein 36.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	128	Total	C	N	O	S	0	5	0
			1088	691	167	224	6			

- Molecule 4 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (three-letter code: GNP) (formula: $C_{10}H_{17}N_6O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			32	10	6	13	3		
4	B	1	Total	C	N	O	P	0	0
			32	10	6	13	3		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Mg	0	0
			1	1		
5	A	1	Total	Mg	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	89	Total 89	O 89	0	0
6	B	23	Total 23	O 23	0	0
6	C	87	Total 87	O 87	0	0
6	D	75	Total 75	O 75	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ADP-ribosylation factor-like protein 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.55Å 67.72Å 98.47Å 90.00° 102.65° 90.00°	Depositor
Resolution (Å)	28.95 – 2.00 28.95 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.6 (28.95-2.00) 99.6 (28.95-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 2.00Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.189 , 0.243 0.189 , 0.243	Depositor DCC
R_{free} test set	2234 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	32.8	Xtriage
Anisotropy	0.353	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 41.8	EDS
Estimated twinning fraction	0.024 for h,-k,-h-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 44701 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5335	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GNP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/1520	0.58	2/2052 (0.1%)
1	B	0.34	0/1390	0.63	2/1878 (0.1%)
2	C	0.46	0/1057	0.52	0/1428
3	D	0.41	0/1105	0.52	0/1491
All	All	0.41	0/5072	0.57	4/6849 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	161	LEU	CB-CG-CD1	-7.41	98.40	111.00
1	A	72	ARG	NE-CZ-NH1	5.80	123.20	120.30
1	B	142	GLY	N-CA-C	-5.06	100.46	113.10
1	A	151	ARG	NE-CZ-NH2	-5.05	117.77	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	137	SER	Peptide
1	B	141	GLU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1497	0	1519	13	0
1	B	1370	0	1391	36	0
2	C	1040	0	1017	10	0
3	D	1088	0	1054	12	0
4	A	32	0	13	0	0
4	B	32	0	13	4	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	89	0	0	1	1
6	B	23	0	0	3	0
6	C	87	0	0	4	0
6	D	75	0	0	4	0
All	All	5335	0	5007	68	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:96:ILE:HD11	6:D:209:HOH:O	1.80	0.81
1:B:9:LYS:HD3	3:D:84:THR:HG22	1.63	0.79
1:B:113[A]:GLU:OE2	6:B:401:HOH:O	2.00	0.78
1:A:87:ILE:HD13	1:A:173:VAL:HG13	1.71	0.73
1:B:87:ILE:HG21	1:B:173:VAL:HG13	1.72	0.72
1:B:79:ARG:HG3	6:B:415:HOH:O	1.90	0.71
2:C:89:LYS:NZ	6:C:202:HOH:O	2.24	0.71
3:D:80:GLN:O	3:D:84:THR:HG23	1.92	0.69
1:B:174:CYS:HA	1:B:177:VAL:HG23	1.73	0.69
1:B:137:SER:HA	1:B:140:ALA:H	1.57	0.68
1:B:79:ARG:NH1	1:B:113[A]:GLU:OE1	2.27	0.66
1:B:161:LEU:H	1:B:161:LEU:HD12	1.60	0.66
1:A:3:LEU:HD22	1:A:4:LEU:H	1.62	0.65
3:D:21:ARG:NH2	6:D:203:HOH:O	2.28	0.65
2:C:31[B]:ASP:OD1	6:C:201:HOH:O	2.15	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:17:ALA:HB2	2:C:92:THR:HG22	1.79	0.63
1:B:113[B]:GLU:CD	6:B:401:HOH:O	2.39	0.60
1:B:137:SER:HB3	1:B:140:ALA:CB	2.33	0.59
1:B:97:ARG:NH1	1:B:138:GLU:OE2	2.35	0.58
1:B:141:GLU:OE1	1:B:145:LEU:N	2.36	0.58
1:B:137:SER:HB3	1:B:140:ALA:HB2	1.86	0.58
1:A:120:PRO:HG3	1:A:179:ALA:HB3	1.86	0.57
3:D:84:THR:HG21	6:D:221:HOH:O	2.05	0.57
1:B:140:ALA:HA	1:B:145:LEU:HD12	1.87	0.56
1:B:152:VAL:HG12	1:B:172:TRP:HZ2	1.71	0.56
1:B:130:LEU:HD22	1:B:132:THR:H	1.71	0.55
1:B:79:ARG:NH1	1:B:113[B]:GLU:OE2	2.40	0.54
1:B:121:VAL:HG12	1:B:151:ARG:HD3	1.88	0.54
1:A:178:ASN:O	1:A:182:LYS:HG3	2.07	0.54
1:B:161:LEU:HB2	1:B:162:THR:HG23	1.89	0.54
3:D:115:GLN:NE2	6:D:201:HOH:O	2.25	0.54
1:B:59:GLN:NE2	1:B:174:CYS:SG	2.81	0.53
3:D:33:VAL:HA	3:D:113[A]:MET:HE1	1.91	0.52
1:A:36:GLN:HG3	1:A:160:ALA:O	2.10	0.52
1:B:3:LEU:HB3	1:B:6:ILE:HD12	1.93	0.51
2:C:14:GLU:OE2	2:C:91:ARG:HD2	2.11	0.51
1:A:13:ALA:O	1:A:16:GLN:HB3	2.12	0.49
1:B:161:LEU:HD11	4:B:301:GNP:N1	2.28	0.48
3:D:92:THR:OG1	3:D:97:LEU:HG	2.14	0.48
1:A:54:LYS:NZ	2:C:44:GLU:OE2	2.46	0.48
1:B:59:GLN:OE1	1:B:174:CYS:HB2	2.14	0.47
1:A:38:ALA:HB2	1:A:54:LYS:HG2	1.96	0.46
1:B:23:LEU:HD21	1:B:68[A]:ILE:HD13	1.96	0.46
2:C:31[A]:ASP:OD1	3:D:126:ARG:NH1	2.46	0.45
1:B:59:GLN:HB2	1:B:61:PHE:CD2	2.52	0.45
1:B:127:LYS:HG2	4:B:301:GNP:C6	2.47	0.45
1:B:72:ARG:HD2	1:B:102:GLU:OE2	2.16	0.45
1:B:58:SER:C	1:B:59:GLN:HG3	2.36	0.45
1:A:8:ARG:O	1:A:11:LYS:HB2	2.16	0.44
1:B:161:LEU:HD11	4:B:301:GNP:HN1	1.83	0.44
3:D:17:ALA:HB2	3:D:92:THR:HG22	1.99	0.44
3:D:124:ALA:O	3:D:128:ILE:HG13	2.18	0.44
1:A:3:LEU:HB3	1:A:6:ILE:HD12	2.00	0.44
2:C:58:LYS:NZ	6:C:211:HOH:O	2.52	0.43
1:B:161:LEU:HD21	4:B:301:GNP:C2	2.48	0.43
1:B:141:GLU:HA	1:B:144:ASN:H	1.84	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:144:ASN:O	1:B:147:THR:OG1	2.24	0.42
1:A:72:ARG:HD2	1:A:102:GLU:OE2	2.19	0.42
1:B:137:SER:CA	1:B:140:ALA:H	2.29	0.42
1:A:40:GLU:OE2	1:A:54:LYS:HE2	2.20	0.41
1:B:137:SER:HB3	1:B:140:ALA:HB3	2.03	0.41
6:A:406:HOH:O	2:C:104:GLU:HB2	2.19	0.41
2:C:115:GLN:HG2	6:C:238:HOH:O	2.20	0.41
3:D:123:GLN:NE2	3:D:126:ARG:HH21	2.18	0.41
1:A:82:PHE:CE2	1:A:113[A]:GLU:HG2	2.55	0.41
1:B:125:ALA:HB3	1:B:157:SER:HA	2.02	0.41
1:B:68[A]:ILE:HD12	1:B:68[A]:ILE:N	2.36	0.40
3:D:33:VAL:HG12	3:D:116:LYS:HG2	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:485:HOH:O	6:A:487:HOH:O[2_656]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	184/190 (97%)	182 (99%)	2 (1%)	0	100	100
1	B	168/190 (88%)	156 (93%)	12 (7%)	0	100	100
2	C	125/135 (93%)	124 (99%)	1 (1%)	0	100	100
3	D	131/133 (98%)	129 (98%)	2 (2%)	0	100	100
All	All	608/648 (94%)	591 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/167 (99%)	163 (98%)	3 (2%)	66	69
1	B	152/167 (91%)	150 (99%)	2 (1%)	76	79
2	C	116/120 (97%)	115 (99%)	1 (1%)	84	88
3	D	121/119 (102%)	121 (100%)	0	100	100
All	All	555/573 (97%)	549 (99%)	6 (1%)	84	83

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	83	GLU
1	A	157[A]	SER
1	A	157[B]	SER
1	B	17	GLU
1	B	130	LEU
2	C	129	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	59	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GNP	A	301	5	29,34,34	2.69	12 (41%)	28,54,54	1.91	7 (25%)
4	GNP	B	301	5	29,34,34	2.78	10 (34%)	28,54,54	1.91	5 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GNP	A	301	5	-	0/16/38/38	0/3/3/3
4	GNP	B	301	5	-	0/16/38/38	0/3/3/3

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	301	GNP	C4-N9	-7.62	1.37	1.47
4	A	301	GNP	C4-N9	-7.33	1.38	1.47
4	B	301	GNP	C5-C6	-6.43	1.41	1.53
4	A	301	GNP	C5-C6	-6.04	1.42	1.53
4	B	301	GNP	PB-O3A	-4.22	1.53	1.59
4	A	301	GNP	PB-O2B	-3.81	1.46	1.56
4	B	301	GNP	PB-O2B	-3.35	1.47	1.56
4	A	301	GNP	PG-O3G	-3.09	1.48	1.56
4	B	301	GNP	C8-N9	-2.86	1.38	1.47
4	A	301	GNP	C8-N9	-2.81	1.38	1.47
4	B	301	GNP	C2-N3	-2.61	1.32	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	301	GNP	C2-N3	-2.47	1.33	1.43
4	A	301	GNP	PG-O2G	-2.11	1.51	1.56
4	A	301	GNP	PB-O3A	-2.08	1.56	1.59
4	B	301	GNP	C1'-N9	2.16	1.46	1.42
4	A	301	GNP	PB-O1B	2.30	1.48	1.46
4	A	301	GNP	C1'-N9	2.40	1.46	1.42
4	B	301	GNP	PB-O1B	3.48	1.49	1.46
4	A	301	GNP	C6-N1	3.84	1.39	1.33
4	B	301	GNP	C6-N1	4.13	1.40	1.33
4	B	301	GNP	PG-O1G	5.44	1.52	1.46
4	A	301	GNP	PG-O1G	5.88	1.52	1.46

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	301	GNP	O3G-PG-O1G	-4.41	101.97	113.58
4	A	301	GNP	O3G-PG-O1G	-3.24	105.05	113.58
4	A	301	GNP	O6-C6-N1	-3.17	118.64	122.80
4	A	301	GNP	O2A-PA-O3A	2.29	115.08	105.27
4	B	301	GNP	O2B-PB-O1B	2.42	114.79	110.02
4	B	301	GNP	O3G-PG-O2G	2.79	115.78	107.67
4	A	301	GNP	O2B-PB-O1B	3.12	116.16	110.02
4	A	301	GNP	O3G-PG-O2G	3.28	117.20	107.67
4	B	301	GNP	O6-C6-C5	4.69	128.65	119.69
4	A	301	GNP	O6-C6-C5	4.86	128.97	119.69
4	A	301	GNP	C4-C5-N7	5.01	110.48	102.67
4	B	301	GNP	C4-C5-N7	5.54	111.31	102.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	301	GNP	4	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	180/190 (94%)	0.13	5 (2%) 56 57	21, 35, 61, 74	0
1	B	170/190 (89%)	1.06	36 (21%) 1 1	31, 67, 95, 110	0
2	C	123/135 (91%)	-0.25	3 (2%) 62 63	23, 35, 59, 69	0
3	D	128/133 (96%)	-0.15	5 (3%) 43 45	24, 37, 60, 97	0
All	All	601/648 (92%)	0.26	49 (8%) 14 15	21, 41, 87, 110	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	136	ALA	7.2
3	D	128	ILE	7.0
1	B	131	LEU	6.0
1	B	149	ARG	6.0
1	B	61	PHE	5.7
1	B	174	CYS	5.6
1	B	142	GLY	5.0
1	B	177	VAL	4.7
1	B	59	GLN	4.5
1	B	57	GLN	4.5
2	C	7	ASP	4.4
1	B	146	HIS	4.3
1	B	161	LEU	4.0
3	D	130	GLU	4.0
1	A	183	LEU	3.6
1	B	130	LEU	3.4
1	B	137	SER	3.3
1	B	60	GLY	3.2
1	B	117	SER	3.1
1	B	43	SER	3.0
3	D	127	ILE	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	141	GLU	2.9
1	B	132	THR	2.9
1	B	140	ALA	2.7
1	B	150	ASP	2.7
1	B	135	PRO	2.7
1	A	16	GLN	2.6
1	A	89	ILE	2.6
1	B	133	ALA	2.5
1	B	129	ASP	2.5
1	B	175	LYS	2.5
1	B	90	TYR	2.5
1	B	91	VAL	2.4
1	B	41	ASP	2.3
3	D	3	ALA	2.3
1	B	173	VAL	2.2
1	B	168	ASP	2.2
1	B	42	ILE	2.2
1	B	148	ILE	2.2
1	B	122	LEU	2.2
1	B	143	LEU	2.1
1	B	58	SER	2.1
2	C	128	ILE	2.1
1	A	23	LEU	2.1
2	C	90	THR	2.1
1	B	8	ARG	2.1
3	D	129	GLN	2.1
1	A	12	SER	2.0
1	B	44	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	GNP	B	301	32/32	0.94	0.11	-0.78	34,51,69,73	0
4	GNP	A	301	32/32	0.98	0.09	-0.87	18,27,33,35	0
5	MG	A	302	1/1	0.99	0.07	-	25,25,25,25	0
5	MG	B	302	1/1	0.83	0.08	-	44,44,44,44	0

6.5 Other polymers

There are no such residues in this entry.