



wwPDB X-ray Structure Validation Summary Report ⓘ

May 23, 2016 – 05:08 AM EDT

PDB ID : 4ZOD
Title : Crystal Structure of beta-glucosidase from *Listeria innocua* in complex with glucose
Authors : Nakajima, M.; Yoshida, R.; Miyanaga, A.; Abe, K.; Takahashi, Y.; Sugimoto, N.; Toyozumi, H.; Nakai, H.; Kitaoka, M.; Taguchi, H.
Deposited on : 2015-05-06
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027457
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	731	<div> <div>4%</div> <div>88%</div> <div>10%</div> <div>..</div> </div>
1	B	731	<div> <div>5%</div> <div>89%</div> <div>9%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BGC	B	803	-	-	-	X
4	GOL	A	803[A]	-	-	-	X
4	GOL	A	803[B]	-	-	-	X
4	GOL	A	804[A]	-	-	-	X
4	GOL	A	804[B]	-	-	-	X
4	GOL	A	805	-	-	-	X
4	GOL	A	808[A]	-	-	-	X
4	GOL	B	805[A]	-	-	-	X
4	GOL	B	805[B]	-	-	-	X
5	PEG	A	806	-	-	-	X
6	SO4	A	807	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 11754 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lin1840 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	720	Total	C	N	O	S	0	2	0
			5578	3533	923	1097	25			
1	B	721	Total	C	N	O	S	0	1	0
			5578	3531	923	1099	25			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	GLU	LYS	engineered mutation	UNP Q92AS9
A	724	LEU	-	expression tag	UNP Q92AS9
A	725	GLU	-	expression tag	UNP Q92AS9
A	726	HIS	-	expression tag	UNP Q92AS9
A	727	HIS	-	expression tag	UNP Q92AS9
A	728	HIS	-	expression tag	UNP Q92AS9
A	729	HIS	-	expression tag	UNP Q92AS9
A	730	HIS	-	expression tag	UNP Q92AS9
A	731	HIS	-	expression tag	UNP Q92AS9
B	2	GLU	LYS	engineered mutation	UNP Q92AS9
B	724	LEU	-	expression tag	UNP Q92AS9
B	725	GLU	-	expression tag	UNP Q92AS9
B	726	HIS	-	expression tag	UNP Q92AS9
B	727	HIS	-	expression tag	UNP Q92AS9
B	728	HIS	-	expression tag	UNP Q92AS9
B	729	HIS	-	expression tag	UNP Q92AS9
B	730	HIS	-	expression tag	UNP Q92AS9
B	731	HIS	-	expression tag	UNP Q92AS9

- Molecule 2 is BETA-D-GLUCOSE (three-letter code: BGC) (formula: $C_6H_{12}O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	1
			12	6	6		
4	A	1	Total	C	O	0	1
			12	6	6		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	1
			12	6	6		
4	B	1	Total	C	O	0	1
			12	6	6		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	242	Total 242	O 242	0	0
7	B	233	Total 233	O 233	0	0

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Chain A:

4% 88% 10%

Chain A: M1 E2 Q3 E4 K5 S11 T14 E17 K18 F28 L29 F30 T33 N34 A37 E38 I39 T40 G41 P42 L43 L44 Q45 E46 V60 L61 L67 R81 L82 L104 A119 H132 L139 D140 L141 V142 R143 L160 L164 G171 Y172 Q173 V196 M210 E214 Y219 L220 P221 M235 Q243 I244 K250 W251 L252 V256 L257 R258 F263 I268 S269 D270 W271 N285 P286 D299 L300 T305 I308 H309 G318 K319 L320 S321 E322 R331 L339 L349 LVS ASN ASN ASP R354 T355 K356 V357 V396 L406 E430 T431 V435 E438 Q455 W456 L462 L481 V501 I502 N506 G529 T530 E531 E534 L550 T575 P576 E577 I578 R579 E580 E606 S613 V626 R655 R658 E664 V676 T677 E683 R702 S709

Chain B:

5% 89% 9% ..

M1 K5 L9 V10 S11 Q12 M13 T14 E17 L23 F28 L29 F30 T33 N34 K35 N36 A37 E38 L39 T40 G41 P42 L43 L44 Q45 D51 E55 V60 L61 G62 S63 A66 L67 K78 R81 L82 I93 H94 L104 D129 H132 F135 M138 M151 K166 L183 P210 E214 Y219 P245 V248 R258 F263 T268 S269 D270 W271 V274 V277 E280 R284 N285 F286 D299 L300 T308 L320 K323 R331 L339 G348 L349 LYS ASN D353 R354 T355 W356 D357 I358 R368 R383 K389 V396 L406 V428 F429 E430 V435 E438 Q455 L462 V501 I502 R508 F576 E577 N578 K579 G580 S613 L614 D635 L636 I653 K658 E664 L680 E683 H690 R702 S709 L724 C725

4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	89.79 Å 95.41 Å 213.23 Å 90.00° 97.27° 90.00°	Depositor
Resolution (Å)	37.10 – 2.10 37.11 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.5 (37.10-2.10) 97.6 (37.11-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 2.10 Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.167 , 0.233 0.177 , 0.236	Depositor DCC
R_{free} test set	5084 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	25.0	Xtriage
Anisotropy	0.062	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 53.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11754	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, GOL, MG, BGC, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/5679	0.92	3/7696 (0.0%)
1	B	0.82	0/5679	0.91	8/7695 (0.1%)
All	All	0.84	0/11358	0.91	11/15391 (0.1%)

There are no bond length outliers.

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	702	ARG	NE-CZ-NH1	7.54	124.07	120.30
1	A	702	ARG	NE-CZ-NH1	7.41	124.00	120.30
1	B	368	ARG	NE-CZ-NH1	-6.76	116.92	120.30
1	B	210[A]	MET	CG-SD-CE	-6.62	89.61	100.20
1	B	210[B]	MET	CG-SD-CE	-6.62	89.61	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5578	0	5534	46	0
1	B	5578	0	5527	41	0
2	A	12	0	12	1	0
2	B	24	0	24	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	42	0	56	5	0
4	B	24	0	32	2	0
5	A	7	0	10	1	0
5	B	7	0	10	1	0
6	A	5	0	0	0	0
7	A	242	0	0	2	0
7	B	233	0	0	3	0
All	All	11754	0	11205	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 86 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:299:ASP:OD1	1:B:331:ARG:NH1	2.08	0.86
1:A:244[A]:ILE:HD11	1:B:653:ILE:HD13	1.66	0.75
1:A:299:ASP:OD1	1:A:331:ARG:NH1	2.20	0.74
1:A:270:ASP:OD1	2:A:801:BGC:H1	1.91	0.71
1:A:210[A]:MET:HE1	1:A:219:TYR:CD1	2.29	0.67

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	718/731 (98%)	696 (97%)	22 (3%)	0	100	100
1	B	718/731 (98%)	694 (97%)	24 (3%)	0	100	100
All	All	1436/1462 (98%)	1390 (97%)	46 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	603/612 (98%)	587 (97%)	16 (3%)	52	56
1	B	603/612 (98%)	585 (97%)	18 (3%)	48	51
All	All	1206/1224 (98%)	1172 (97%)	34 (3%)	51	55

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	683	GLU
1	B	78	LYS
1	B	664	GLU
1	B	61	LEU
1	A	320	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	185	GLN
1	B	691	GLN
1	B	625	HIS
1	B	182	ASN
1	B	690	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 2 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BGC	A	801	-	12,12,12	0.91	0	17,17,17	1.55	3 (17%)
4	GOL	A	803[A]	-	5,5,5	0.68	0	5,5,5	1.29	0
4	GOL	A	803[B]	-	5,5,5	0.73	0	5,5,5	0.82	0
4	GOL	A	804[A]	-	5,5,5	0.65	0	5,5,5	0.68	0
4	GOL	A	804[B]	-	5,5,5	0.34	0	5,5,5	0.37	0
4	GOL	A	805	-	5,5,5	0.44	0	5,5,5	0.70	0
5	PEG	A	806	-	6,6,6	0.47	0	5,5,5	1.35	1 (20%)
6	SO4	A	807	-	4,4,4	0.61	0	6,6,6	0.30	0
4	GOL	A	808[A]	-	5,5,5	0.45	0	5,5,5	1.20	0
4	GOL	A	808[B]	-	5,5,5	0.51	0	5,5,5	0.80	0
5	PEG	B	801	-	6,6,6	0.66	0	5,5,5	0.25	0
2	BGC	B	802	-	12,12,12	0.67	0	17,17,17	1.35	3 (17%)
2	BGC	B	803	-	12,12,12	0.96	0	17,17,17	2.69	8 (47%)
4	GOL	B	805[A]	-	5,5,5	0.46	0	5,5,5	0.52	0
4	GOL	B	805[B]	-	5,5,5	0.45	0	5,5,5	0.63	0
4	GOL	B	806	-	5,5,5	0.21	0	5,5,5	0.55	0
4	GOL	B	807	-	5,5,5	0.48	0	5,5,5	1.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	A	801	-	-	0/2/22/22	0/1/1/1
4	GOL	A	803[A]	-	-	0/4/4/4	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	803[B]	-	-	0/4/4/4	0/0/0/0
4	GOL	A	804[A]	-	-	0/4/4/4	0/0/0/0
4	GOL	A	804[B]	-	-	0/4/4/4	0/0/0/0
4	GOL	A	805	-	-	0/4/4/4	0/0/0/0
5	PEG	A	806	-	-	0/4/4/4	0/0/0/0
6	SO4	A	807	-	-	0/0/0/0	0/0/0/0
4	GOL	A	808[A]	-	-	0/4/4/4	0/0/0/0
4	GOL	A	808[B]	-	-	0/4/4/4	0/0/0/0
5	PEG	B	801	-	-	0/4/4/4	0/0/0/0
2	BGC	B	802	-	-	0/2/22/22	0/1/1/1
2	BGC	B	803	-	-	0/2/22/22	0/1/1/1
4	GOL	B	805[A]	-	-	0/4/4/4	0/0/0/0
4	GOL	B	805[B]	-	-	0/4/4/4	0/0/0/0
4	GOL	B	806	-	-	0/4/4/4	0/0/0/0
4	GOL	B	807	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	803	BGC	C3-C4-C5	-3.85	103.36	110.23
2	A	801	BGC	O5-C5-C4	-3.78	102.45	109.67
2	B	803	BGC	O2-C2-C3	-3.58	102.28	110.36
2	B	803	BGC	O1-C1-O5	-3.07	101.78	110.33
2	B	803	BGC	O6-C6-C5	-2.79	101.98	111.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	BGC	1	0
4	A	803[B]	GOL	1	0
4	A	804[A]	GOL	3	0
4	A	805	GOL	1	0
5	A	806	PEG	1	0
5	B	801	PEG	1	0
2	B	802	BGC	1	0
4	B	805[B]	GOL	1	0
4	B	807	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	720/731 (98%)	0.09	31 (4%) 39 48	12, 26, 50, 73	0
1	B	721/731 (98%)	0.11	34 (4%) 35 44	12, 27, 50, 70	0
All	All	1441/1462 (98%)	0.10	65 (4%) 37 46	12, 27, 50, 73	0

The worst 5 of 65 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	579	LYS	6.0
1	B	579	LYS	5.2
1	A	39	LEU	4.7
1	A	576	PRO	4.2
1	B	34	ASN	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	GOL	A	805	6/6	0.93	0.26	5.35	39,44,47,50	0
2	BGC	B	803	12/12	0.92	0.21	5.03	33,41,55,57	0
4	GOL	A	808[A]	6/6	0.92	0.18	4.00	9,18,23,25	6
4	GOL	B	805[A]	6/6	0.96	0.21	3.88	22,23,23,23	6
4	GOL	B	805[B]	6/6	0.96	0.21	3.52	17,22,24,26	6
4	GOL	A	803[B]	6/6	0.90	0.17	3.52	19,20,21,22	6
4	GOL	A	803[A]	6/6	0.90	0.17	3.24	18,23,24,25	6
4	GOL	A	804[A]	6/6	0.94	0.19	3.22	10,14,14,17	6
5	PEG	A	806	7/7	0.94	0.12	3.09	32,36,39,43	0
4	GOL	A	804[B]	6/6	0.94	0.19	3.04	18,26,28,36	6
6	SO4	A	807	5/5	0.97	0.20	2.71	16,17,18,18	5
4	GOL	B	806	6/6	0.96	0.14	1.60	40,47,50,50	0
4	GOL	B	807	6/6	0.94	0.13	1.42	32,39,44,48	0
5	PEG	B	801	7/7	0.92	0.10	-0.13	39,44,50,53	0
2	BGC	B	802	12/12	0.97	0.14	-0.28	20,24,26,28	0
2	BGC	A	801	12/12	0.97	0.13	-0.41	15,19,20,22	0
3	MG	A	802	1/1	0.96	0.05	-2.31	24,24,24,24	0
3	MG	B	804	1/1	0.99	0.03	-4.83	24,24,24,24	0
4	GOL	A	808[B]	6/6	0.92	0.18	-	22,23,24,26	6

6.5 Other polymers [i](#)

There are no such residues in this entry.