



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:31 PM GMT

PDB ID : 3ZPH
Title : Bacterial chalcone isomerase in closed conformation from Eubacterium ramulus at 2.8 Å resolution
Authors : Thomsen, M.; Palm, G.J.; Hinrichs, W.
Deposited on : 2013-02-27
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	<div> <div>%</div> <div>84% 9% 8%</div> </div>
1	B	282	<div> <div>2%</div> <div>91% 9%</div> </div>
1	C	282	<div> <div>%</div> <div>82% 9% 9%</div> </div>
1	D	282	<div> <div>%</div> <div>83% 8% 9%</div> </div>
1	E	282	<div> <div>%</div> <div>81% 11% 9%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	282	<div> <div></div> <div>84%</div> <div>7% • 9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	301	-	-	-	X
2	GOL	C	301	-	-	-	X
2	GOL	D	401	-	-	-	X
2	GOL	F	301	-	-	-	X

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 14021 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHALCONE ISOMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	0	0	0
			2132	1392	347	380	13			
1	B	282	Total	C	N	O	S	0	0	0
			2287	1478	379	417	13			
1	C	258	Total	C	N	O	S	0	1	0
			2125	1388	345	379	13			
1	D	258	Total	C	N	O	S	0	2	0
			2131	1392	345	381	13			
1	E	258	Total	C	N	O	S	0	1	0
			2125	1388	345	379	13			
1	F	258	Total	C	N	O	S	0	1	0
			2125	1388	345	379	13			

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0
2	E	1	Total C O 6 3 3	0	0
2	F	1	Total C O 6 3 3	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Cl 1 1	0	0
3	A	1	Total Cl 1 1	0	0
3	C	1	Total Cl 1 1	0	0
3	E	1	Total Cl 1 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	214	Total O 215 215	0	1
4	B	196	Total O 196 196	0	0
4	C	177	Total O 177 177	0	0
4	D	163	Total O 163 163	0	0
4	E	144	Total O 144 144	0	0
4	F	167	Total O 167 167	0	0



• Molecule 1: CHALCONE ISOMERASE



4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	177.73Å 203.12Å 206.11Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.09 – 2.80 20.08 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.0 (20.09-2.80) 99.3 (20.08-2.80)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 2.79Å)	Xtriage
Refinement program	REFMAC 5.7.0027	Depositor
R, R_{free}	0.248 , 0.297 0.250 , 0.292	Depositor DCC
R_{free} test set	4539 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	42.9	Xtriage
Anisotropy	0.323	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 31.2	EDS
Estimated twinning fraction	0.000 for -h,-l,-k	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.55$, $\langle L^2 \rangle = 0.40$	Xtriage
Outliers	2 of 90731 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	14021	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	2/2203 (0.1%)	0.77	1/2995 (0.0%)
1	B	0.70	1/2360 (0.0%)	0.77	0/3209
1	C	0.70	0/2199	0.76	0/2989
1	D	0.69	1/2208 (0.0%)	0.72	0/3001
1	E	0.72	3/2199 (0.1%)	0.78	0/2989
1	F	0.69	0/2199	0.75	0/2989
All	All	0.70	7/13368 (0.1%)	0.76	1/18172 (0.0%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	220	TRP	CD2-CE2	5.68	1.48	1.41
1	E	156	GLU	CG-CD	5.66	1.60	1.51
1	E	243	TRP	CD2-CE2	5.45	1.47	1.41
1	B	144	TRP	CD2-CE2	5.20	1.47	1.41
1	E	220	TRP	CD2-CE2	5.17	1.47	1.41
1	A	100	TRP	CD2-CE2	5.10	1.47	1.41
1	A	215	TRP	CD2-CE2	5.10	1.47	1.41

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	233	ASP	CB-CG-OD1	5.32	123.09	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2132	0	2062	12	0
1	B	2287	0	2197	17	0
1	C	2125	0	2057	14	0
1	D	2131	0	2063	17	0
1	E	2125	0	2057	14	0
1	F	2125	0	2057	13	0
2	A	6	0	8	0	0
2	C	6	0	8	0	0
2	D	6	0	8	2	0
2	E	6	0	8	0	0
2	F	6	0	8	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	E	1	0	0	0	0
4	A	215	0	0	6	0
4	B	196	0	0	5	0
4	C	177	0	0	3	0
4	D	163	0	0	3	0
4	E	144	0	0	1	0
4	F	167	0	0	2	0
All	All	14021	0	12533	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:85:GLU:HG2	4:B:2077:HOH:O	1.88	0.74
1:B:108:ALA:O	1:B:109:HIS:CD2	2.45	0.70
1:A:183:PHE:CE2	4:A:2174:HOH:O	2.48	0.67
1:A:266:TYR:OH	4:A:2201:HOH:O	2.11	0.67
1:A:182:LEU:HD23	4:A:2174:HOH:O	1.94	0.66
1:A:167:ILE:HD11	1:A:182:LEU:HD22	1.81	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:206:LYS:CE	1:E:214:ASP:OD1	2.48	0.62
1:E:206:LYS:HE2	1:E:214:ASP:OD1	2.01	0.61
1:C:206:LYS:HE3	1:C:213:MET:O	2.04	0.58
1:D:135:PHE:HE1	2:D:401:GOL:HO1	1.50	0.58
1:D:106:ALA:C	4:D:2074:HOH:O	2.41	0.57
1:D:131:GLU:N	4:D:2078:HOH:O	2.38	0.57
1:E:81:ASP:HB3	1:E:84:LEU:HD13	1.88	0.56
1:D:135:PHE:HE1	2:D:401:GOL:O1	1.89	0.56
1:A:14:VAL:O	1:A:66:ALA:HB1	2.08	0.54
4:C:2048:HOH:O	1:F:282:ARG:HG3	2.08	0.54
1:C:15:ASP:HB2	4:C:2019:HOH:O	2.08	0.53
1:F:106:ALA:C	4:F:2066:HOH:O	2.47	0.52
1:E:106:ALA:C	4:E:2071:HOH:O	2.48	0.51
1:E:206:LYS:HE3	1:E:214:ASP:OD1	2.11	0.50
1:B:146:LYS:HE3	4:B:2116:HOH:O	2.12	0.50
1:B:108:ALA:O	1:B:109:HIS:CG	2.64	0.50
1:A:182:LEU:HD21	1:A:217:LEU:HD11	1.93	0.50
1:B:195:GLU:HG3	4:B:2069:HOH:O	2.12	0.49
1:F:249:PHE:CG	1:F:250:PRO:HA	2.47	0.49
1:E:34:ILE:HB	1:E:35:PRO:HD3	1.94	0.49
1:F:22:ARG:HB3	1:F:23:PRO:HD3	1.95	0.49
1:B:1:ALA:HB1	4:B:2003:HOH:O	2.12	0.49
1:F:168:GLY:O	1:F:257:ASN:HB3	2.13	0.48
1:C:239:GLU:HG2	4:C:2069:HOH:O	2.12	0.48
1:C:103:GLN:HA	1:C:133:ASN:O	2.13	0.48
1:E:42:GLU:HB3	1:E:43:PRO:HD3	1.95	0.48
1:E:51:TYR:CE2	1:E:267:THR:HG22	2.49	0.48
1:C:249:PHE:CD1	1:C:250:PRO:HA	2.49	0.48
1:B:41:PHE:CD1	1:B:45:VAL:HB	2.49	0.48
1:B:91:GLU:OE2	1:B:121:GLY:HA3	2.15	0.47
1:C:149:LYS:O	1:C:198:ARG:HA	2.14	0.47
1:A:205:LYS:HE2	4:A:2168:HOH:O	2.14	0.47
1:C:159:ALA:HB2	1:F:27:ARG:HD2	1.96	0.46
1:F:42:GLU:N	1:F:43:PRO:CD	2.79	0.46
1:A:7:PRO:HA	1:A:75:TRP:O	2.16	0.46
1:A:27:ARG:HD2	1:E:159:ALA:HB2	1.98	0.46
1:D:17:VAL:HG11	1:D:102:GLY:HA3	1.97	0.46
1:D:182:LEU:HD23	1:D:183:PHE:CE2	2.52	0.45
1:D:106:ALA:C	4:D:2073:HOH:O	2.53	0.45
1:D:257:ASN:OD1	1:D:257:ASN:N	2.50	0.45
1:F:230:VAL:O	1:F:234:ASP:HB2	2.18	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:47:LYS:HA	1:D:272:LEU:HD12	2.00	0.44
1:B:96:ASP:HB3	1:B:100:TRP:CZ2	2.52	0.44
1:E:54:PHE:HB3	1:E:55:PRO:HD2	1.99	0.44
1:B:22:ARG:HB3	1:B:23:PRO:HD3	1.98	0.44
1:F:103:GLN:HG3	1:F:133:ASN:O	2.18	0.44
1:C:42:GLU:HB3	1:C:43:PRO:HD3	2.00	0.44
1:D:30:TYR:HA	1:D:34:ILE:HD12	1.99	0.44
1:D:149:LYS:O	1:D:198:ARG:HA	2.17	0.43
1:A:107:ALA:N	4:A:2097:HOH:O	2.50	0.43
1:C:12:ILE:HB	1:C:71:THR:HB	2.00	0.43
1:E:153:ARG:NH2	1:E:160:ASN:OD1	2.51	0.43
1:C:180:LYS:O	1:C:184:GLU:HB2	2.19	0.43
1:D:22:ARG:HB3	1:D:23:PRO:HD3	2.00	0.42
1:D:249:PHE:CD1	1:D:250:PRO:HA	2.54	0.42
1:B:142:MET:HB2	1:B:143:TRP:CZ3	2.54	0.42
1:B:37:SER:HB3	1:B:41:PHE:CE2	2.55	0.42
1:A:17:VAL:HB	4:A:2011:HOH:O	2.19	0.42
1:F:2:ASP:HB2	4:F:2003:HOH:O	2.20	0.42
1:B:41:PHE:HD1	1:B:45:VAL:HB	1.85	0.41
1:D:182:LEU:HD23	1:D:183:PHE:CZ	2.55	0.41
1:F:149:LYS:HD2	1:F:191:GLN:HA	2.01	0.41
1:B:103:GLN:HA	1:B:133:ASN:O	2.21	0.41
1:B:109:HIS:HB3	4:B:2090:HOH:O	2.19	0.41
1:B:159:ALA:HB2	1:D:27:ARG:HD2	2.03	0.41
1:E:155:ILE:HD12	1:E:155:ILE:HA	1.92	0.41
1:F:103:GLN:HA	1:F:133:ASN:O	2.20	0.41
1:A:42:GLU:N	1:A:43:PRO:CD	2.84	0.41
1:C:47:LYS:HA	1:C:272:LEU:HD12	2.03	0.41
1:E:140:LEU:HD21	1:E:216:VAL:CG2	2.51	0.41
1:C:249:PHE:CG	1:C:250:PRO:HA	2.56	0.41
1:D:168:GLY:O	1:D:257:ASN:HB3	2.21	0.40
1:F:150:GLY:O	1:F:151:LYS:C	2.59	0.40
1:C:155:ILE:HA	1:C:155:ILE:HD12	1.89	0.40
1:E:37:SER:HB3	1:E:41:PHE:CZ	2.56	0.40
1:B:225:SER:OG	1:D:20:ASP:OD2	2.36	0.40
1:C:229:LYS:O	1:C:233:ASP:HB2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/282 (91%)	246 (96%)	10 (4%)	0	100	100
1	B	280/282 (99%)	267 (95%)	12 (4%)	1 (0%)	39	74
1	C	255/282 (90%)	246 (96%)	8 (3%)	1 (0%)	39	74
1	D	256/282 (91%)	250 (98%)	6 (2%)	0	100	100
1	E	255/282 (90%)	243 (95%)	12 (5%)	0	100	100
1	F	255/282 (90%)	245 (96%)	10 (4%)	0	100	100
All	All	1557/1692 (92%)	1497 (96%)	58 (4%)	2 (0%)	56	87

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	207	ASP
1	C	207	ASP

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/240 (94%)	219 (97%)	6 (3%)	52	85
1	B	240/240 (100%)	237 (99%)	3 (1%)	76	94
1	C	226/240 (94%)	222 (98%)	4 (2%)	66	91
1	D	227/240 (95%)	226 (100%)	1 (0%)	93	98
1	E	226/240 (94%)	220 (97%)	6 (3%)	52	85

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	226/240 (94%)	223 (99%)	3 (1%)	76	94
All	All	1370/1440 (95%)	1347 (98%)	23 (2%)	68	92

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ASP
1	A	41	PHE
1	A	85	GLU
1	A	87	LYS
1	A	103	GLN
1	A	131	GLU
1	B	93	PHE
1	B	146	LYS
1	B	276	ARG
1	C	41	PHE
1	C	93	PHE
1	C	104	ILE
1	C	239	GLU
1	D	68	MET
1	E	95	MET
1	E	96	ASP
1	E	131	GLU
1	E	151	LYS
1	E	185	LYS
1	E	197	THR
1	F	41	PHE
1	F	103	GLN
1	F	151	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	133	ASN
1	B	109	HIS
1	C	101	GLN
1	C	103	GLN
1	D	101	GLN
1	D	103	GLN
1	E	101	GLN
1	F	101	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	301	-	5,5,5	0.30	0	5,5,5	0.45	0
2	GOL	C	301	-	5,5,5	0.44	0	5,5,5	0.70	0
2	GOL	D	401	-	5,5,5	0.41	0	5,5,5	0.56	0
2	GOL	E	301	-	5,5,5	0.24	0	5,5,5	0.60	0
2	GOL	F	301	-	5,5,5	0.36	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	301	-	-	0/4/4/4	0/0/0/0
2	GOL	C	301	-	-	0/4/4/4	0/0/0/0
2	GOL	D	401	-	-	0/4/4/4	0/0/0/0
2	GOL	E	301	-	-	0/4/4/4	0/0/0/0
2	GOL	F	301	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	401	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	260/282 (92%)	-0.55	4 (1%)	76 68	25, 32, 44, 61	0
1	B	282/282 (100%)	-0.38	7 (2%)	61 48	29, 36, 57, 92	0
1	C	258/282 (91%)	-0.52	3 (1%)	81 73	26, 34, 49, 60	0
1	D	258/282 (91%)	-0.47	4 (1%)	74 66	27, 34, 50, 63	0
1	E	258/282 (91%)	-0.42	3 (1%)	81 73	27, 36, 51, 66	0
1	F	258/282 (91%)	-0.38	4 (1%)	74 66	29, 38, 54, 77	0
All	All	1574/1692 (93%)	-0.45	25 (1%)	74 66	25, 35, 53, 92	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	109	HIS	4.7
1	A	1	ALA	4.2
1	B	117	ASP	3.4
1	C	106	ALA	3.3
1	F	1	ALA	3.3
1	B	115	ASP	3.3
1	B	1	ALA	3.1
1	D	106	ALA	3.1
1	F	131	GLU	3.0
1	E	131	GLU	2.9
1	E	1	ALA	2.8
1	F	210	GLY	2.6
1	D	171	GLU	2.5
1	D	172	GLY	2.5
1	C	207	ASP	2.4
1	B	239	GLU	2.3
1	C	171	GLU	2.3
1	A	130	ALA	2.2
1	B	131	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	1	ALA	2.1
1	E	106	ALA	2.1
1	B	130	ALA	2.1
1	A	171	GLU	2.1
1	A	106	ALA	2.1
1	F	106	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GOL	A	301	6/6	0.91	0.22	5.24	43,47,48,51	0
2	GOL	F	301	6/6	0.92	0.23	2.56	44,46,50,52	0
2	GOL	C	301	6/6	0.92	0.23	2.33	34,36,39,47	0
2	GOL	D	401	6/6	0.89	0.19	2.11	40,48,50,51	0
2	GOL	E	301	6/6	0.89	0.20	1.45	45,49,50,51	0
3	CL	A	401	1/1	0.89	0.25	-	54,54,54,54	0
3	CL	B	401	1/1	0.80	0.20	-	50,50,50,50	0
3	CL	E	401	1/1	0.88	0.13	-	43,43,43,43	0
3	CL	C	401	1/1	0.83	0.24	-	42,42,42,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.