



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 12:10 AM GMT

PDB ID : 1ZYC
Title : Crystal Structure of eIF2alpha Protein Kinase GCN2: Wild-Type in Apo Form.
Authors : Padyana, A.K.; Qiu, H.; Roll-Mecak, A.; Hinnebusch, A.G.; Burley, S.K.
Deposited on : 2005-06-09
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	303	 42% 37% 8% 14%
1	B	303	 42% 34% 8% 16%
1	C	303	 38% 43% 7% 11%
1	D	303	 36% 40% 8% 16%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 8601 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase GCN2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	261	Total 2153	C 1378	N 375	O 391	S 9	5	0	0
1	B	254	Total 2091	C 1341	N 364	O 377	S 9	0	0	0
1	C	271	Total 2235	C 1435	N 388	O 403	S 9	0	0	0
1	D	256	Total 2108	C 1352	N 366	O 381	S 9	0	0	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	592	SER	-	CLONING ARTIFACT	UNP P15442
A	593	LEU	-	CLONING ARTIFACT	UNP P15442
B	592	SER	-	CLONING ARTIFACT	UNP P15442
B	593	LEU	-	CLONING ARTIFACT	UNP P15442
C	592	SER	-	CLONING ARTIFACT	UNP P15442
C	593	LEU	-	CLONING ARTIFACT	UNP P15442
D	592	SER	-	CLONING ARTIFACT	UNP P15442
D	593	LEU	-	CLONING ARTIFACT	UNP P15442

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total 3	O 3	0	0
2	B	6	Total 6	O 6	0	0
2	C	5	Total 5	O 5	0	0



• Molecule 1: Serine/threonine-protein kinase GCN2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.89Å 95.70Å 175.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.99 – 3.00 41.99 – 2.81	Depositor EDS
% Data completeness (in resolution range)	94.6 (41.99-3.00) 93.1 (41.99-2.81)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 2.81Å)	Xtrriage
Refinement program	CNX 2000.1	Depositor
R, R_{free}	0.234 , 0.299 0.226 , 0.283	Depositor DCC
R_{free} test set	562 reflections (2.20%)	DCC
Wilson B-factor (Å ²)	81.8	Xtrriage
Anisotropy	0.487	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 68.9	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Outliers	0 of 31126 reflections	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8601	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/2196	0.71	1/2956 (0.0%)
1	B	0.45	0/2132	0.76	4/2869 (0.1%)
1	C	0.46	0/2279	0.71	2/3066 (0.1%)
1	D	0.46	0/2150	0.70	0/2895
All	All	0.46	0/8757	0.72	7/11786 (0.1%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	633	THR	N-CA-C	11.42	141.84	111.00
1	C	898	GLY	N-CA-C	-8.85	90.99	113.10
1	C	820	LEU	CA-CB-CG	-6.17	101.10	115.30
1	B	859	ASN	N-CA-C	5.95	127.07	111.00
1	A	858	LYS	N-CA-C	5.81	126.68	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2153	0	2160	133	0
1	B	2091	0	2104	144	0
1	C	2235	0	2259	158	0
1	D	2108	0	2122	142	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	3	0	0	0	0
2	B	6	0	0	1	0
2	C	5	0	0	0	0
All	All	8601	0	8645	559	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 559 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:924:THR:HG22	1:D:926:MET:H	1.09	1.14
1:B:769:ARG:HB3	1:B:769:ARG:HH11	1.22	1.00
1:B:769:ARG:NH1	1:B:769:ARG:HB3	1.78	0.97
1:D:858:LYS:HD2	1:D:858:LYS:H	1.34	0.92
1:C:657:ARG:HB2	1:C:789:GLU:HB2	1.52	0.92

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	255/303 (84%)	201 (79%)	42 (16%)	12 (5%)	3 17
1	B	246/303 (81%)	191 (78%)	46 (19%)	9 (4%)	4 23
1	C	265/303 (88%)	200 (76%)	51 (19%)	14 (5%)	2 14
1	D	250/303 (82%)	202 (81%)	34 (14%)	14 (6%)	2 13
All	All	1016/1212 (84%)	794 (78%)	173 (17%)	49 (5%)	3 17

5 of 49 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	858	LYS
1	C	770	ASN
1	D	904	GLU
1	A	639	THR
1	A	937	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/272 (86%)	216 (92%)	18 (8%)	16	50
1	B	227/272 (84%)	201 (88%)	26 (12%)	7	28
1	C	243/272 (89%)	220 (90%)	23 (10%)	11	38
1	D	229/272 (84%)	212 (93%)	17 (7%)	17	52
All	All	933/1088 (86%)	849 (91%)	84 (9%)	12	41

5 of 84 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	906	ILE
1	C	621	ASP
1	D	905	LYS
1	B	942	PHE
1	B	973	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	840	ASN
1	C	770	ASN
1	D	833	HIS
1	B	964	HIS
1	B	967	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	261/303 (86%)	-0.17	1 (0%) 93 80	41, 71, 112, 125	2 (0%)
1	B	254/303 (83%)	-0.20	3 (1%) 81 55	40, 78, 116, 132	0
1	C	271/303 (89%)	-0.17	0 100 100	41, 76, 110, 117	0
1	D	256/303 (84%)	-0.26	6 (2%) 64 33	37, 67, 106, 118	0
All	All	1042/1212 (85%)	-0.20	10 (0%) 84 60	37, 73, 111, 132	2 (0%)

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	899	THR	4.3
1	B	886	GLY	3.1
1	D	897	ASP	3.0
1	D	859	ASN	2.7
1	B	897	ASP	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.