



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 07:12 AM GMT

PDB ID : 2ZYA
Title : Dimeric 6-phosphogluconate dehydrogenase complexed with 6-phosphogluconate
Authors : Chen, Y.-Y.; Ko, T.-P.; Lo, L.-P.; Lin, C.-H.; Wang, A.H.-J.
Deposited on : 2009-01-19
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	2475 (1.60-1.60)
Clashscore	102246	2732 (1.60-1.60)
Ramachandran outliers	100387	2654 (1.60-1.60)
Sidechain outliers	100360	2653 (1.60-1.60)
RSRZ outliers	91569	2479 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 8643 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 6-phosphogluconate dehydrogenase, decarboxylating.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	465	Total	C 3602	N 2290	O 606	S 697	9	0	0
1	B	466	Total	C 3610	N 2296	O 607	S 698	9	0	0

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	EXPRESSION TAG	UNP P00350
A	-10	GLY	-	EXPRESSION TAG	UNP P00350
A	-9	HIS	-	EXPRESSION TAG	UNP P00350
A	-8	HIS	-	EXPRESSION TAG	UNP P00350
A	-7	HIS	-	EXPRESSION TAG	UNP P00350
A	-6	HIS	-	EXPRESSION TAG	UNP P00350
A	-5	HIS	-	EXPRESSION TAG	UNP P00350
A	-4	HIS	-	EXPRESSION TAG	UNP P00350
A	-3	HIS	-	EXPRESSION TAG	UNP P00350
A	-2	HIS	-	EXPRESSION TAG	UNP P00350
A	-1	HIS	-	EXPRESSION TAG	UNP P00350
A	0	HIS	-	EXPRESSION TAG	UNP P00350
A	378	CYS	TYR	ENGINEERED	UNP P00350
B	-11	MET	-	EXPRESSION TAG	UNP P00350
B	-10	GLY	-	EXPRESSION TAG	UNP P00350
B	-9	HIS	-	EXPRESSION TAG	UNP P00350
B	-8	HIS	-	EXPRESSION TAG	UNP P00350
B	-7	HIS	-	EXPRESSION TAG	UNP P00350
B	-6	HIS	-	EXPRESSION TAG	UNP P00350
B	-5	HIS	-	EXPRESSION TAG	UNP P00350
B	-4	HIS	-	EXPRESSION TAG	UNP P00350
B	-3	HIS	-	EXPRESSION TAG	UNP P00350
B	-2	HIS	-	EXPRESSION TAG	UNP P00350
B	-1	HIS	-	EXPRESSION TAG	UNP P00350
B	0	HIS	-	EXPRESSION TAG	UNP P00350

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	378	CYS	TYR	ENGINEERED	UNP P00350

- Molecule 2 is SUGAR (6-PHOSPHOGLUCONIC ACID) (three-letter code: 6PG) (formula: C₆H₁₃O₁₀P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O P 17 6 10 1	0	0
2	B	1	Total C O P 17 6 10 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	680	Total O 680 680	0	0
3	B	717	Total O 717 717	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 6-phosphogluconate dehydrogenase, decarboxylating



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.00 Å 108.72 Å 133.50 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.60 31.98 – 1.60	Depositor EDS
% Data completeness (in resolution range)	93.6 (50.00-1.60) 93.6 (31.98-1.60)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	3.68 (at 1.60 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R , R_{free}	0.153 , 0.181 0.153 , 0.180	Depositor DCC
R_{free} test set	6192 reflections (5.05%)	DCC
Wilson B-factor (Å ²)	13.7	Xtriage
Anisotropy	0.281	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 52.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 122724 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8643	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 6PG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.82	1/3670 (0.0%)	0.84	2/4969 (0.0%)
1	B	0.89	1/3678 (0.0%)	0.91	3/4980 (0.1%)
All	All	0.86	2/7348 (0.0%)	0.87	5/9949 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	263	GLY	C-O	10.02	1.39	1.23
1	A	263	GLY	C-O	9.81	1.39	1.23

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	141	MET	CG-SD-CE	-8.35	86.84	100.20
1	A	261	GLY	N-CA-C	6.44	129.19	113.10
1	A	111	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	B	111	ARG	NE-CZ-NH2	-5.14	117.73	120.30
1	B	111	ARG	NE-CZ-NH1	5.12	122.86	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	427	TYR	Sidechain
1	B	353	TYR	Sidechain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3602	0	3569	57	0
1	B	3610	0	3580	48	0
2	A	17	0	10	0	0
2	B	17	0	10	0	0
3	A	680	0	0	20	1
3	B	717	0	0	12	1
All	All	8643	0	7169	100	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (100) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:309:GLN:HG2	1:A:414:ASN:ND2	1.83	0.93
1:A:410:TYR:CZ	1:A:414:ASN:OD1	2.22	0.92
1:A:262:THR:HG22	3:A:947:HOH:O	1.70	0.90
1:A:413:GLN:HE22	1:B:402:GLN:HE22	1.16	0.90
1:A:262:THR:CG2	3:A:947:HOH:O	2.22	0.84
1:B:18:LEU:HD23	1:B:71:LEU:HD21	1.58	0.83
1:B:358:LYS:HD3	3:B:1169:HOH:O	1.82	0.79
1:A:3:LYS:HE2	3:A:472:HOH:O	1.82	0.78
1:B:410:TYR:CZ	1:B:414:ASN:OD1	2.36	0.77
1:A:309:GLN:CG	1:A:414:ASN:ND2	2.47	0.77
1:B:5:GLN:HE22	1:B:28:THR:H	1.30	0.77
1:A:410:TYR:CE1	1:A:414:ASN:OD1	2.38	0.77
1:B:307:GLN:HB3	3:B:874:HOH:O	1.86	0.75
1:A:395:GLN:HG3	3:A:1347:HOH:O	1.86	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3:LYS:HE3	3:B:1263:HOH:O	1.87	0.74
1:B:5:GLN:H	1:B:5:GLN:HE21	1.36	0.73
1:A:355:GLU:HG2	3:A:684:HOH:O	1.89	0.71
1:B:414:ASN:HD22	1:B:414:ASN:N	1.87	0.70
1:B:307:GLN:H	1:B:307:GLN:NE2	1.89	0.70
1:A:309:GLN:HG2	1:A:414:ASN:HD22	1.56	0.68
1:A:401:GLN:NE2	1:A:405:ARG:HH21	1.91	0.68
1:B:316:GLU:OE2	3:B:1265:HOH:O	2.13	0.66
1:A:413:GLN:HE22	1:B:402:GLN:NE2	1.90	0.66
1:B:309:GLN:HB2	1:B:414:ASN:HD21	1.60	0.65
1:B:251:ASP:OD2	3:B:1067:HOH:O	2.15	0.65
1:A:191:TYR:OH	1:A:259:ASN:HB2	1.97	0.64
1:B:108:THR:OG1	1:B:180:HIS:HD2	1.79	0.64
1:B:309:GLN:HG2	1:B:414:ASN:ND2	2.13	0.64
1:A:345:GLU:HG3	3:A:1354:HOH:O	1.97	0.64
1:A:309:GLN:HB2	1:A:414:ASN:HD21	1.63	0.62
1:A:345:GLU:OE2	3:A:814:HOH:O	2.16	0.62
1:A:466:TRP:C	3:A:611:HOH:O	2.38	0.61
1:B:401:GLN:NE2	1:B:405:ARG:HH21	1.99	0.61
1:A:108:THR:OG1	1:A:180:HIS:HD2	1.83	0.60
1:A:58:LYS:HE2	3:A:1304:HOH:O	1.99	0.60
1:B:270:ALA:HB2	1:B:339:GLN:HE22	1.68	0.59
1:A:341:ARG:HH12	1:A:383:GLN:NE2	2.02	0.57
1:A:461:VAL:HG12	3:A:855:HOH:O	2.05	0.56
1:A:413:GLN:NE2	1:B:402:GLN:HE22	1.97	0.56
1:B:213:GLU:HG2	3:B:1068:HOH:O	2.05	0.56
1:A:309:GLN:CG	1:A:414:ASN:HD21	2.18	0.56
1:A:186:HIS:CE1	1:A:364:CYS:HB2	2.41	0.55
1:B:244:GLU:HG2	3:B:1384:HOH:O	2.07	0.55
1:A:380:GLU:HG3	3:A:755:HOH:O	2.06	0.55
1:B:309:GLN:H	1:B:414:ASN:CG	2.10	0.54
1:B:309:GLN:CG	1:B:414:ASN:ND2	2.70	0.54
1:B:5:GLN:NE2	1:B:28:THR:H	2.03	0.54
1:A:102:ASN:HA	3:A:1297:HOH:O	2.07	0.54
1:B:186:HIS:CE1	1:B:364:CYS:HB2	2.43	0.53
1:B:270:ALA:HB2	1:B:339:GLN:NE2	2.25	0.52
1:B:395:GLN:NE2	3:B:734:HOH:O	2.43	0.52
1:A:309:GLN:CB	1:A:414:ASN:HD21	2.21	0.52
1:B:401:GLN:HE21	1:B:405:ARG:HE	1.55	0.52
1:B:2:SER:N	3:B:955:HOH:O	2.45	0.50
1:A:401:GLN:HE21	1:A:405:ARG:HH21	1.60	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:414:ASN:HD22	1:A:414:ASN:N	2.07	0.50
1:B:358:LYS:HG3	1:B:361:ARG:NH2	2.27	0.49
1:B:309:GLN:CB	1:B:414:ASN:HD21	2.24	0.49
1:B:108:THR:OG1	1:B:180:HIS:CD2	2.63	0.49
1:B:126:GLY:HA3	1:B:183:LYS:HB2	1.95	0.49
1:B:307:GLN:H	1:B:307:GLN:CD	2.14	0.49
1:A:414:ASN:ND2	1:A:414:ASN:N	2.61	0.48
1:A:270:ALA:HB2	1:A:339:GLN:HE22	1.78	0.48
1:A:187:ASN:HB3	1:A:262:THR:OG1	2.14	0.48
1:A:136:LYS:HE3	3:A:1157:HOH:O	2.12	0.48
1:A:376:ASP:O	1:A:380:GLU:HG2	2.12	0.48
1:B:191:TYR:OH	1:B:259:ASN:HB2	2.13	0.48
1:B:186:HIS:HD2	1:B:187:ASN:OD1	1.96	0.48
1:A:243:ASP:CG	3:A:1231:HOH:O	2.51	0.48
1:A:358:LYS:HE3	3:A:685:HOH:O	2.14	0.47
1:A:108:THR:OG1	1:A:180:HIS:CD2	2.65	0.47
1:A:38:LYS:HD2	3:A:638:HOH:O	2.16	0.45
1:A:309:GLN:CB	1:A:414:ASN:ND2	2.79	0.45
1:B:395:GLN:HG3	3:B:1019:HOH:O	2.17	0.45
1:B:18:LEU:CD2	1:B:71:LEU:HD21	2.39	0.45
1:A:395:GLN:NE2	3:A:1053:HOH:O	2.50	0.44
1:B:307:GLN:CB	3:B:874:HOH:O	2.57	0.44
1:A:243:ASP:OD1	1:A:244:GLU:N	2.51	0.43
1:A:308:ALA:HA	1:A:414:ASN:HB3	2.00	0.43
1:A:261:GLY:HA2	3:A:1001:HOH:O	2.19	0.43
1:B:5:GLN:H	1:B:5:GLN:NE2	2.12	0.43
1:B:233:ASP:O	1:B:236:LYS:HG3	2.18	0.42
1:A:270:ALA:HB2	1:A:339:GLN:NE2	2.34	0.42
1:A:410:TYR:CE2	1:A:414:ASN:OD1	2.71	0.42
1:B:461:VAL:HG23	3:B:509:HOH:O	2.20	0.42
1:A:461:VAL:CG1	3:A:855:HOH:O	2.64	0.42
1:B:186:HIS:HE1	1:B:366:ILE:H	1.67	0.42
1:A:401:GLN:HE22	1:B:305:GLY:HA2	1.85	0.42
1:A:453:TYR:CD2	1:B:253:ILE:HG12	2.55	0.42
1:B:358:LYS:HG3	1:B:361:ARG:CZ	2.49	0.42
1:A:309:GLN:H	1:A:414:ASN:CG	2.24	0.41
1:B:66:THR:HA	1:B:67:PRO:C	2.41	0.41
1:A:126:GLY:HA3	1:A:183:LYS:HB2	2.01	0.41
1:B:410:TYR:CE2	1:B:414:ASN:OD1	2.74	0.41
1:A:243:ASP:CG	1:A:244:GLU:N	2.73	0.41
1:A:245:ASP:OD1	1:A:247:ASN:HB2	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:164:GLU:OE1	1:A:164:GLU:HA	2.21	0.41
1:A:213:GLU:O	1:A:217:GLN:HG3	2.21	0.41
1:A:186:HIS:HD2	1:A:187:ASN:OD1	2.03	0.41
1:A:345:GLU:CG	3:A:1354:HOH:O	2.64	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:678:HOH:O	3:B:948:HOH:O[2_564]	2.18	0.02

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	463/480 (96%)	451 (97%)	11 (2%)	1 (0%)	52 28
1	B	464/480 (97%)	452 (97%)	11 (2%)	1 (0%)	52 28
All	All	927/960 (97%)	903 (97%)	22 (2%)	2 (0%)	52 28

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	245	ASP
1	B	129	GLY

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	373/387 (96%)	370 (99%)	3 (1%)	86 75
1	B	374/387 (97%)	370 (99%)	4 (1%)	80 63
All	All	747/774 (96%)	740 (99%)	7 (1%)	84 71

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	GLU
1	A	106	GLN
1	A	243	ASP
1	B	5	GLN
1	B	106	GLN
1	B	141	MET
1	B	414	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	106	GLN
1	A	180	HIS
1	A	186	HIS
1	A	217	GLN
1	A	268	GLN
1	A	339	GLN
1	A	383	GLN
1	A	401	GLN
1	A	413	GLN
1	A	414	ASN
1	B	5	GLN
1	B	106	GLN
1	B	112	ASN
1	B	145	GLN
1	B	180	HIS
1	B	186	HIS
1	B	224	ASN
1	B	268	GLN
1	B	307	GLN
1	B	339	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	401	GLN
1	B	414	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	6PG	A	2002	-	12,16,16	1.94	5 (41%)	14,23,23	1.00	0
2	6PG	B	2001	-	12,16,16	1.98	6 (50%)	14,23,23	1.04	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	6PG	A	2002	-	-	0/18/22/22	0/0/0/0
2	6PG	B	2001	-	-	0/18/22/22	0/0/0/0

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2002	6PG	C4-C3	2.07	1.57	1.53
2	B	2001	6PG	O4-C4	2.21	1.48	1.43
2	A	2002	6PG	P-O6	2.33	1.68	1.60
2	B	2001	6PG	P-O3P	2.33	1.63	1.54
2	B	2001	6PG	P-O6	2.37	1.68	1.60
2	B	2001	6PG	O3-C3	2.57	1.49	1.43
2	A	2002	6PG	P-O3P	2.58	1.64	1.54
2	A	2002	6PG	O5-C5	2.59	1.49	1.43
2	B	2001	6PG	P-O2P	2.60	1.64	1.54
2	B	2001	6PG	C6-C5	3.36	1.56	1.51
2	A	2002	6PG	C6-C5	3.89	1.57	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2001	6PG	O3P-P-O6	2.08	112.55	106.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	465/480 (96%)	-0.25	11 (2%) 62 60	8, 16, 30, 52	0
1	B	466/480 (97%)	-0.40	6 (1%) 79 79	7, 13, 24, 46	0
All	All	931/960 (96%)	-0.32	17 (1%) 71 70	7, 15, 28, 52	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	244	GLU	5.3
1	A	2	SER	5.0
1	A	244	GLU	4.5
1	A	243	ASP	4.1
1	B	467	LEU	4.0
1	A	461	VAL	3.9
1	B	461	VAL	3.8
1	A	245	ASP	3.5
1	A	263	GLY	3.2
1	B	245	ASP	2.9
1	B	262	THR	2.7
1	A	246	GLY	2.5
1	A	164	GLU	2.5
1	A	262	THR	2.3
1	A	260	LYS	2.3
1	B	263	GLY	2.2
1	A	459	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	6PG	A	2002	17/17	0.83	0.18	1.25	25,30,35,42	0
2	6PG	B	2001	17/17	0.86	0.14	0.94	19,27,29,29	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.