



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:35 PM GMT

PDB ID : 3ZYC
Title : DYNAMIN 1 GTPASE GED FUSION DIMER COMPLEXED WITH GMP-PCP
Authors : Chappie, J.S.; Mears, J.A.; Fang, S.; Leonard, M.; Schmid, S.L.; Milligan, R.A.; Hinshaw, J.E.; Dyda, F.
Deposited on : 2011-08-22
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	
1	D	353	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5719 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DYNAMIN-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	330	Total	C	N	O	S	0	0	0
			2573	1618	454	491	10			
1	D	340	Total	C	N	O	S	0	0	0
			2667	1678	472	507	10			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	CLONING ARTIFACT	UNP Q05193
A	2	PRO	-	CLONING ARTIFACT	UNP Q05193
A	3	GLU	-	CLONING ARTIFACT	UNP Q05193
A	4	PHE	-	CLONING ARTIFACT	UNP Q05193
A	5	SER	-	CLONING ARTIFACT	UNP Q05193
A	321	LYS	-	LINKER	UNP Q05193
A	322	HIS	-	LINKER	UNP Q05193
A	323	GLY	-	LINKER	UNP Q05193
A	324	THR	-	LINKER	UNP Q05193
A	325	ASP	-	LINKER	UNP Q05193
A	326	SER	-	LINKER	UNP Q05193
A	327	ARG	-	LINKER	UNP Q05193
A	328	VAL	-	LINKER	UNP Q05193
A	744	ASN	ASP	VARIANT	UNP Q05193
D	1	GLY	-	CLONING ARTIFACT	UNP Q05193
D	2	PRO	-	CLONING ARTIFACT	UNP Q05193
D	3	GLU	-	CLONING ARTIFACT	UNP Q05193
D	4	PHE	-	CLONING ARTIFACT	UNP Q05193
D	5	SER	-	CLONING ARTIFACT	UNP Q05193
D	321	LYS	-	LINKER	UNP Q05193
D	322	HIS	-	LINKER	UNP Q05193
D	323	GLY	-	LINKER	UNP Q05193
D	324	THR	-	LINKER	UNP Q05193
D	325	ASP	-	LINKER	UNP Q05193
D	326	SER	-	LINKER	UNP Q05193

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	327	ARG	-	LINKER	UNP Q05193
D	328	VAL	-	LINKER	UNP Q05193
D	744	ASN	ASP	VARIANT	UNP Q05193

- Molecule 2 is PHOSPHOMETHYLPHOSPHONIC ACID GUANYLATE ESTER (three-letter code: GCP) (formula: $C_{11}H_{18}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			32	11	5	13	3		
2	D	1	Total	C	N	O	P	0	0
			32	11	5	13	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	229	Total	O	0	0
			229	229		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	184	Total 184	O 184	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DYNAMIN-1



• Molecule 1: DYNAMIN-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	44.15Å 82.64Å 95.01Å 90.00° 96.68° 90.00°	Depositor
Resolution (Å)	30.00 – 2.20 29.40 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.0 (30.00-2.20) 99.1 (29.40-2.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 2.20Å)	Xtriage
Refinement program	CNS 1.3	Depositor
R, R_{free}	0.225 , 0.269 0.224 , 0.266	Depositor DCC
R_{free} test set	941 reflections (2.75%)	DCC
Wilson B-factor (Å ²)	18.7	Xtriage
Anisotropy	0.331	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 50.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	2 of 34205 reflections (0.006%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5719	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 51.02 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 5.9503e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GCP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.30	0/2610	0.58	0/3532
1	D	0.30	0/2707	0.55	0/3662
All	All	0.30	0/5317	0.56	0/7194

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2573	0	2632	21	0
1	D	2667	0	2723	31	0
2	A	32	0	14	0	0
2	D	32	0	14	0	0
3	A	1	0	0	0	0
3	D	1	0	0	0	0
4	A	229	0	0	2	0
4	D	184	0	0	3	0
All	All	5719	0	5383	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:67:ARG:HH12	1:D:118:VAL:HB	1.52	0.72
1:D:60:GLY:HA2	1:D:239:GLN:HE22	1.60	0.66
1:D:10:ILE:HB	1:D:11:PRO:HD3	1.82	0.62
1:D:71:LEU:HD11	1:D:124:VAL:HG23	1.82	0.61
1:A:75:ASN:HA	1:A:126:SER:O	2.01	0.60
1:A:23:ILE:HG22	1:A:25:GLN:HG2	1.83	0.60
1:A:20:PHE:HE1	1:A:300:LEU:HD11	1.69	0.57
1:D:40:GLN:HA	1:D:139:GLY:HA3	1.86	0.57
1:D:34:ILE:HB	1:D:134:LEU:HD22	1.88	0.56
1:D:79:GLU:HA	1:D:125:TYR:O	2.07	0.55
1:A:15:ARG:HH11	1:A:15:ARG:HG3	1.71	0.55
1:D:38:GLY:HA2	1:D:186:ALA:HB2	1.89	0.55
1:A:274:THR:HB	1:A:275:PRO:HD3	1.88	0.55
1:A:249:THR:HG23	4:A:2194:HOH:O	2.10	0.51
1:D:86:CYS:HB2	4:D:2056:HOH:O	2.10	0.51
1:D:67:ARG:NH1	1:D:118:VAL:HB	2.23	0.51
1:D:17:GLN:HG3	1:D:29:LEU:HD12	1.91	0.51
1:A:40:GLN:HA	1:A:139:GLY:HA3	1.93	0.49
1:D:87:LYS:HZ2	1:D:87:LYS:H	1.61	0.49
1:A:732:TYR:O	1:A:736:LYS:HB2	2.13	0.47
1:D:298:ASN:ND2	4:D:2168:HOH:O	2.47	0.47
1:D:188:LYS:O	1:D:192:GLU:HG3	2.15	0.46
1:D:74:VAL:O	1:D:125:TYR:HA	2.16	0.46
1:D:87:LYS:HD3	1:D:87:LYS:H	1.81	0.46
1:A:240:LYS:HE2	4:A:2185:HOH:O	2.15	0.45
1:D:274:THR:HB	1:D:275:PRO:HD3	1.98	0.45
1:D:87:LYS:HB2	1:D:87:LYS:NZ	2.32	0.45
1:A:6:MET:HE2	1:A:130:LEU:HD21	1.99	0.44
1:A:29:LEU:N	1:A:29:LEU:HD12	2.32	0.44
1:D:73:LEU:HD21	1:D:134:LEU:HG	2.00	0.43
1:D:76:ALA:HB3	1:D:125:TYR:HB3	2.00	0.43
1:D:113:LYS:HD2	1:D:113:LYS:N	2.34	0.43
1:A:309:LYS:HB3	1:A:309:LYS:NZ	2.32	0.43
1:A:67:ARG:HG2	1:A:116:SER:HB3	2.01	0.43
1:D:327:ARG:NE	1:D:327:ARG:HA	2.33	0.43
1:A:87:LYS:HE2	1:A:88:GLY:N	2.33	0.42
1:D:297:ARG:NH2	1:D:746:ASN:HD22	2.18	0.42
1:D:75:ASN:HA	1:D:126:SER:O	2.20	0.42
1:A:143:VAL:HG21	1:D:225:LEU:HD21	2.02	0.41
1:A:149:PRO:HG3	1:A:155:GLN:OE1	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:104:GLU:HG3	4:D:2047:HOH:O	2.20	0.41
1:A:737:GLU:O	1:A:741:ILE:HG12	2.20	0.41
1:A:29:LEU:H	1:A:29:LEU:HD12	1.85	0.41
1:A:728:MET:SD	1:A:730:ARG:HD3	2.61	0.41
1:D:23:ILE:HG22	1:D:25:GLN:HG2	2.02	0.41
1:D:262:HIS:HA	1:D:263:PRO:HD2	1.89	0.41
1:D:87:LYS:HD3	1:D:87:LYS:N	2.36	0.40
1:A:307:ILE:N	1:A:307:ILE:HD12	2.36	0.40
1:D:282:ASN:HA	1:D:282:ASN:HD22	1.71	0.40
1:D:135:VAL:HG12	1:D:137:LEU:HG	2.04	0.40
1:A:134:LEU:HD12	1:A:134:LEU:N	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/353 (92%)	320 (98%)	6 (2%)	0	100	100
1	D	336/353 (95%)	319 (95%)	15 (4%)	2 (1%)	30	29
All	All	662/706 (94%)	639 (96%)	21 (3%)	2 (0%)	46	50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	30	ASP
1	D	79	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/309 (93%)	281 (98%)	7 (2%)	57	69
1	D	298/309 (96%)	292 (98%)	6 (2%)	63	76
All	All	586/618 (95%)	573 (98%)	13 (2%)	60	72

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	54	ARG
1	A	69	LEU
1	A	87	LYS
1	A	240	LYS
1	A	299	LYS
1	A	301	GLN
1	A	727	GLU
1	D	78	THR
1	D	87	LYS
1	D	99	LEU
1	D	113	LYS
1	D	248	ILE
1	D	304	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	162	GLN
1	A	168	ASN
1	A	198	GLN
1	A	301	GLN
1	A	746	ASN
1	D	25	GLN
1	D	33	GLN
1	D	72	GLN
1	D	148	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	155	GLN
1	D	162	GLN
1	D	239	GLN
1	D	282	ASN
1	D	303	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GCP	A	1749	3	26,34,34	1.40	4 (15%)	34,54,54	2.51	8 (23%)
2	GCP	D	1749	3	26,34,34	1.34	4 (15%)	34,54,54	2.51	8 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GCP	A	1749	3	-	0/15/38/38	0/3/3/3
2	GCP	D	1749	3	-	0/15/38/38	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1749	GCP	PB-O2B	-3.21	1.48	1.56
2	D	1749	GCP	PB-O2B	-2.98	1.49	1.56
2	D	1749	GCP	PG-O2G	-2.59	1.48	1.54
2	A	1749	GCP	PG-O3G	-2.56	1.48	1.54
2	A	1749	GCP	PG-O2G	-2.53	1.48	1.54
2	D	1749	GCP	PG-O3G	-2.24	1.49	1.54
2	D	1749	GCP	C6-N1	3.40	1.39	1.33
2	A	1749	GCP	C6-N1	3.47	1.39	1.33

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1749	GCP	C5-C6-N1	-8.87	111.46	123.59
2	A	1749	GCP	C5-C6-N1	-8.80	111.56	123.59
2	A	1749	GCP	PA-O3A-PB	-6.49	114.50	132.73
2	D	1749	GCP	PA-O3A-PB	-6.47	114.56	132.73
2	A	1749	GCP	O1G-PG-C3B	-2.75	104.86	111.13
2	D	1749	GCP	N3-C2-N1	-2.28	123.97	127.44
2	A	1749	GCP	N3-C2-N1	-2.27	123.99	127.44
2	D	1749	GCP	O1G-PG-C3B	-2.26	105.98	111.13
2	D	1749	GCP	O2G-PG-O1G	2.31	118.30	112.40
2	D	1749	GCP	O3G-PG-O1G	2.37	118.47	112.40
2	A	1749	GCP	O2G-PG-O1G	2.39	118.53	112.40
2	A	1749	GCP	O3G-PG-O1G	2.42	118.58	112.40
2	A	1749	GCP	O2B-PB-O1B	3.07	119.77	110.12
2	D	1749	GCP	O2B-PB-O1B	3.11	119.91	110.12
2	A	1749	GCP	C6-N1-C2	6.58	125.08	115.94
2	D	1749	GCP	C6-N1-C2	6.60	125.10	115.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	330/353 (93%)	0.21	20 (6%) 25 24	4, 19, 51, 79	0
1	D	340/353 (96%)	0.26	18 (5%) 30 29	5, 23, 54, 72	0
All	All	670/706 (94%)	0.23	38 (5%) 27 27	4, 21, 53, 79	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	726	ASP	6.6
1	A	24	GLY	6.2
1	A	309	LYS	5.3
1	A	78	THR	5.2
1	A	26	ASN	5.1
1	D	26	ASN	4.9
1	D	328	VAL	4.6
1	A	729	LEU	4.5
1	D	748	THR	4.3
1	A	307	ILE	4.0
1	A	127	PRO	3.9
1	D	27	ALA	3.9
1	D	25	GLN	3.8
1	A	25	GLN	3.3
1	D	23	ILE	3.1
1	D	29	LEU	3.1
1	D	314	TYR	3.0
1	A	728	MET	3.0
1	D	78	THR	3.0
1	D	305	LEU	2.9
1	A	5	SER	2.8
1	A	23	ILE	2.8
1	D	128	HIS	2.7
1	A	128	HIS	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	327	ARG	2.6
1	A	77	THR	2.5
1	A	310	GLU	2.5
1	A	8	ASP	2.4
1	A	730	ARG	2.4
1	A	306	SER	2.3
1	D	28	ASP	2.3
1	D	79	GLU	2.2
1	A	298	ASN	2.1
1	A	311	VAL	2.1
1	D	174	VAL	2.1
1	D	317	PHE	2.1
1	D	306	SER	2.1
1	D	24	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	GCP	D	1749	32/32	0.98	0.11	-0.48	3,8,12,14	0
2	GCP	A	1749	32/32	0.98	0.11	-0.70	5,8,12,13	0
3	MG	D	1750	1/1	0.83	0.09	-1.60	9,9,9,9	0
3	MG	A	1750	1/1	0.98	0.04	-3.96	5,5,5,5	0

6.5 Other polymers [i](#)

There are no such residues in this entry.