



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:12 AM GMT

PDB ID : 2ZZL  
Title : Structure of bacteriorhodopsin's M intermediate at pH 7  
Authors : Yamamoto, M.; Kouyama, T.  
Deposited on : 2009-02-18  
Resolution : 2.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*


The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	7858 (2.04-2.00)
Clashscore	102246	9060 (2.04-2.00)
Ramachandran outliers	100387	8952 (2.04-2.00)
Sidechain outliers	100360	8951 (2.04-2.00)
RSRZ outliers	91569	7873 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	262	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	L3P	A	260	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	L2P	A	270	-	-	-	X
5	L1P	A	280	-	-	-	X
5	L1P	A	290	-	-	-	X
5	L1P	A	300	-	-	-	X
7	SOG	A	430	-	-	-	X

## 2 Entry composition [i](#)


There are 8 unique types of molecules in this entry. The entry contains 2127 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bacteriorhodopsin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	228	Total	C	N	O	S	0	0	0
			1765	1185	269	302	9			

- Molecule 2 is RETINAL (three-letter code: RET) (formula: C<sub>20</sub>H<sub>28</sub>O).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	C	0	0
			20	20		

- Molecule 3 is 2,3-DI-O-PHYTANLY-3-SN-GLYCERO-1-PHOSPHORYL-3'-SN-GLYCEROL-1'-PHOSPHATE (three-letter code: L3P) (formula: C<sub>46</sub>H<sub>94</sub>O<sub>11</sub>P<sub>2</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			59	46	11	2		

- Molecule 4 is 2,3-DI-PHYTANYL-GLYCEROL (three-letter code: L2P) (formula:  $C_{43}H_{88}O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			46	43	3		

- Molecule 5 is 3-PHOSPHORYL-[1,2-DI-PHYTANYL]GLYCEROL (three-letter code: L1P) (formula:  $C_{43}H_{89}O_6P$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O	P	0	0
			40	33	6	1		
5	A	1	Total	C	O	P	0	0
			50	43	6	1		
5	A	1	Total	C	O	P	0	0
			50	43	6	1		

- Molecule 6 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	3	Total	C	O	S	0	0
			37	18	18	1		

- Molecule 7 is SUGAR (2-HYDROXYMETHYL-6-OCTYLSULFANYL-TETRAHYDRO-PYRAN-3,4,5-TRIOL) (three-letter code: SOG) (formula: C<sub>14</sub>H<sub>28</sub>O<sub>5</sub>S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	O	S	0	0
			20	14	5	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	40	Total	O	0	0
			40	40		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Bacteriorhodopsin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 6 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.21Å 102.21Å 112.48Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	15.00 – 2.03 41.18 – 2.03	Depositor EDS
% Data completeness (in resolution range)	88.3 (15.00-2.03) 93.8 (41.18-2.03)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.33 (at 2.03Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.207 , 0.238 0.244 , 0.231	Depositor DCC
$R_{free}$ test set	1050 reflections (4.87%)	DCC
Wilson B-factor (Å <sup>2</sup> )	32.9	Xtriage
Anisotropy	0.259	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 73.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 21618 reflections	Xtriage
$F_o, F_c$ correlation	0.85	EDS
Total number of atoms	2127	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.79% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SOG, GLC, L1P, RET, L2P, L3P, SGA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/1813	0.53	0/2476

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1765	0	1819	45	0
2	A	20	0	27	2	0
3	A	59	0	93	5	0
4	A	46	0	87	1	0
5	A	140	0	238	0	0
6	A	37	0	27	3	0
7	A	20	0	28	3	0
8	A	40	0	0	1	0
All	All	2127	0	2319	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:LYS:HG3	7:A:430:SOG:H62	1.67	0.77
1:A:101:VAL:HG22	1:A:159:LYS:HB3	1.69	0.74
1:A:7:ARG:H	1:A:8:PRO:HD2	1.53	0.71
1:A:101:VAL:CG2	1:A:159:LYS:HB3	2.21	0.71
1:A:6:GLY:HA2	1:A:9:GLU:OE1	1.95	0.67
1:A:129:LYS:H	7:A:430:SOG:H5	1.62	0.65
1:A:40:LYS:NZ	3:A:260:L3P:H12	2.12	0.64
1:A:40:LYS:HZ3	3:A:260:L3P:H12	1.63	0.63
1:A:104:ASP:HB2	1:A:107:THR:HG22	1.82	0.61
1:A:199:VAL:HG12	1:A:203:ILE:HB	1.84	0.58
1:A:66:LEU:HG	1:A:79:TYR:CE1	2.42	0.55
1:A:206:LEU:HD13	1:A:206:LEU:C	2.27	0.55
1:A:44:ALA:HA	3:A:260:L3P:H111	1.88	0.54
1:A:101:VAL:HG21	1:A:156:PHE:O	2.08	0.53
1:A:82:ARG:HG2	8:A:604:HOH:O	2.09	0.53
1:A:186:PRO:HB3	2:A:250:RET:H183	1.91	0.53
1:A:104:ASP:O	1:A:107:THR:HG23	2.09	0.52
1:A:203:ILE:O	1:A:207:LEU:HG	2.10	0.52
1:A:15:LEU:HB3	1:A:209:MET:HE2	1.92	0.52
1:A:12:TRP:CE3	1:A:206:LEU:HD23	2.46	0.51
1:A:40:LYS:HE2	3:A:260:L3P:H31	1.95	0.48
1:A:128:THR:HG23	7:A:430:SOG:O4	2.15	0.47
1:A:196:ALA:O	1:A:198:ILE:N	2.48	0.46
1:A:156:PHE:HB3	1:A:171:PHE:CZ	2.52	0.45
1:A:15:LEU:HD23	1:A:209:MET:CE	2.45	0.45
1:A:53:ALA:HB2	1:A:216:LYS:CD	2.46	0.45
1:A:101:VAL:HG22	1:A:159:LYS:CB	2.44	0.45
2:A:250:RET:H7	2:A:250:RET:H181	1.85	0.45
1:A:101:VAL:HG21	1:A:159:LYS:HB3	1.98	0.44
1:A:129:LYS:HE2	6:A:420:SGA:H61	2.00	0.44
1:A:199:VAL:CG1	1:A:203:ILE:HB	2.47	0.44
1:A:53:ALA:HB2	1:A:216:LYS:HD3	2.00	0.43
1:A:70:PRO:HA	1:A:74:GLU:O	2.19	0.43
1:A:49:VAL:HG21	1:A:93:LEU:CD1	2.50	0.42
1:A:101:VAL:HG23	1:A:160:ALA:CA	2.50	0.42
1:A:206:LEU:O	1:A:210:VAL:HG23	2.20	0.42
1:A:168:ALA:O	1:A:172:LYS:HG2	2.20	0.42
1:A:129:LYS:HZ1	6:A:410:MAN:H61	1.84	0.42
1:A:101:VAL:CG2	1:A:160:ALA:N	2.82	0.42
1:A:120:GLY:HA3	4:A:270:L2P:H172	2.02	0.42
1:A:46:THR:O	1:A:50:PRO:HD2	2.19	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:95:LEU:O	1:A:99:LEU:HG	2.21	0.41
1:A:129:LYS:NZ	6:A:410:MAN:H61	2.35	0.41
1:A:48:LEU:HG	3:A:260:L3P:H143	2.02	0.41
1:A:104:ASP:O	1:A:107:THR:CG2	2.69	0.41
1:A:97:LEU:HD13	1:A:219:PHE:HZ	1.86	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	226/262 (86%)	220 (97%)	4 (2%)	2 (1%)	21 12

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	7	ARG
1	A	197	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	183/206 (89%)	173 (94%)	10 (6%)	27 20

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	ARG
1	A	66	LEU
1	A	93	LEU
1	A	94	LEU
1	A	95	LEU
1	A	97	LEU
1	A	101	VAL
1	A	107	THR
1	A	161	GLU
1	A	211	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	105	GLN
1	A	202	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

3 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	GLC	A	400	6	11,11,12	2.87	5 (45%)	14,15,17	1.14	1 (7%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	MAN	A	410	6	11,11,12	2.18	4 (36%)	14,15,17	1.84	4 (28%)
6	SGA	A	420	6	15,15,16	1.14	0	18,22,24	1.07	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GLC	A	400	6	-	0/2/19/22	0/1/1/1
6	MAN	A	410	6	-	0/2/19/22	0/1/1/1
6	SGA	A	420	6	-	0/7/24/27	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	400	GLC	C4-C5	2.34	1.58	1.53
6	A	410	MAN	C1-C2	2.43	1.58	1.52
6	A	410	MAN	O5-C1	2.80	1.48	1.43
6	A	410	MAN	O2-C2	3.05	1.50	1.43
6	A	400	GLC	C2-C3	4.13	1.58	1.52
6	A	410	MAN	C2-C3	4.32	1.58	1.52
6	A	400	GLC	C1-C2	4.35	1.62	1.52
6	A	400	GLC	O5-C5	4.80	1.54	1.43
6	A	400	GLC	O5-C1	4.81	1.51	1.43

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	410	MAN	O5-C1-C2	-2.95	106.07	110.86
6	A	420	SGA	C3-O3-S	-2.68	113.66	118.77
6	A	410	MAN	C2-C3-C4	-2.53	106.75	111.04
6	A	400	GLC	C2-C3-C4	-2.19	107.31	111.04
6	A	420	SGA	C1-O5-C5	2.03	114.83	112.25
6	A	410	MAN	O6-C6-C5	2.99	121.20	111.33
6	A	410	MAN	C1-O5-C5	4.32	117.73	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	410	MAN	2	0
6	A	420	SGA	1	0

## 5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	RET	A	250	-	19,20,21	1.59	4 (21%)	27,27,28	2.09	11 (40%)
3	L3P	A	260	-	58,58,58	1.18	7 (12%)	64,73,73	1.23	6 (9%)
4	L2P	A	270	-	45,45,45	1.81	12 (26%)	50,53,53	1.51	8 (16%)
5	L1P	A	280	-	39,39,49	1.92	10 (25%)	43,48,60	1.02	2 (4%)
5	L1P	A	290	-	49,49,49	1.74	15 (30%)	55,60,60	1.45	7 (12%)
5	L1P	A	300	-	49,49,49	1.63	11 (22%)	55,60,60	1.13	6 (10%)
7	SOG	A	430	-	20,20,20	1.39	4 (20%)	23,25,25	1.29	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RET	A	250	-	-	0/13/30/31	0/1/1/1
3	L3P	A	260	-	-	0/67/67/67	0/0/0/0
4	L2P	A	270	-	-	0/51/51/51	0/0/0/0
5	L1P	A	280	-	-	0/43/43/55	0/0/0/0
5	L1P	A	290	-	-	0/55/55/55	0/0/0/0
5	L1P	A	300	-	-	0/55/55/55	0/0/0/0
7	SOG	A	430	-	-	0/11/31/31	0/1/1/1

All (63) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	250	RET	C2-C3	-3.10	1.44	1.52
3	A	260	L3P	P1-O2P	-2.12	1.45	1.54
7	A	430	SOG	C1'-S1	-2.05	1.78	1.81
5	A	290	L1P	C54-C53	2.02	1.59	1.52
2	A	250	RET	C7-C6	2.07	1.53	1.45
3	A	260	L3P	C6-C5	2.09	1.59	1.51
5	A	280	L1P	C52-C53	2.14	1.63	1.52
5	A	290	L1P	C26-C25	2.15	1.62	1.52
5	A	300	L1P	C14-C13	2.15	1.60	1.52
5	A	290	L1P	C56-C55	2.20	1.62	1.52
3	A	260	L3P	C42-C41	2.21	1.58	1.50
5	A	280	L1P	C42-C41	2.22	1.58	1.50
5	A	300	L1P	C26-C27	2.25	1.62	1.52
5	A	300	L1P	C19-C18	2.26	1.60	1.52
5	A	280	L1P	O1-C1	2.26	1.48	1.42
5	A	300	L1P	C55-C53	2.28	1.64	1.52
5	A	290	L1P	C19-C18	2.29	1.60	1.52
4	A	270	L2P	C25-C23	2.29	1.64	1.52
5	A	280	L1P	P-O3	2.30	1.68	1.60
5	A	290	L1P	C26-C27	2.31	1.62	1.52
5	A	290	L1P	C14-C13	2.34	1.60	1.52
5	A	280	L1P	C56-C55	2.34	1.62	1.52
4	A	270	L2P	C24-C23	2.37	1.60	1.52
5	A	300	L1P	O1-C11	2.38	1.52	1.42
5	A	300	L1P	C17-C18	2.38	1.65	1.52
5	A	290	L1P	C17-C18	2.41	1.65	1.52
4	A	270	L2P	C56-C57	2.43	1.63	1.52
5	A	290	L1P	C25-C23	2.50	1.65	1.52
5	A	290	L1P	C55-C53	2.51	1.65	1.52
3	A	260	L3P	P2-O6	2.52	1.67	1.60
4	A	270	L2P	C50-C48	2.52	1.65	1.52
5	A	290	L1P	O1-C11	2.54	1.52	1.42
5	A	300	L1P	C15-C13	2.59	1.66	1.52
3	A	260	L3P	P1-O3	2.60	1.70	1.59
5	A	290	L1P	C15-C13	2.67	1.66	1.52
4	A	270	L2P	O2-C2	2.70	1.50	1.43
4	A	270	L2P	O2-C41	2.77	1.50	1.42
4	A	270	L2P	C22-C23	2.78	1.67	1.52
7	A	430	SOG	C4-C5	2.81	1.59	1.53
3	A	260	L3P	O2-C2	2.82	1.51	1.43
5	A	280	L1P	O2-C41	2.84	1.50	1.42
4	A	270	L2P	C47-C48	2.86	1.67	1.52
4	A	270	L2P	C42-C41	2.87	1.61	1.50

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	290	L1P	C12-C11	2.89	1.61	1.50
5	A	300	L1P	C12-C11	2.91	1.61	1.50
7	A	430	SOG	O5-C5	2.95	1.51	1.44
3	A	260	L3P	O2-C41	2.99	1.51	1.42
4	A	270	L2P	C3-C2	3.10	1.59	1.50
5	A	300	L1P	P-O3P	3.16	1.66	1.54
5	A	290	L1P	P-O3P	3.17	1.66	1.54
5	A	300	L1P	C3-C2	3.25	1.59	1.50
4	A	270	L2P	C1-C2	3.32	1.60	1.50
2	A	250	RET	C5-C6	3.32	1.39	1.34
5	A	290	L1P	C3-C2	3.39	1.60	1.50
7	A	430	SOG	O5-C1	3.40	1.48	1.42
2	A	250	RET	C1-C6	3.65	1.59	1.53
5	A	280	L1P	C1-C2	3.68	1.61	1.50
4	A	270	L2P	C26-C25	3.87	1.70	1.52
5	A	300	L1P	P-O3	4.00	1.73	1.60
5	A	280	L1P	C3-C2	4.02	1.62	1.50
5	A	290	L1P	P-O3	4.15	1.74	1.60
5	A	280	L1P	P-O3P	4.36	1.70	1.54
5	A	280	L1P	O2-C2	5.20	1.57	1.43

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	250	RET	C20-C13-C14	-3.69	112.05	123.36
4	A	270	L2P	C22-C21-C20	-3.51	95.57	112.99
4	A	270	L2P	C24-C23-C22	-3.47	97.75	111.08
7	A	430	SOG	C3'-C2'-C1'	-3.24	99.77	113.18
4	A	270	L2P	C16-C17-C18	-3.20	104.86	115.49
2	A	250	RET	C8-C9-C10	-2.95	114.23	118.98
2	A	250	RET	C17-C1-C2	-2.64	99.32	108.79
3	A	260	L3P	C26-C25-C23	-2.59	106.90	115.49
3	A	260	L3P	C46-C45-C43	-2.50	107.18	115.49
3	A	260	L3P	C51-C50-C48	-2.49	107.24	115.49
5	A	300	L1P	C47-C46-C45	-2.38	101.20	112.99
5	A	290	L1P	C47-C46-C45	-2.32	101.48	112.99
4	A	270	L2P	C26-C25-C23	-2.28	107.94	115.49
2	A	250	RET	C1-C6-C5	-2.20	119.42	122.66
4	A	270	L2P	C54-C53-C52	-2.19	102.66	111.08
5	A	290	L1P	O1-C11-C12	2.08	115.57	109.52
2	A	250	RET	C2-C3-C4	2.08	116.78	111.53
5	A	300	L1P	C21-C20-C18	2.10	122.44	115.49

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	260	L3P	O6P-P2-O6	2.14	110.58	105.35
5	A	290	L1P	C16-C17-C18	2.21	122.83	115.49
5	A	300	L1P	O3-P-O1P	2.22	112.79	107.14
5	A	290	L1P	C21-C20-C18	2.33	123.21	115.49
5	A	290	L1P	O3-P-O1P	2.33	113.08	107.14
5	A	300	L1P	C16-C17-C18	2.37	123.34	115.49
2	A	250	RET	C1-C6-C7	2.46	122.70	115.82
5	A	280	L1P	O1-C1-C2	2.50	112.92	108.71
4	A	270	L2P	O1-C1-C2	2.54	113.00	108.71
2	A	250	RET	C16-C1-C6	2.55	114.31	110.30
5	A	300	L1P	O3P-P-O3	2.60	114.06	106.56
5	A	290	L1P	O3P-P-O3	2.76	114.51	106.56
2	A	250	RET	C17-C1-C6	2.95	114.93	110.30
2	A	250	RET	C20-C13-C12	2.98	123.05	118.10
2	A	250	RET	C3-C4-C5	3.22	118.98	113.87
3	A	260	L3P	C11-O1-C1	3.42	124.59	113.44
4	A	270	L2P	C21-C20-C18	3.63	127.51	115.49
5	A	280	L1P	C51-C50-C48	3.69	127.71	115.49
7	A	430	SOG	C1'-S1-C1	3.86	105.62	100.30
2	A	250	RET	C2-C1-C6	4.00	116.70	110.36
5	A	300	L1P	O1-C1-C2	4.09	115.61	108.71
4	A	270	L2P	C21-C22-C23	4.14	129.22	115.49
3	A	260	L3P	C41-O2-C2	4.98	126.86	115.50
5	A	290	L1P	O1-C1-C2	7.67	121.65	108.71

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	250	RET	2	0
3	A	260	L3P	5	0
4	A	270	L2P	1	0
7	A	430	SOG	3	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	228/262 (87%)	1.05	33 (14%) 3 4	22, 31, 51, 71	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	5	THR	11.8
1	A	196	ALA	4.3
1	A	230	PHE	4.0
1	A	198	ILE	3.7
1	A	87	LEU	3.4
1	A	232	GLU	3.2
1	A	231	GLY	3.1
1	A	36	ASP	3.1
1	A	37	PRO	3.0
1	A	7	ARG	3.0
1	A	84	ALA	3.0
1	A	88	PHE	2.9
1	A	227	ARG	2.9
1	A	10	TRP	2.9
1	A	92	LEU	2.8
1	A	119	ILE	2.8
1	A	140	ILE	2.6
1	A	161	GLU	2.5
1	A	123	LEU	2.4
1	A	199	VAL	2.4
1	A	225	ARG	2.3
1	A	32	MET	2.2
1	A	117	ILE	2.2
1	A	38	ASP	2.2
1	A	113	GLY	2.2
1	A	154	PHE	2.2
1	A	26	TYR	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	120	GLY	2.1
1	A	94	LEU	2.1
1	A	80	TRP	2.1
1	A	143	ALA	2.0
1	A	83	TYR	2.0
1	A	6	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	MAN	A	410	11/12	0.77	0.20	1.56	38,43,52,60	0
6	GLC	A	400	11/12	0.91	0.17	0.27	41,44,45,47	0
6	SGA	A	420	15/16	0.67	0.21	-	65,69,71,71	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
7	SOG	A	430	20/20	0.47	0.60	32.97	86,88,89,89	0
4	L2P	A	270	46/46	0.72	0.71	14.04	45,73,85,86	0
3	L3P	A	260	59/59	0.40	0.54	8.67	73,85,94,94	13
5	L1P	A	300	50/50	0.61	0.48	6.38	0,0,0,0	0
5	L1P	A	290	50/50	0.44	0.58	5.87	0,0,0,0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	L1P	A	280	40/50	0.64	0.47	5.54	0,0,0,0	0
2	RET	A	250	20/21	0.88	0.21	1.14	24,27,31,31	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.