



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 20, 2016 – 12:58 PM EDT

PDB ID : 4ZZA
Title : Raffinose and panose binding protein from Bifidobacterium animalis subsp. lactis Bl-04, bound with raffinose, selenomethionine derivative
Authors : Fredslund, F.; Ejby, M.; Andersen, J.M.; Slotboom, D.J.; Abou Hachem, M.
Deposited on : 2015-05-22
Resolution : 2.02 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027790
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027790


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	7858 (2.04-2.00)
Clashscore	102246	9060 (2.04-2.00)
Ramachandran outliers	100387	8952 (2.04-2.00)
Sidechain outliers	100360	8951 (2.04-2.00)
RSRZ outliers	91569	7873 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	396	 87% 5% 8%
1	B	396	 88% 5% 7%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 12256 atoms, of which 5568 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Sugar binding protein of ABC transporter system.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	364	Total	C	H	N	O	S	Se	0	0	0
			5603	1830	2737	472	554	1	9			
1	B	368	Total	C	H	N	O	S	Se	0	0	0
			5669	1853	2768	477	561	1	9			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	43	SER	ASN	conflict	UNP C6A9Y6
A	44	HIS	SER	conflict	UNP C6A9Y6
A	45	MSE	ALA	conflict	UNP C6A9Y6
B	-2	SER	ASN	conflict	UNP C6A9Y6
B	-1	HIS	SER	conflict	UNP C6A9Y6
B	0	MSE	ALA	conflict	UNP C6A9Y6

- Molecule 2 is raffinose (three-letter code: RAF) (formula: C₁₈H₃₂O₁₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			66	18	32	16		
2	B	1	Total	C	H	O	0	0
			65	18	31	16		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	413	Total	O	0	0
			413	413		
3	B	440	Total	O	0	0
			440	440		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sugar binding protein of ABC transporter system



- Molecule 1: Sugar binding protein of ABC transporter system



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.38Å 90.95Å 145.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.00 – 2.02 47.30 – 2.02	Depositor EDS
% Data completeness (in resolution range)	99.7 (43.00-2.02) 99.7 (47.30-2.02)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.43 (at 2.01Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.169 , 0.210 0.165 , 0.209	Depositor DCC
R_{free} test set	2447 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	14.4	Xtriage
Anisotropy	0.520	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 50.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12256	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 55.66 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.0643e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: RAF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/2924	0.63	0/3950
1	B	0.58	0/2960	0.63	0/3999
All	All	0.57	0/5884	0.63	0/7949

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2866	2737	2780	16	1
1	B	2901	2768	2810	12	0
2	A	34	32	31	0	0
2	B	34	31	32	0	0
3	A	413	0	0	11	3
3	B	440	0	0	8	4
All	All	6688	5568	5653	28	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:52:PHE:N	3:B:503:HOH:O	2.03	0.91
1:A:50:VAL:N	3:A:1002:HOH:O	2.07	0.88
1:B:222:LYS:NZ	3:B:501:HOH:O	1.91	0.85
1:B:213:GLU:OE2	3:B:502:HOH:O	1.97	0.83
1:A:266:GLU:OE2	3:A:1001:HOH:O	2.00	0.80
1:A:196:MSE:HE3	1:A:200:PHE:CE2	2.24	0.72
1:A:196:MSE:HE3	1:A:200:PHE:HE2	1.58	0.69
1:A:385:GLU:OE2	3:A:1003:HOH:O	2.12	0.67
1:B:311:ASN:OD1	3:B:505:HOH:O	2.14	0.65
1:A:134:GLU:HB2	3:A:1091:HOH:O	1.98	0.64
1:B:196:GLY:O	3:B:506:HOH:O	2.16	0.62
1:B:133:LYS:NZ	3:B:508:HOH:O	2.25	0.58
1:A:391:ASP:HB2	3:A:1035:HOH:O	2.03	0.58
1:A:385:GLU:OE1	3:A:1004:HOH:O	2.17	0.57
1:B:343:ARG:NH1	3:B:513:HOH:O	2.36	0.53
1:B:270:ASP:OD1	3:B:504:HOH:O	2.19	0.53
1:A:198:LYS:HD2	3:A:1015:HOH:O	2.09	0.52
1:A:426:GLU:OE2	3:A:1005:HOH:O	2.19	0.50
1:A:84:ASN:C	3:A:1237:HOH:O	2.50	0.48
1:B:99:LYS:O	1:B:110:LYS:HG3	2.13	0.48
1:A:220:ALA:HB3	1:A:221:PRO:HD3	1.96	0.46
1:A:50:VAL:HA	3:A:1002:HOH:O	2.15	0.46
1:A:196:MSE:HE3	1:A:200:PHE:CD2	2.52	0.45
1:B:324:GLU:HG2	1:B:324:GLU:O	2.18	0.44
1:B:175:ALA:HB3	1:B:176:PRO:HD3	2.01	0.42
1:A:50:VAL:CA	3:A:1002:HOH:O	2.59	0.42
1:A:144:LYS:O	1:A:155:LYS:HG3	2.19	0.41
1:B:333:GLU:N	1:B:334:PRO:CD	2.84	0.41

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1368:HOH:O	3:B:719:HOH:O[3_655]	2.06	0.14
3:A:1186:HOH:O	3:B:504:HOH:O[3_555]	2.07	0.13
3:A:1027:HOH:O	3:B:504:HOH:O[3_555]	2.11	0.09
1:A:53:ASP:OD1	3:B:504:HOH:O[3_555]	2.12	0.08

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	360/396 (91%)	354 (98%)	6 (2%)	0	100	100
1	B	364/396 (92%)	358 (98%)	6 (2%)	0	100	100
All	All	724/792 (91%)	712 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	305/321 (95%)	301 (99%)	4 (1%)	76	78
1	B	309/321 (96%)	305 (99%)	4 (1%)	76	78
All	All	614/642 (96%)	606 (99%)	8 (1%)	76	78

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	54	PHE
1	A	56	GLN
1	A	108	PHE
1	A	387	ASN
1	B	9	PHE
1	B	11	GLN
1	B	63	PHE
1	B	154	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	219	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	RAF	A	901	-	36,36,36	1.25	3 (8%)	54,54,54	1.06	4 (7%)
2	RAF	B	401	-	36,36,36	1.11	4 (11%)	54,54,54	0.86	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RAF	A	901	-	-	0/17/76/76	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RAF	B	401	-	-	0/17/76/76	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	RAF	C4'-C5'	-3.04	1.44	1.53
2	A	901	RAF	O3-C3	-2.73	1.36	1.43
2	B	401	RAF	C4'-C5'	-2.15	1.47	1.53
2	B	401	RAF	O2X-C2X	-2.14	1.37	1.43
2	A	901	RAF	C4-C5	-2.10	1.48	1.53
2	B	401	RAF	C4-C5	-2.04	1.48	1.53
2	B	401	RAF	O5X-C5X	-2.02	1.39	1.44

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	RAF	C6-C5-C4	-2.22	106.94	112.00
2	A	901	RAF	C6-C5-C4	-2.11	107.18	112.00
2	A	901	RAF	O6-C6-C5	2.01	112.72	109.14
2	A	901	RAF	O5-C5-C4	2.78	114.97	109.67
2	A	901	RAF	O6-C1X-C2X	2.83	111.48	108.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	355/396 (89%)	-0.39	0 100 100	6, 13, 26, 35	0
1	B	359/396 (90%)	-0.44	0 100 100	5, 12, 25, 37	0
All	All	714/792 (90%)	-0.41	0 100 100	5, 13, 25, 37	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	RAF	A	901	34/34	0.97	0.09	0.18	4,10,17,21	0
2	RAF	B	401	34/34	0.98	0.08	-0.36	5,8,14,19	0

6.5 Other polymers ⓘ

There are no such residues in this entry.